



Spatio-Temporal Assessment of Groundwater Quality in the Town of Moundou in South-Western Chad

Doumtoudjinodji Prosper^{1,2,†}, Elegbede Manou Bernadin¹, Doumnang Mbaigane Jean Claude², Socohou Amadou Akilou¹ and Djoueingue Nguérassem²

¹University of Abomey-Calavi (UAC), National Institute of Water (INE), Laboratory of Water and Environmental Science and Technology (LSTEE), Cotonou, Benin

²University of N'Djaména, Faculty of Exact and Applied Sciences (FSEA), Department of Geology, Research Laboratory for Hydro-Geosciences and Reservoirs (LHGR), N'Djaména, Chad

†Corresponding author: Doumtoudjinodji Prosper; doumto@gamil.com

Abbreviation: Nat. Env. & Poll. Technol.

Website: www.neptjournal.com

Received: 25-04-2025

Revised: 09-06-2025

Accepted: 10-06-2025

Key Words:

Groundwater
Drinking water quality
Irrigation water quality
Hydrochemistry

Citation for the Paper:

Doumtoudjinodji, P., Elegbede, M.B., Doumnang, M.J.C., Socohou, A.A. and Djoueingue, N., 2026. Spatio-temporal assessment of groundwater quality in the town of Moundou in south-western Chad. *Nature Environment and Pollution Technology*, 25(1), D1800. <https://doi.org/10.46488/NEPT.2026.v25i01.D1800>

Note: From 2025, the journal has adopted the use of Article IDs in citations instead of traditional consecutive page numbers. Each article is now given individual page ranges starting from page 1.



Copyright: © 2026 by the authors

Licensee: Technoscience Publications

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

Assessment of groundwater quality is becoming essential for effective resource management. In this study, we conducted a seasonal assessment of groundwater quality, hydrogeochemical processes, and statistical analyses in the city of Moundou. A total of 62 groundwater samples were taken in the 27 districts of the city, in August 2022 (rainy season) and May 2023 (dry season), respectively. From a geochemical point of view, the results highlighted two dominant geochemical facies in both the dry and wet seasons: the calcium-magnesium bicarbonate facies (Ca-Mg-HCO₃) and the sodium-potassium bicarbonate facies (Na-K-HCO₃). The multivariate analysis showed that the mineralisation of gypsum (CaSO₄ · 2H₂O), CaCl₂ salts, silicates, carbonates and the decomposition of organic matter are the main processes affecting the quality of Moundou's water. The physico-chemical results show that of the parameters monitored, only pH, iron and ammonium do not comply with the WHO standard, and almost 89% of the sites sampled are considered acceptable according to QWI values. Only the sites in the north-east of the city showed poor water quality during the rainy season. On the whole, this water is of better quality for irrigation.

INTRODUCTION

The degradation of groundwater quality worldwide has intensified in recent years due to uncontrolled industrial discharges, excessive use of chemical fertilisers in agriculture and inefficient management of water resources. (Khaldi et al. 2018). These factors alter the chemical composition of water, making it unsuitable for its intended uses. (Naima et al. 2022). Water resources, whether groundwater or surface water, are used by humans for a wide range of purposes (Ahoussi et al. 2010). The chemical composition of water from the natural environment varies considerably. It is influenced by the geological nature of the soil from which it comes, as well as by the reactive substances in contact during its flow. The presence of suspended and dissolved matter, whether of mineral or organic origin, affects the quality of groundwater, both quantitatively and qualitatively (Ngouala Mabonzo 2020). In developing countries, groundwater is a key resource for supplying people with drinking water. They generally offer satisfactory quality at a low cost, unlike surface water, which is expensive to treat and often inaccessible to nations with fragile economies, as is the case in many African countries (Yao et al. 2016).

In many parts of the world, particularly in developing countries, access to water is now essential for public health and socio-economic growth (Kettab et al. 2008). Of all the freshwater on Earth (2.53%), only 0.63% is groundwater. Groundwater is the world's main source of drinking water, as it is generally better protected from

pollution than surface water (Edith et al. 2023). Groundwater quality can deteriorate when it is exposed to harmful or even toxic substances as a result of human activities. Although essential to life, water can also become a vector for water-borne diseases due to its contamination by various mineral and organic wastes, as well as by excrement (Ndahama et al. 2014). In Chad, the expansion of major urban centres is leading to the establishment of new neighbourhoods in areas that are sometimes unsuitable for housing. At the same time, the water supply network is not expanding at the same pace as this growth, and the few neighbourhoods that are served suffer frequent interruptions. Faced with this situation, the most affluent households invest in human-powered boreholes, whose water quality is not systematically checked by qualified experts.

On the other hand, the most precarious households, located far from these boreholes, are resigned to using water from traditional wells (Mahamat et al. 2015). For several decades now, water quality has been a major issue, contributing to the spread of diarrhoeal diseases. In Chad, this problem affects almost the entire country, with greater intensity in the Saharan and Sahelian regions. To meet their water needs, whether domestic, agricultural or for livestock, people draw on traditional and modern boreholes, as well as rivers and ponds (Kriga et al. 2016).

Like most towns in Chad, the city of Moundou has a low drinking water supply. This situation can be explained on the one hand by strong urbanisation and exponential population growth. On the other hand, the drinking water supply network, with a coverage rate of 47%, is not keeping up with the high rate of urbanisation. In addition, the network is outdated, leading to water leaks. In addition, residents of neighbourhoods not served by the current network obtain their water directly from the river or from the surface water table, via boreholes and private wells. However, due to a lack of sanitation, these sources are often contaminated by the infiltration of wastewater, black water and other pollutants, leading to cases of water-borne diseases among the population (Urbaplan 2008). Despite this high anthropogenic pressure on water, very few studies have looked at the quality of groundwater in the town of Moundou. The only studies identified in the literature are those by Bandé (2016) and Schneider & Wolff (1992), which looked at hydrochemical facies, among other things. No study has examined the spatial and seasonal variation in water quality in the town of Moundou. This study was initiated to fill this information gap, which is crucial for the integrated management of the city's water resources. The aim of the study is therefore to assess the spatial and seasonal distribution of water quality in Moundou. This will involve determining the hydrochemical facies of the water, assessing

the spatial and seasonal variability of the physico-chemical parameters and, finally, the potential uses (drinking water and water for agriculture).

MATERIALS AND METHODS

Study Area

Moundou is located in south-western Chad, in the Logone Occidental region, of which it is the capital. The country's second most populous city after N'Djaména, Moundou is considered to be the economic capital of Chad, due to the presence of numerous industries, particularly agri-food and brewing. It lies between 8°33' and 8°36' North and 16°03' and 16°06' East (Djako 2018). Moundou lies between 382 and 439 metres above sea level (Schneider & Wolff 1992). Demographic growth and accelerated urbanisation are leading to increasing pressure on water resources, with risks of pollution and overexploitation of groundwater (Mahamat et al. 2015). Moundou is located in the Sudanian climate zone, characterised by two contrasting seasons (the dry season and the rainy season). The rainy season runs from May to October, with average annual rainfall of between 900 and 1,200 mm, while the dry season extends from November to April and is dominated by the harmattan, a dry, dusty wind from the Sahara (Djebe et al. 2022). Temperatures are high, averaging between 25 and 35°C, with peaks reaching 40°C during periods of intense heat (Khaldi et al. 2018). The Moundou region belongs to the geological formations of the Terminal Continental, characterised by varied sedimentary deposits composed mainly of obliquely stratified sands and sandstones, multicoloured clays and kaolin and gravel and conglomerate lenses, bearing witness to ancient fluvial environments (Schneider & Wolff 1992). Underlying these formations is the Precambrian crystalline and metamorphic basement, which forms an impermeable barrier limiting the storage of groundwater (Mathieu 1980). Two types of aquifer dominate the hydrogeology of Moundou. The Continental Terminal water table, which is shallow (between 5 and 15 metres), is rapidly recharged in the rainy season and is highly vulnerable to anthropogenic pollution (latrines, waste water, solid waste, etc.). The Precambrian basement aquifers, which are continuous and located in rock fractures, are deep (between 50 and 150 metres) and generally of better quality, but require costly drilling (Schneider & Wolff 1992). Groundwater supply is therefore based on the joint exploitation of these aquifers, with disparities in quality and availability. The Logone River, which is the main watercourse and a source of supply for a number of uses (domestic, agricultural and industrial); Lake Wey and Lake Taba, located to the west and east of the town respectively, which contribute to groundwater recharge by infiltration




Fig. 1: Map of the study area and sampling points.

during flood periods; Lake Taba and the numerous temporary pools, which form during the rainy season and can influence the recharge of surface water (Djebe et al. 2022).

Sample Collection and Laboratory Analysis

A total of 62 groundwater samples from wells and boreholes were collected during the dry and rainy seasons. These samples were analysed at the Laboratoire National des Eaux (LNE) to assess their quality. The analyses focused on chemical parameters such as Na^+ , K^+ , Mg^{2+} , Ca^{2+} , HCO_3^- , SO_4^{2-} , NO_3^- and Cl^- . As soon as they were collected, the samples were placed in a refrigerated cooler and transported to the analysis laboratory. Samples were taken in the 27 districts of the town of Moundou, as shown in Fig. 1, which shows the geographical distribution of the collection points. Physical parameters, including pH, temperature, conductivity and turbidity, were measured directly on site using a multiparameter. The colour of the water was determined in accordance with French standard NF T 90-034. As regards chemical analyses, the concentrations of Fe^{2+} , Na^+ , Cl^- , SO_4^{2-} , F^- , HCO_3^- , NO_3^- and NO_2^- were assessed in a specially equipped laboratory (LNE) in N'djaména, the Chadian capital. These analyses were carried out using a spectrophotometer, in accordance with the standard methods recommended by the French standards AFNOR.

Hydrochemistry of Groundwater

The Piper diagram is a widely used approach for identifying the hydrochemical type and facies of groundwater (Ousmane et al. 2022). According to this classification, groundwater is divided into main groups according to the dominant cations and anions: Mg^{2+} , Ca^{2+} , Na^+ and K^+ ; SO_4^{2-} , HCO_3^- , Cl^- . This diagram thus provides a better understanding of mineralisation processes and the geochemical evolution of groundwater. The factors influencing the chemical composition of groundwater will be analysed using the semi-logarithmic diagrams developed by Gibbs and McIntyre (1970). These diagrams represent the relationship between TDS and the ratios $\text{Cl}/(\text{Cl}+\text{HCO}_3^-)$ for anions, and $\text{Na}/(\text{Na}+\text{Ca})$ for cations. According to this approach, the main mechanisms controlling the hydrochemical characteristics of groundwater can be grouped into three dominant categories: dominance of water-rock interaction, influence of precipitation, and impact of evaporation (Ngouala Mabonzo 2020). Interpretation of these diagrams provides a better understanding of the geochemical processes behind groundwater mineralisation.

Multivariate Statistical Analysis

For the study of groundwater in the city of Moundou, multivariate statistical analyses were performed to examine the interactions between the different hydrochemical

Table 1: Standard values, weights and relative weights of hydrochemical parameters (WHO 2011).

Hydrochemical parameters	Unit	Standard value (WHO 2011)	Weight [Wi]	Relative weight [Wi']
pH	-	6,5 - 8,5	4	0,085
Conductivity (CE)	$\mu\text{S.cm}^{-1}$	2500	5	0,106
TDS (Dissolved Solids)	mg.L^{-1}	500	5	0,106
Calcium (Ca^{2+})	mg.L^{-1}	100	2	0,043
Magnesium (Mg^{2+})	Mg.L^{-1}	50	2	0,043
Sodium (Na^+)	mg.L^{-1}	200	4	0,085
Potassium (K^+)	mg.L^{-1}	12	4	0,085
Bicarbonates (HCO_3^-)	mg.L^{-1}	500	4	0,085
Chlorides (Cl^-)	mg.L^{-1}	250	4	0,085
Sulphates (SO_4^{2-})	mg.L^{-1}	250	2	0,043
Nitrates (NO_3^-)	mg.L^{-1}	50	5	0,106
Iron (Fe^{2+})	mg.L^{-1}	0,3	3	0,064
Ammonium (NH_4^+)	mg.L^{-1}	0,5	3	0,064
			47	

parameters (Amroune 2018). In addition, the Pearson correlation matrix was used to identify relationships between hydrochemical parameters. A correlation coefficient between 0.5 and 0.7 was interpreted as a moderate relationship, while a value greater than 0.7 was considered highly significant (Bourjila 2023). Principal Component Analysis (PCA) was implemented to reduce the dimensionality of the data and identify the main sources of influence of groundwater hydrochemical parameters (Soro et al. 2019). When calculating the PCA, a rotation of the principal components was carried out using the Varimax method, allowing better interpretation of the results (Hassan & Firat Ersoy 2022).

Water Quality for Drinking

The Water Quality Index (WQI) is an essential tool for the overall assessment of the quality of groundwater intended for human consumption (Chaima 2024). It is calculated by assigning weights, ranging from 2 to 5, to the various hydrochemical parameters, according to their influence on water quality. First, the relative weights (Wi) were determined using equation (1), making it possible to quantify the impact of each parameter on water classification.

$$W_i = \frac{w_i}{\sum_1^n w_i} \quad \dots(1)$$

The weights assigned and the relative weights of each parameter are listed in Table 1 (Soro et al. 2022). In the second phase, the quality index (Qi) for each hydrochemical parameter was determined by calculating the ratio between the measured value and the reference standard established by WHO (2011), in accordance with equation (2).

$$Q_i = \frac{C_i}{S_i} \times 100 \quad \dots(2)$$

Ci: The measured value of the parameters, Si: standard value of the parameters

In the final step, the water quality sub-index and the WQI are obtained using equations (3) and (4)

$$SL_i = Q_i \times W_i \quad \dots(3)$$

$$WQI = \sum_1^n SL_i \quad \dots(4)$$

After calculating the WQI, Table 2 is used to classify the different types of groundwater quality.

To determine the areas where the risk of water pollution is high, the groundwater quality indices (WQI) calculated at each site were used as input data for extrapolating the WQI in the non-sampled areas using the inverse distance interpolation (IDW) method.

Water Quality for Agriculture

To assess the suitability of groundwater for agricultural purposes, Wilcox diagrams have been drawn using the value of the sodium absorption ratio (SAR) as a function of EC (Wilcox 1955). The SAR can be calculated by equation (5).

$$SAR = \frac{Na^+}{\sqrt{\frac{Mg^{2+} + Ca^{2+}}{2}}} \quad \dots(5)$$

Table 2: WQI water quality index classification.

Classification	Groundwater quality
WQI < 25	Excellent water
25 < WQI < 50	Good water
50 < WQI < 75	Poor water
75 < WQI < 100	Very poor water
WQI > 100	Unsuitable water

Table 3: The quality of groundwater for irrigation has been classified into four categories (Fallahati et al. 2020).

Class	Water quality for irrigation
C1S1	Sweet
C1S2 C2S2 C2S1	A little salty
C1S3 C2S3 C3S1 C3S2 C3S3	Salty
C1S4 C2S4 C3S4 C4S4 C4S3 C4S2 C4S1	Very salty

The groundwater quality for irrigation was categorized into four classes, as presented in Table 3.

RESULTS AND DISCUSSION

Global Statistics of Physicochemical Analyses

The overall characteristics of the hydrogeochemical variables used in this study concern the minimum-maximum values, the mean, the standard deviation and the coefficient of variation (Table 4). The average contents in the rainy season and in the dry season were compared with the admissible values for drinking water proposed by the WHO (2011). The results show that the pH of the groundwater of the city of Moundou was acidic, with an average of 5.14 in the rainy season and 5.77 in the dry season, i.e., an overall average of 5.45, which is lower than the limit recommended by the WHO (2011). The low conductivity values recorded suggest

that the waters of this city are weakly mineralized. The average value of electrical conductivity in the rainy season ($427 \mu\text{S}\cdot\text{cm}^{-1}$) is significantly higher than that in the dry season ($262 \mu\text{S}\cdot\text{cm}^{-1}$), which corroborates well with the fact that the average value of most ions measured in the rainy season is also higher than that measured in the dry season. This seems contradictory because of the dilution phenomena that take place in the rainy season. This phenomenon could be explained by anthropogenic pollution sources, such as wastewater or agricultural activities, that contribute to an increase in ions in groundwater during the rainy season following infiltration and runoff. These waters are also soft and very aggressive, whatever the season. Of all the parameters monitored, only iron and ammonium are higher than the WHO standard. Iron would come from rocks, such as silicates, oxides, hydroxides, carbonates and sulfides or from organic matter, while ammonium has a purely organic origin. In the rainy season, in addition to temperature and pH, which have their standard deviation significantly lower than the mean, most of the monitored parameters have their standard deviation close to or higher than the mean, suggesting that all parameters except temperature and pH vary greatly from one sampling site to another. The same observations were made in the dry season, except that nitrate and ammonium did not experience a strong spatial variation. This low spatial variation of nitrate and ammonium indicates

Table 4: Statistical characteristics of physicochemical analyses (August 2022 and May 2023).

	Moyenne		Écart-type			Minimum			Maximum			WHO	
	RS	DS	AN	RS	DS	AN	RS	DS	AN	RS	DS		AN
pH	5.14	5.77	5.45	0.632	0.441	0.629	3.26	4.32	3.26	6.30	6.47	6.47	6.5–8.5
CE [$\mu\text{S}\cdot\text{cm}^{-1}$]	427	262	345	572	266	452	14.5	15.5	14.5	3670	1000	3670	750
TDS [$\text{mg}\cdot\text{L}^{-1}$]	213	152	182	287	176	239	7.27	3.86	3.86	1840	961	1840	500
Temp [$^{\circ}\text{C}$]	29.2	30.0	29.6	1.21	1.60	1.46	26.2	25.1	25.1	33.0	33.7	33.7	-
Turb [NTU]	2.14	1.01	1.58	3.65	0.601	2.66	0.100	0.100	0.100	26.0	4.00	26.0	5
CaCO ₃ [$\text{mg}\cdot\text{L}^{-1}$]	43.6	31.0	37.3	54.2	39.0	47.5	4.00	0.400	0.400	300	300	300	-
Ca ²⁺ [$\text{mg}\cdot\text{L}^{-1}$]	12.9	9.58	11.3	14.8	10.6	12.9	1.40	0.100	0.100	80.0	80.0	80.0	75
Mg ²⁺ [$\text{mg}\cdot\text{L}^{-1}$]	2.74	1.71	2.23	4.48	3.20	3.91	0.01	0.00	0.00	24.3	24.3	24.3	30
K ⁺ [$\text{mg}\cdot\text{L}^{-1}$]	1.64	1.05	1.35	2.05	1.13	1.67	0.100	0.00	0.00	14.0	7.00	14.0	30
Na ⁺ [$\text{mg}\cdot\text{L}^{-1}$]	16.4	9.30	12.8	17.8	10.1	14.8	0.400	0.200	0.200	89.0	66.0	89.0	200
HCO ₃ ⁻ [$\text{mg}\cdot\text{L}^{-1}$]	53.2	35.3	44.2	66.4	34.8	53.6	3.70	0.500	0.500	366	244	366	500
Cl ⁻ [$\text{mg}\cdot\text{L}^{-1}$]	14.2	10.3	12.3	16.2	12.2	14.4	0.800	0.200	0.200	100	90.0	100	250
SO ₄ ²⁻ [$\text{mg}\cdot\text{L}^{-1}$]	4.73	2.66	3.70	7.71	8.80	8.30	0.00	0.00	0.00	36.0	63.0	63.0	250
NO ₃ ⁻ [$\text{mg}\cdot\text{L}^{-1}$]	12.7	6.32	9.50	15.9	4.93	12.1	0.660	0.200	0.200	88.0	37.0	88.0	45
Fe ²⁺ [$\text{mg}\cdot\text{L}^{-1}$]	0.0271	0.143	0.08	0.052	0.271	0.195	0.00	0.00	0.00	0.220	1.00	1.00	0.01
NH ₄ ⁺ [$\text{mg}\cdot\text{L}^{-1}$]	1.82	0.740	1.29	8.29	0.248	5.92	0.110	0.110	0.110	66.0	1.12	66.0	0.25

DS = Dry season, RS = Rainy season, AM = Annual mean, Temp = température, Turb = turbidity




Fig. 2: Piper diagram for the rainy and dry seasons.

that runoff plays an important role in the contamination of groundwater with nitrogen in this city.

Geochemical Facies of Groundwater

Piper's diagram allows us to characterize the chemical composition of the water by representing the distribution of the main cations (Ca^{2+} , Mg^{2+} , Na^+ , K^+) and anions (SO_4^{2-} , Cl^- , HCO_3^- , etc.) (Piper 1944). As the 62 water points were sampled during both the wet and dry seasons, it makes sense to describe them by season. The geochemical facies given by Piper's diagrams are shown in Fig. 2a and 2b, which represent Piper's diagrams for the rainy and dry seasons, respectively. There was no significant difference between the two seasons, suggesting that season does not influence water facies in Moundou. In the lower left-hand section of each diagram, the dominant cations in all samples are calcium, followed by magnesium; while the anions, in the lower right-hand section, are dominated by HCO_3^- and sulfates (SO_4^{2-}), with a non-negligible contribution from chlorides (Cl^-). This results in two types of geochemical facies dominated by the calcium-magnesium bicarbonate facies (Ca-Mg-HCO_3), generally observed in recent, relatively undeveloped groundwater resulting from infiltration of precipitation and interaction with carbonate and silicate rocks (Olivier 2015), and the sodium-potassium bicarbonate facies (Na-K-HCO_3), which results mainly from chemical alteration processes of silicate rock minerals. A slight scattering of points suggests mixed influences, with possible anthropogenic input (urban and agricultural runoff) (Edmunds & Smedley 2000). These hydrochemical facies obtained by the diagram software reflect the local homogeneity of the geology of the town

of Moundou, as indicated by Bandé (2016). Aquifers in sedimentary environments are more or less continuous, unlike those in basement environments, where several studies have revealed the discontinuous nature of the latter environment; this is linked to the polyphasic alteration that occurs there (Mfonka et al. 2021). These results are comparable to those obtained by Mfonka et al. (2024) at N'Djaména in Chad.

Groundwater Type and Facies Control Factors

The Gibbs diagram is a tool for identifying the dominant processes that control groundwater chemistry, as a function of total salinity (TDS - Total Dissolved Solids) and ionic ratios (Meryem 2021). He distinguishes three major processes, namely Precipitation Dominance (Influence of precipitation), Rock Dominance (Water-rock interaction) and Evaporation Dominance (Concentration by evaporation). The diagram in Fig. 3 (a) shows that the majority of points are located in the 'Rock Dominance' zone, with a high concentration which reflects the influence of rock alteration on groundwater chemistry; the presence of Na^+ and Ca^{2+} suggests mineral alteration processes such as the dissolution of carbonates (calcite, dolomite) and silicates (feldspars) and finally a small proportion of samples are located in the 'Precipitation Dominance' zones showing that precipitation has a relatively limited influence on water chemistry. The diagram in Fig. 3 (b) shows a high concentration of all points in the 'Rock Dominance' zone, confirming that water-rock interaction is the main factor influencing the chemical composition of Moundou groundwater. The Gibbs diagram for the dry season in Fig. 4 (a) and (b) shows a similarity

to that for the rainy season, but there is a slight scattering of points in both the cation and anion diagrams. This phenomenon can be attributed to an increased concentration of dissolved salts due to evaporation, leading to a change in ionic ratios and greater variability in dissolution and mineral precipitation processes, linked to fluctuations in water table levels and greater mineralisation of groundwater. This result is consistent with that of Bouchagoura (2019). This analysis confirms the major influence of water-rock interactions on groundwater mineralisation, with a possible impact of precipitation and evaporation depending on the season

Multivariate Statistical Analysis (Correlation Analysis and Principal Component Analysis)

Correlation Analysis

The correlation matrix obtained from the rainy and dry season data is presented in Table 5. All the major ions showed a strong correlation with each other in both the rainy

and dry seasons. Indeed, Ca^{2+} and Mg^{2+} correlated strongly with CaCO_3 (0.99 and 0.95, respectively), confirming their role in water hardness due to the dissolution of carbonates (calcite, dolomite). HCO_3^- is also correlated with CaCO_3 (1.00), reinforcing the hypothesis of carbonate alteration as a major source of ions. Na^+ and Cl^- show a strong positive correlation (0.96), suggesting a common origin, possibly the alteration of evaporitic rocks (halite) or anthropogenic inputs (saline intrusion, pollution). Sulphate (SO_4^{2-}) correlates well with Na^+ (0.86) and Cl^- (0.77), suggesting a common source, such as dissolution of gypsum or other sulphide minerals. NO_3^- was moderately correlated with CE/TDS (0.36), suggesting a possible influence of precipitation and infiltration containing nitrates of agricultural or domestic origin. In the dry season, iron showed no strong correlation with the other parameters, indicating that it could be due to reducing conditions (dissolution of iron oxides in oxygen-poor environments). In the rainy season, however,




Fig. 3: Gibbs diagram for the rainy season.




Fig. 4: Gibbs diagram for the dry season.

Table 5: Correlation matrix of physicochemical parameters in rainy seasons (SP) and dry seasons (SS).

SP	pH	CE	TDS	Temp	Turb	CaCO ₃	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻	NO ₃ ⁻	Fe ²⁺	NH ₄ ⁺
pH	1															
CE	0,25	1,00														
TDS	0,25	1,00	1,00													
Temp	-0,37	-0,28	-0,28	1,00												
Turb	0,20	0,00	-0,01	-0,11	1,00											
CaCO ₃	0,20	0,32	0,32	-0,07	0,28	1,00										
Ca ²⁺	0,21	0,29	0,29	-0,06	0,36	0,99	1,00									
Mg ²⁺	0,16	0,36	0,36	-0,09	0,08	0,95	0,90	1,00								
K ⁺	0,16	0,21	0,21	-0,06	0,10	0,55	0,55	0,52	1,00							
Na ⁺	0,15	0,32	0,32	-0,02	0,21	0,97	0,95	0,94	0,58	1,00						
HCO ₃ ⁻	0,20	0,32	0,32	-0,07	0,28	1,00	0,99	0,95	0,56	0,97	1,00					
Cl ⁻	0,17	0,32	0,31	-0,02	0,23	0,99	0,98	0,94	0,53	0,96	0,99	1,00				
SO ₄ ²⁻	0,13	0,18	0,17	-0,01	0,16	0,79	0,79	0,75	0,63	0,86	0,79	0,77	1,00			
NO ₃ ⁻	0,07	0,36	0,36	0,05	0,23	0,87	0,86	0,84	0,50	0,92	0,87	0,88	0,68	1,00		
Fe ²⁺	-0,04	0,09	0,09	0,25	-0,03	0,18	0,17	0,18	0,08	0,24	0,18	0,19	0,09	0,35	1,00	
NH ₄ ⁺	-0,21	-0,05	-0,05	0,13	-0,01	-0,05	-0,05	-0,05	-0,03	-0,04	-0,05	-0,03	-0,08	-0,02	-0,06	1,00

SS	pH	CE	TDS	Temp	Turb	CaCO ₃	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻	NO ₃ ⁻	Fe ²⁺	NH ₄ ⁺
pH	1,00															
CE	0,24	1,00														
TDS	0,21	0,92	1,00													
Temp	-0,34	-0,13	-0,18	1,00												
Turb	0,13	0,33	0,37	-0,05	1,00											
CaCO ₃	0,27	0,48	0,45	-0,11	0,59	1,00										
Ca ²⁺	0,29	0,48	0,45	-0,10	0,58	0,99	1,00									
Mg ²⁺	0,23	0,47	0,44	-0,13	0,59	0,97	0,94	1,00								
K ⁺	0,27	0,60	0,63	-0,15	0,42	0,87	0,86	0,85	1,00							
Na ⁺	0,31	0,48	0,40	-0,13	0,45	0,93	0,92	0,92	0,91	1,00						
HCO ₃ ⁻	0,31	0,50	0,45	-0,13	0,50	0,98	0,98	0,93	0,90	0,97	1,00					
Cl ⁻	0,29	0,49	0,44	-0,14	0,55	0,99	0,98	0,96	0,89	0,96	0,98	1,00				
SO ₄ ²⁻	0,19	0,49	0,44	-0,08	0,59	0,93	0,89	0,96	0,81	0,90	0,88	0,91	1,00			
NO ₃ ⁻	0,31	0,42	0,44	-0,06	0,56	0,88	0,88	0,84	0,83	0,83	0,86	0,85	0,79	1,00		
Fe ²⁺	0,03	0,23	0,28	-0,20	0,27	-0,12	-0,09	-0,17	-0,20	-0,24	-0,17	-0,16	-0,11	-0,14	1,00	
NH ₄ ⁺	-0,03	0,13	0,16	-0,12	-0,11	0,28	0,31	0,21	0,30	0,28	0,29	0,31	0,22	0,26	0,07	1,00

it was positively correlated with nitrate, indicating that some of the iron comes from anthropogenic activities. Ammonium (NH₄⁺) is weakly correlated with the other parameters, which may indicate a local origin linked to organic sources (agricultural or domestic pollution or decomposition of organic matter). In the dry season, electrical conductivity (EC), total dissolved solids (TDS) and turbidity showed moderate positive correlation with all major ions, which was not the case in the wet season due to dilution.

Principal Component Analysis (PCA)

Principal component analyses were carried out to determine the relationship between physico-chemical variables and potential sources of groundwater pollution in the dry and rainy seasons. Concerning the number of principal components to be retained, we chose the eigenvalue analysis method (Table 6); thus, the factorial axes with eigenvalues of less than 1 were eliminated, which made it possible to retain the first four principal components in both the wet and dry seasons (Table 6). In the rainy season, the four

Table 6: Contributions of the components.

	Rainy season				Dry season			
	PC1	PC2	PC3	PC4	PC1	PC2	PC3	PC4
Eigenvalue	7.80	1.75	1.14	1.09	7.46	1.48	1.42	1.31
% variance	51.98	11.65	7.59	7.25	49.72	9.85	9.48	8.76
pH	0.11	0.72	0.22	0.18	0.23	0.82	0.21	-0.04
CE	0.34	0.5	0.09	-0.5	0.49	-0.03	0.07	0.4
Temp	0	-0.82	0.18	0.03	-0.02	-0.83	0.2	-0.12
Turb	0.22	0.15	0.01	0.82	0.07	0	0.73	0.48
CaCO ₃	0.98	0.09	0.06	0.08	0.96	0.12	-0.05	0
Ca ²⁺	0.96	0.09	0.06	0.17	0.91	0.13	-0.07	0.02
Mg ²⁺	0.94	0.09	0.05	-0.11	0.91	0.07	-0.01	-0.04
K ⁺	0.63	0.11	-0.06	-0.06	0.87	0.07	-0.08	-0.1
Na ⁺	0.98	0.03	0.09	0.01	0.95	0.09	-0.05	-0.13
HCO ₃ ⁻	0.98	0.09	0.06	0.08	0.96	0.12	-0.06	-0.07
Cl ⁻	0.97	0.04	0.06	0.04	0.94	0.13	-0.11	-0.03
SO ₄ ²⁻	0.84	0.02	-0.01	0.07	0.77	-0.09	-0.01	0.12
NO ₃ ⁻	0.91	-0.06	0.19	-0.03	0.64	0.16	0.17	-0.31
Fe ²⁺	0.2	-0.34	0.72	-0.2	-0.23	0.11	-0.01	0.85
NH ₄ ⁺	0.04	-0.35	-0.69	-0.14	0.2	0.02	-0.86	0.19

principal components explained 78.5% of the total variance. The first principal component (PC1) explained 51.98% of the total variability and was positively correlated with all the major elements. The correlation matrix also showed a strong correlation between all major elements. Thus, PC1 reflects mineralization of natural origin (water-casement contact). The inclusion of nitrates in this batch indicates that PC1 also highlights mineralization of anthropogenic origin. The second principal component (PC2) explains 11.65% of the total variability. PC2 correlates positively with pH and electrical conductivity (EC) and negatively with temperature, highlighting the influence of climatic conditions on the mineralisation process. With 7.59% of the total variance explained, the third principal component (PC3) is strongly and positively correlated with iron and ammonium. This component seems to highlight the process of decomposition of organic matter. The fourth principal component (PC4), associated negatively with electrical conductivity and positively with turbidity, highlighting the dilution process resulting from the infiltration of rain water.

In the dry season, the four principal components explained 77.81% of the total variance (Table 6). The first principal component is positively correlated with all the major ions and electrical conductivity. PC1 explains the same processes in the wet and dry seasons. The absence of a strong correlation observed between major ions and electrical conductivity in the rainy season is linked to the dilution phenomenon. The

second principal component (PC2) is positively related to pH and negatively related to temperature. PC2 also explains the same processes in the wet and dry seasons. PC3 is strongly and negatively correlated with ammonium and positively with turbidity, indicating pollution of organic origin. PC4 is positively associated with iron and turbidity, suggesting that iron is the cause of water turbidity.

Groundwater Quality Assessment

Water Quality for Drinking

Calculation of the WQI of 62 samples for the rainy and dry seasons made it possible to express the proportions of 5 classes of groundwater quality, as shown in Fig. 5 (a) and (b). Fig. 5 (a) shows a marked deterioration in water quality in the rainy season at certain points, such as the Lac Taba carré borehole, where the water was non-drinkable (1.61%); the Guelkoura 1 carré 4 well and the Guelbé Résidence Sœur borehole, where the water was of mediocre quality (3.23%); poor quality water was observed at the 15 ans II carré 3 well, the Doyon carré 2 well and the Mbomia carré 2 borehole (4.84%). 48.39% of samples were of excellent quality, and 41.94% of the samples were of good quality. Unlike the rainy season, the dry season (Fig. 5 (b)) shows a reduction in pollution, as no water point was found to be either undrinkable or of poor quality. The percentage of poor quality water (4.84%) fell to 3.23%. More than half (54.84%) of the samples had excellent water quality, and




Fig. 5: Proportions of groundwater quality classes for the rainy season (a) and the dry season (b).




Fig. 6: WQI spatial distribution map of groundwater for the rainy season (a) and the dry season (b).




Fig. 7: Wilcox diagram for the rainy season (a) and the dry season (b).

41.84% of the samples had good water quality. Fig. 6 (a) and (b) for the rainy and dry seasons, respectively, show the spatial distribution of water quality indices on the map of the town of Moundou.

Water Quality for Irrigation

Overall, the distribution of points shown in Fig. 7 for the wet season (a) and dry season (b) shows that the majority

of points are found in zones C1-S1 and C2-S1, suggesting low to medium salinity with a low Sodicity Index (SAR). Very few points appear in zones C3 and C4, indicating a low accumulation of dissolved salts. Finally, no points are located in zones S3 and S4, confirming a low risk of soil sodification. In the Wilcox diagram in Fig. 7 for the rainy season, the majority of samples are located in zones C1-S1 and C2-S1, indicating that rainfall dilutes the salinity of the

groundwater. The electrical conductivity (EC) is lower due to the freshwater input. One sample is located in zone C4S1, indicating high salinity, and a few samples are located in zone C3S1, reflecting medium salinity. In the dry season, a similar distribution of points is observed, but the electrical conductivity appears to be slightly higher, with the majority of points in zones C1S1 and C2S1, indicating a higher concentration of dissolved salts due to evaporation. A few points are also found in zone C3S1, suggesting average salinity. Nevertheless, the water is generally of good quality for irrigation, with no significant risk of sodification.

CONCLUSIONS

This study assessed the quality of groundwater in the town of Moundou according to seasonal variations and identified the main natural and anthropogenic influences on its chemical composition. Analysis of the hydrochemical facies revealed the predominance of calcium and magnesium bicarbonate facies (Ca-Mg-HCO_3), reflecting an interaction between seepage water and carbonate and silicate rocks. The sodium and potassium bicarbonate facies (Na-K-HCO_3) was also identified, resulting from alteration processes of silicate minerals. These facies were observed during the two seasons studied, with a slight dispersion in the dry season attributed to evaporation. Multivariate statistical analyses revealed sources of contamination of natural origin, linked to the weathering of rocks, as well as anthropogenic sources, notably industrial waste, fertilisers, pesticides, wastewater and metal carcasses. Assessment of drinking water quality revealed that pollution is more marked in the rainy season due to infiltration of run-off water, while in the dry season, quality improves with an increase in the number of samples showing good to excellent water quality. With regard to the suitability of groundwater for irrigation, the Wilcox diagram showed low to medium salinity, making the water suitable for irrigation in both seasons. However, vigilance is still required during the rainy season because of the increased risk of contamination. This study highlights the importance of rigorous monitoring of groundwater quality, particularly in urban areas, in order to prevent pollution risks and ensure sustainable management of water resources. Further investigations, including isotope studies and hydrogeological modelling, could provide a better understanding of the evolution of aquifers and the associated pollution mechanisms.

In light of the results of our study, we suggest that the local authorities in the town of Moundou extend the public drinking water distribution network, which currently has insufficient coverage (47%), in order to limit the population's reliance on polluted surface water. In addition, the authorities must ban groundwater abstraction for drinking water supply

in the north-east of Moundou, identified as the area most vulnerable to pollution. Set up a permanent groundwater quality monitoring programme, particularly in densely urbanised areas, as well as an awareness-raising programme.

ACKNOWLEDGMENTS

The authors would like to thank the National School of Public Works of the University of N'Djaména.

REFERENCES

- Ahoussi, E.K., Soro, N., Kouassi, A.M., Soro, G., Koffi, Y.B. and Zade, S.P., 2010. Application of multivariate statistical analysis methods to study the origin of heavy metals (Cu^{2+} , Mn^{2+} , Zn^{2+} and Pb^{2+}) in groundwater aquifers of Abidjan city. *International Journal of Biological and Chemical Sciences*, 4(5), pp.1201–1218.
- Amroune, A., 2018. *Hydrochemical contribution study for understanding the functioning of the alluvial aquifer in the northern Hodna region (Southeast Algeria)*. PhD Thesis, University of Batna 2, pp.1–250.
- Bande, F., 2016. *Hydrodynamic and hydrochemical characterisation of groundwater in an urban context: Case of Moundou city (Southwest Chad)*. Master's Thesis, University of N'Gaoundéré, pp.1–120.
- Bouchagoura, L., 2019. Study of groundwater quality and vulnerability to pollution of a shallow aquifer in extreme Eastern Algeria: Case of Tébessa region. *Journal of Environmental Hydrogeology*, 12(2), pp.55–78.
- Bourjila, A., 2023. *Hydrochemical modelling of marine intrusion in the Ghiss–Nekor coastal aquifer and characterisation of potential groundwater zones in the Ghiss and Nekor basins (Morocco) using geospatial methods*. PhD Thesis, Abdelmalek Essaâdi University, pp.1–300.
- Chaima, A., 2024. *Assessment of physicochemical quality of groundwater using the water quality index (WQI) and GIS – Case of M'Sila*. PhD Thesis, University of Bordj Bou Arreridj, pp.1–200.
- Djako, A., 2018. Administration report. *Journal of Scientific Management*, 5(1), pp.1–10.
- Djebe, M., Mbaihadjim, J., Dombor, D.D. and Abbo, F.M., 2022. Urban dynamics and morpho-hydrological constraints in Moundou (Southwest Chad). *Journal of Urban Geography Studies*, 14(2), pp.45–62.
- Edith, K., Bati, G. and Florent, B., 2023. Study of groundwater quality and its suitability for consumption: Case of Moundou city (Southwest Chad). *African Journal of Water Resources*, 9(3), pp.101–120.
- Edmunds, W.M. and Smedley, P.L., 2000. Residence time indicators in groundwater: The East Midlands Triassic Sandstone aquifer. *Applied Geochemistry*, 15(6), pp.737–752.
- Fallahati, A., Soleimani, H., Alimohammadi, M., Dehghanifard, E., Askari, M., Eslami, F. and Karami, L., 2020. Impacts of drought phenomenon on the chemical quality of groundwater resources in central Iran: Application of GIS technique. *Environmental Monitoring and Assessment*, 192(1), p.64.
- Gibbs, A.J. and McIntyre, G.A., 1970. The diagram, a method for comparing sequences: Its use with amino acid and nucleotide sequences. *European Journal of Biochemistry*, 16(1), pp.1–11.
- Hassan, A.M. and Firat Ersoy, A., 2022. Statistical assessment of seasonal variation of groundwater quality in the Çarşamba coastal plain, Samsun (Turkey). *Environmental Monitoring and Assessment*, 194(2), p.135.
- Kettab, A., Mitiche, R. and Bennaçar, N., 2008. Water for sustainable development: Issues and strategies. *Water Sciences Review*, 21(2), pp.247–256.
- Khalidi, F., Smati, H., Grara, N., Smati, N., Maizi, N., Boukehili, K. and Gheid, A., 2018. Measurement and control of properties of

- selected groundwater from the Ouled Driss agglomeration, Algeria. *Environment, Risks & Health*, 17(3), pp.253–261.
- Kriga, A., Barka, M. and Mahamat, A., 2016. Advantages of solar pumping systems relating to potable water challenges in Chad. *Journal of Sustainable Water Systems*, 27(1), pp.19–30.
- Mahamat, S.A.M., Maoudombaye, T., Abdelsalam, T., Ndoumtamia, G. and Loukhman, B., 2015. Evaluation of the physicochemical quality of public water supply distributed by the Chadian Water Company in N'Djaména, Chad. *Journal of Applied Biosciences*, 95, pp.8973–8980.
- Mathieu, P., 1980. *Upper Quaternary on the South Shore of Lake Chad*. Geological Research Monograph, pp.1–203.
- Meryem, L., 2021. *Contribution to the Biogeochemical Study of Water from the Hammam Grouz Dam*. PhD Thesis, Abdelhafid Boussouf University Centre, Mila, pp.1–200.
- Mfonka, Z., Kaledje, P.K., Onana, A.A., Nsangou, D., Kpoumie, A., Zammouri, M., Ndjigui, P.D. and Ngoupayou, J.N., 2024. Geosystems and Geoenvironment.
- Naima, B., Saida, K., Abdellah, O., Labed, A. and Faisa, M., 2022. Physicochemical quality of the Sigus River (Northeast Algeria): Characterisation and principal component analysis. *Journal of Biological Resources*, 12(2), pp.20–35.
- Ndahama, N., Bagalwa, M. and Bayomgwa, C., 2014. Study of total and faecal organic pollution in aquatic systems of the Eastern Democratic Republic of Congo. *Africa Science: International Journal of Science and Technology*, 10(2), pp.101–118.
- Ngouala Mabonzo, M., 2020. Hydrochemical characterisation of groundwater in the contact zone between the coastal sedimentary basin and the Lower Precambrian basement in Southwest Congo. *Five Continents*, 10(21), pp.1–20.
- Olivier, A., 2015. *Chemistry and Pollution of Groundwater*. Lavoisier, pp.1–500.
- Ousmane, B.I., Babaye, M.S.A., Nazoumou, Y., Favreau, G., Mahaman, R.A. and Aboubacar, M.S.I., 2022. Groundwater assessment: Application of geochemistry for a case study in the Djirataoua irrigation perimeter, Southeast Niger. *Afrique Science*, 20(1), pp.56–70.
- Piper, A.M., 1944. A graphic procedure in the geochemical interpretation of water-analyses. *Eos, Transactions American Geophysical Union*, 25(6), pp.914–928.
- Schneider, J.L. and Wolff, J.P., 1992. *Geological Map Volume 1*. Geological Survey Publications, pp.1–302.
- Schneider, J.-L. and Wolff, J.P., 1992. *Geological and Hydrogeological Maps at 1/1,500,000 of the Republic of Chad: Explanatory Memoir (Vol. 1)*. BRGM Editions, pp.1–250.
- Soro, G., Soro, T.D., Adjiri, O.A. and Soro, N., 2019. Application of multivariate statistical methods to the hydrochemical study of groundwater in the lakes region (Central Côte d'Ivoire). *International Journal of Biological and Chemical Sciences*, 13(3), pp.1870–1889.
- Soro, T.D., Fossou, N.M.R., Kone, G.Y.M., Oga, Y.M.S. and Soro, N., 2022. Evaluation of groundwater quality in the Grand-Lahou department (Southwest Côte d'Ivoire) using the WQI index. *International Journal of Innovation and Applied Studies*, 38(2), pp.417–427.
- Urbaplan, 2008. Environmental and social impact study of PADUR sub-projects in the City of Moundou. Environmental Assessment Report, pp.1–120.
- Wilcox, L.V., 1955. Classification and use of irrigation waters. U.S. Department of Agriculture, *Technical Bulletin*, 21(4), pp.1–30.
- World Health Organization, 2011. *Strategies for the Safe Management of Drinking Water for Human Consumption: Report of the Secretariat*. WHO, pp.1–150.
- Yao, A.B., Goula, B.T.A., Kane, A., Mangoua, O.M.J. and Kouassi, K.A., 2016. Groundwater potential mapping of the Lobo watershed (Central-West Côte d'Ivoire): A multi-criteria analysis approach. *Hydrological Sciences Journal*, 61(12), pp.1–12. [DOI]