

Potential Efficiency of Green Algae *Scenedesmus quadricauda* in Bioremediation of Polycyclic Aromatic Hydrocarbon, Benzo[a] Pyrene (BaP)

Hala R. Mohammed¹, Jasim Mohammed Salman²† and Adi Jassim Abd Al-Rezzaq²

¹College of Pharmacy, University of Babylon, Hilla, Iraq

²Department of Biology, College of Science, University of Babylon, Hilla, Iraq

†Corresponding author: Jasim Mohammed Salman; jasimsalman@uobabylon.edu.iq

Abbreviation: Nat. Env. & Poll. Technol.

Website: www.neptjournal.com

Received: 14-08-2024

Revised: 02-09-2024

Accepted: 12-09-2024

Key Words:

Bioremediation
Scenedesmus quadricauda
 Benzo[a]pyrene
 BG11 medium

Citation for the Paper:

Mohammed, H. R., Salman, J. M. and Abd Al-Rezzaq, A. J., 2025. Potential efficiency of green algae *Scenedesmus quadricauda* in bioremediation of polycyclic aromatic hydrocarbon benzo[a]pyrene (BaP). *Nature Environment and Pollution Technology*, 24(2), p. D1712. <https://doi.org/10.46488/NEPT.2025.v24i02.D1712>

Note: From year 2025, the journal uses Article ID instead of page numbers in citation of the published articles.

ABSTRACT

Using algae to break down or detoxify dangerous environmental pollutants, thereby changing them into a non-hazardous condition, is known as bioremediation. Investigating the ability of the green algae *Scenedesmus quadricauda* (Turpin) Brébisson to break a particular polycyclic aromatic hydrocarbon (PAH) known as Benzo[a]pyrene (BaP), Under regulated laboratory circumstances and on BG11 media, the alga was cultivated and exposed to different BaP dosages (0.5, 1, and 1.5 mM). High-performance liquid chromatography (HPLC) study helped to ascertain the BaP concentration. Involving the growth curve, doubling time, photosynthetic pigments, total protein, carbohydrates, and Lipid peroxidation (Malondialdehyde MDA) levels, the research investigated various physiological and biochemical aspects. Furthermore, measured were the levels of catalase (CAT), superoxide dismutase (SOD), and reactive oxygen species (ROS). Whereas the lowest growth rate was 0.00047 on the 15th day at a concentration of 1.5 mM, the maximum growth rate (k) recorded was 0.391 on the 7th day at a concentration of 0.5 mM. Doubling time also varied from 0.00014 throughout the 15th day with 1.5 mM and from 0.1179 throughout the 7th day with 0.5 mM BaP. The results showed a definite influence of the different quantities of BaP degradation by *S. quadricauda*; the greatest magnitude was 40.13 throughout the 15th with 0.5 mg.L⁻¹, while the lowest magnitude was 0 throughout the 1st day with 0.5 Mm. While the min magnitude was 0.41 µg.mL⁻¹ in 0.5 mM throughout 1st day, the max magnitude of chlorophyll-a was 18.71 (µg.mL⁻¹) in 1.5 mM throughout the 15th day. Whereas the greatest magnitude was 9.19 µg.mL⁻¹ in 1.5 mM throughout the 15th day, the lowest magnitude of chlorophyll b was 0.36 µg.mL⁻¹ in 1.5 mM throughout the 1st day. While the min was 0.013 on 1st day with 1 mM, the max magnitude of ROS was 0.28 until the 15th day with 1.5 mM. With 1 mM over 1st day, the carbohydrate showed a max magnitude of 35.13 µm.mL⁻¹, and with 1.5 mM over the 15th day, the min magnitude was 12.25(µm.mL⁻¹). While the min protein content was 1.83 µg.mL⁻¹ in 1.5 Mm throughout the 8th day, the max protein content was 2.14 µg.mL⁻¹ in 1 mM throughout the 8th day; moreover, SOD fluctuated between 22.22 µg.mL⁻¹ in 0.5 mM throughout 1st day, and 60 µg.mL⁻¹ in as the min magnitude throughout 8th day with 1.5 mM. The results show that magnitudes of CAT fluctuated between 13.33 µg.mL⁻¹ in the 8th and 15th mM throughout the 15th day and 73.33 µg.mL⁻¹ in 1 mM throughout the 15th day. MDA showed the largest magnitude 59.92 µmoL.L⁻¹ in 1.5 mM over the 1st day, while the lowest magnitude, 36.58 µmoL.L in 1 mM over the 15th day.

INTRODUCTION

In many environmental media, bioremediation is the use of biological systems to either eradicate or reduce pollution (Luka et al. 2018). Studies have shown that it is a safe, effective, cheap, environmentally friendly replacement for sustained remedial action against hazardous and recurring pollutants (Shukla et al. 2010, Singh 2006). Unwanted effects of pollution and environmental harm follow naturally from the advancement in industry and technology. The main cause of the environmental effects connected with industrial activity is the development of industrial waste.



Copyright: © 2025 by the authors

Licensee: Technoscience Publications

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Because of problems with conventional approaches to pollution control (Nandal et al. 2015). Reducing pollutants and sources of pollution in water is of utmost importance given the severe scarcity of freshwater worldwide, and ensuring that these treatments are economical and environmentally friendly makes them a suitable option compared to traditional alternatives such as chemical precipitation and ion exchange. These alternatives are used to produce waste that is difficult to dispose of (Tewari et al. 2023). Many sophisticated operations integrating biological, physical, and chemical technologies were developed and used recently to clean polluted regions. In this regard, biological therapy—which uses microorganisms like bacteria, fungi, and algae to break down pollutants such as hydrocarbons and convert them into ecologically friendly compounds—is very promising (Amran et al. 2022). Bioremediation is the process of breaking down organic contaminants into water and carbon dioxide or turning organic pollutants into non-toxic metabolites (Alexander 1999). Often used for the removal of different aquatic pollutants, microalgae are also very important in bioremediation (Hwang et al. 2016). Specifically employed for the conversion and breakdown of polycyclic aromatic compounds (PAHs) and removal from the environment, green algae, As shown by García de Llasera et al. (2016), were used to break down enduring toxins like *Scenedesmus* sp.

Organic substances with carbon and hydrogen atoms bound together in a chemical structure called polycyclic aromatic hydrocarbons (PAHs). One may say of them as benzene rings joined in different configurations without any extra atoms or substituents present (Lawal, 2017). Though there are more than 100 kinds of PAHs, the United States Environmental Protection Agency (US EPA) has ranked 16 polycyclic aromatic hydrocarbons (PAHs) based on their unique characteristics as top priority (Keith 2015). Based on their ring count, the molecules listed might be classified as high molecular weight, medium molecular weight, or low molecular weight compounds (Alegbeleye et al. 2017, Pandya & Kumar 2021). Each of the low molecular weight compounds (LMWC), fluoranthene [Fla], phenanthrene [Phen], anthracene [Ant], fluorene [Flu], acenaphthene [Acp], acenaphthylene [Acpy], and naphthalene [Nap], has 2–3 rings.

Benzo[a]pyrene (BaP) is a major polycyclic aromatic hydrocarbon (PAH) with a high molecular weight. Comprising a polycyclic aromatic hydrocarbon with a chemical formula $C_{20}H_{12}$ and a molecular weight of 252.3 g.mol^{-1} , BaP develops as a side effect of incomplete combustion at temperatures between 300 and 600 degrees centigrade. At 25 degrees centigrade, BaP has a density of 1.24 g.cm^{-3} , a melting point of 179 degrees centigrade, and a boiling point of 495 degrees centigrade (Liu et al. 2019).

Mostly produced by industrial activities, Benzo[a]pyrene (BaP) results from industries releasing smoke and incomplete combustion of fossil fuels. It also results from household activities such as cooking and smoking (Sinha et al. 2005, Sun et al. 2013). Both the United States Environmental Protection Agency (EPA) and the International Agency for Research on Cancer (IARC) have labeled it as a Set 1 carcinogen. It is also a major contaminant in soil, air, and water (Bhatt et al. 2018). In the environment, BaP may undergo chemical degradation, photolysis, volatilization, and adsorption. Still, microbial degradation is mostly responsible for its loss (Haritash & Kaushik 2009).

Furthermore, affecting all kinds of biological life, it shows the process of accumulation in the food chain (Haritash & Kaushik 2009, Okpishi et al. 2017). Petroleum pollutants are easy to form mucous membranes on the surface of plant roots, obstructing plant root respiration and nutrient absorption and even causing root rot and plant death in severe cases (Yang et al. 2024). The current investigation aimed to explore the ability of green algae *S. quadricauda* in the bioremediation of various Benzo[a]pyrene amounts from contaminated media and used some physiological and biochemical parameters as tools for the bioremediation process.

MATERIALS AND METHODS

Algal Strain

Chlorophyceae species *Scenedesmus quadricauda* originally from the Algae Culture Collection (UTEX) at the University of Texas in Austin, Texas, USA. *S. quadricauda* was cultivated under controlled circumstances with a light intensity of $286 \mu\text{E.m}^{-2}.\text{s}^{-1}$, a light/dark cycle of 16:8 hours, and a temperature of 25 ± 2 degree centigrade (Chia et al. 2013), to prevent grouping and hasten their rate of expansion, the civilizations were also gently moved by hand every day (Selvan et al. 2013).

Preparation of Algal Culture Media

The improvement of chlorophyta (Patel et al. 2015) has come from the BG-11 medium. Autoclave the BG-11 medium at 121 degrees Celsius for 15 minutes to sterilize it. After freezing to try to prevent too strong precipitation, using a pH meter revealed that the pH of the medium changed to around 7.4 using 1N NaOH and 1N HCl solutions. As Gour et al. (2014) report, a 10 mL algal culture is placed in a flask holding 100 mL of BG11 medium and allowed to flourish for 15 days. Following that, this culture is placed in 1000 mL of media and allowed to incubate for two weeks minimum before beginning the experiment, therefore ensuring continuous lab conditions. Photoperiod greatly affects cell

division in asexual reproduction. Constant illumination accentuates this division that occurs all of the light time (Salman et al. 2023). Therefore, the period of light exposure may vary based on the objectives of the farming: a light-dark cycle mirroring the natural solar cycle favors regular and vigorous growth, while continuous lighting stimulates fast development (Sánchez-Bayo et al. 2020, Allen 2003).

Experimental Design

S. quadricauda's culture medium was exposed to different benzo (a) pyrene (0.5, 1, and 1.5) mM dosages for 15 days. The research primarily aims to track daily variations in algae during fifteen days. To determine the time to doubling (G) and the rate of growth (k), the algae are incubated. On the first, eighth, and fifteenth days assessments also include chlorophyll a and b, total protein, and many enzymatic responses, including reactive oxygen species (ROS), lipid peroxidation MDA, superoxide dismutase (SOD), and catalase (CAT).

Determination of Growth Rate and Doubling Time

Every day, the cell count of green algae *S. quadricauda* is determined using a UV-Vis spectrophotometer. This is accomplished by measuring the optical density (OD) at 685 nm, therefore enabling the cell density in cells per milliliter. This measurement clarifies many development stages, lets us estimate the biomass size, and ascertain growth rates. Especially, the formula of Richmond (2004) helps one find the doubling time and growth rate during the exponential development phase.

$$K = 3.322 \times (\log OD_t - \log OD_0)/t$$

$$G = 0.301/K$$

K: growth rate, G: doubling time, t: time

OD₀: optical density at the experiment beginning (zero time).

OD_t: optical density after (t) day.

Estimation of Chlorophyll

A technique detailed in Ref. (Juneja et al. 2013) was used to measure the total chlorophyll concentration. A 2 mL sample was taken and then subjected to a centrifugal force of 12,500 revolutions per minute for five minutes. The solid residue left behind from the algae was then mixed with 2 mL of methanol (90%). After an incubation period of twenty hours in a dark atmosphere at a temperature of twenty degrees, the sample was submerged in a water bath at a temperature of 64°C for five minutes. At last, the material was discarded after five minutes of centrifugation at 12,500 rpm. Spectrophotometry at three separate wavelengths— 470 nm, 652 nm, and 665 nm—quantified the filtrate.

The chlorophyll content was determined by the following Equations (1) – (2) (Lin et al. 2013):

$$\text{Chlorophyll a (mg.L}^{-1}\text{)} = (12.7 \times A663) - (2.698 \times A645) \dots(1)$$

$$\text{Chlorophyll b (mg.L}^{-1}\text{)} = (22.9 \times A645) - (4.68 \times A663) \dots(2)$$

Estimation of Reactive Oxygen Species (ROS)

Taking a 2 mL specimen, the estimate technique entails two times washing it under a Phosphate-buffer solution. The resultant precipitate-which comprises algal cells-is then combined with two milliliters of perchloric acid (200 micromol). For three minutes, the cells break down under sonication with pauses every twenty seconds. The specimen is then rapidly thrown by half an hour of spinning it at 10,000 r.m⁻¹. The volume is set at 5 mL after the filtrate is mixed using a micropipette. One 1.5 mL extract is then gathered and combined with 0.1 mL working solution. The working solution consists of 0.28 mL of sulfuric acid, 19.6 mg of ammonium ferrous sulfite, 14.3 mg of xylenol, and 3.64 g of sorbitol. The sample is kept for half an h at 30 degrees Celsius under incubation (Salman et al. 2023). The sample is next evaluated at 560 nm, and the equation helps one to ascertain the quantity of ROS:

$$\% \text{inhibition ROS} = \Delta \text{uninhibited} / \text{Min}(10)$$

Antioxidant Enzymes

The process consisted of discarding 2 mL of the specimen at a speed of 4000 – 6000 r.m⁻¹. After washing the resultant algal precipitate in a buffer solution at pH 7, the discarding process was repeated. The volume was then changed back to 2 mL. The algae cells were then disturbed for three minutes using an ultrasonic gadget called Sonication. Every 20 seconds the model pauses, then is thrown at 10,000 rpm for 30 min. The resultant filtrate is then mixed in small quantities, and the volume is changed to 5 mL (Bradford 1976).

Estimation of Carbohydrates

Following a phosphate-buffer solution wash and sonication-assisted breakdown, a 2 mL sample of the material was removed and aerobically dried. Then, using distilled water, the resultant solution was diluted to a final volume of 5 mL. 1 mL sample of the substance was then mixed with 1 milliliter of phenol (5%) and five milliliters of sulfuric acid (96%); the mixture was then swirled for ten minutes nonstop. The sample then spent thirty minutes submerged in a water bath set at 30-35°C. After that, it was evaluated at 490 nm and matched the known glucose standard curve produced by dissolving 100 mg of glucose in 100 mL of distilled water (Herbert 1971).

Estimation of Total Protein

Several methods—including centrifugation, dialysis, chromatography, precipitation, and ultrafiltration—have been used to separate and concentrate microalgal proteins (Salman et al. 2023). Based on the updated method by Ermis et al. (2020) based on the original procedure by Lowry et al. (1951), the total protein content was determined. The operation included mixing 2 mL of Biuret solution with 0.5 mL of the already produced extract (2–4). The liquid was then thoroughly stirred for thirty minutes using a preheater set to 30 degrees centigrade. The sample was then evaluated at 555 nm and matched with the reference solution depending on the Bovine Serum Albumin Protein level ranging from 0 to 0.1 mL. Dissolving 0.1 g of Bovine Serum Albumin in 100 mL of buffer solution produced a quantity of $100 \mu\text{g.L}^{-1}$, the reference solution.

Estimation of Superoxide dismutase (SOD)

The estimate calls for mixing two milliliters of Tris buffer with one hundred microliters of the extract (2–4). The first reading (ΔA_0) is then obtained by measuring the absorbance at 420 nm after five minutes. After adding 0.2 mL of Pyrogallol solution, the absorbance is once more measured at 420 nm following five minutes to get the second reading (ΔA_1). ΔA_1 is computed utilizing the formula by Zhao (2017) at hand.

SOD activity (μmL) = $[(\Delta A_0 - \Delta A_1 / \Delta A_0) / 50\%] \times \text{Volume of sample}$

Estimation of Catalase (CAT)

The estimate calls for blending one milliliter of hydrogen peroxide with one hundred microliter of the extract (2–4). The combination is then ascertained at 240 nm of wavelength. Ten minutes later, the measurement is taken once again after hydrogen peroxide addition. Apply Frary et al. (2010) formula to get the catalyst's quantity.

$\text{CAT} = [(\Delta \text{Abs}_{240} / \text{Min}_{10}) \times (\text{Reaction volume}_{100})] / 0.001$

Lipid Peroxidation (Malondialdehyde MDA)

The quantification of lipid peroxidation was conducted using the Thiobarbituric acid method to measure the quantity of Malondialdehyde (MDA), as described by Aust (1985) and Burtis and Ashwood (1999). 100 μL specimens were combined with 1000 μL of TCA (20%) and 1000 μL of TBA (0.6%). Tubes are agitated by vortexing. The sample was heated in a water bath at a temp of 100 degrees centigrade for 15 min. Afterward, it was allowed to drop down to room temp. The sample was then subjected to centrifugation at a speed of 4000 revolutions per minute

(4 degrees centigrade) for 15 min. Finally, the absorbance of the sample was measured at a wavelength of 532 nm, using a blank as a reference. The findings were quantified in terms of millimoles per milligram of protein ($\text{mmol.L}^{-1} \text{mg}^{-1} \text{protein}$).

Amount of Malondialdehyde (MDA) = Absorbance at 532nm / E \times b

Where,

E = Extinction coefficient (153mmol/cm),

b = light bath (1cm)

Estimation of Benzo[a]Pyrene

Briefly, after a 7-day incubation period, the cultures were filtered, and the B(a)P compound was extracted from the specimens using an ultrasonic bath using a solvent combination of acetonitrile and dichloromethane (at a ratio of 3:1 v/v). A Dionex UltiMate 3000 system and a fluorescence detector were used in high-performance liquid chromatography (HPLC) to determine the B(a)P amount. Benzo(a)pyrene set the excitation wavelength ($\lambda_{\text{ex}}\%$) at 250 nm and the emission wavelength ($\lambda_{\text{em}}\%$) at 400 nm. Chromatography with a gradient under the use of a mobile phase comprising a combination of water and acetonitrile was the separation method used. Thermo Scientific HYPERSIL GOLD C18 PAH column with dimensions of 250 x 4.6 mm and particle size of 5 μm was the chromatographic column utilized. Made by Merc, the solvents used in the tests were of high-performance liquid chromatography (HPLC) grade. Calibrating curves with magnitudes ranging from 0.1 to 10 ng.cm^{-3} were generated using the Sigma-Aldrich benzo(a) pyrene standard at $1000 \mu\text{g.mL}^{-1}$. The standard solvent used in the construction of the calibration curves was methanol. The method demonstrated a linearity higher than 0.999%. Still, the coefficient of variance revealed a level of accuracy below 15%. With a signal-to-noise proportion of 10 for a specimen with a very low amount of B(a)P, approaching the detection limit, the limit of quantification (LoQ) for the method was found. Measuring the LoQ magnitude, we found 0.01 ng.cm^{-3} . The correlation between the recovery rate and the reference material (SRM2585) was 83%. Once used to find B(a)P in the atmosphere, this approach Nine strains in all were exposed to various dosages of B(a)P (7.8, 15, 78, 312, and 624 ng.L^{-1}) and temperatures (10, 15, 20, 25, and 30 degree centigrade) in Wiśniewska et al. (2023). The experiment also included blank specimens.

Statistical Analysis

For every treatment, the experimental data in this work came from a minimum of three replicas. SPSS version 26 was used for statistical investigation. The data were expressed

as mean and standard deviation. The relevance of variances was evaluated using an analysis of variance (ANOVA). If the related p-magnitudes were less than 0.05, variations were judged noteworthy.

RESULTS AND DISCUSSION

Growth Rate and Doubling Time

Several studies have demonstrated the effect of exposure to PAHs on algal growth rate and doubling time. This study showed that different amounts of benzo[a]pyrene inhibited the growth rate while increasing the doubling time to a long period slowed cell division. There is a significant variance ($p \leq 0.05$) between the various treatments of Benzo[a]pyrene (0.5, 1, 1.5) mM, the present study demonstrates, the maximum growth rate (k) at the amount of (0.5) mM was 0.391 throughout 7th day, while the minimum growth rate was 0.00047 at the amount (1.5) mM throughout 15th day. The shorter doubling time (G) was 0.00014 during 15th day with (1.5)mM, while the longest value 0.1179 during 7th day with (0.5) mM benzo (a) pyrene (Fig. 1,2). The inhibitory impact of toxicity could be attributed to the extremely lipophilic characteristics of Benzo[a]pyrene, which enables it to easily

enter cells via the plasma membrane and rapidly incorporate into all biological lipoprotein membranes (Miller & Ramos 2001). It induces alterations in the permeability and structure of membranes and disrupts crucial metabolic activities, such as photosynthesis, respiration, and transmembrane transport (Sikkema et al. 1994, Tripuranthakam et al. 1999, Aksmann & Tukaj 2004). Calow & Sibly (1990) proposed that various forms of toxic stress cause metabolic alterations in organisms, depleting their energy reserves and negatively impacting their growth and biochemical composition. This finding was reported by Echeveste et al. (2010), who noted a reduction in algae growth when exposed to high amounts of PAHs.

Algal Pigments

In the study, benzo[a]pyrene treatments at various concentrations led to a reduction in chlorophyll a and b in *S. quadricauda*. The results show chlorophyll a in cell *S. quadricauda* treatment with different concentrations of Benzo[a]pyrene decreased to $0.41 \pm (\mu\text{g} \cdot \text{mL}^{-1})$ in 0.5 mM during 1st day, while the high value was $18.71 \pm (\mu\text{g} \cdot \text{mL}^{-1})$ in 1.5 mM during 15th day, The results show chlorophyll b decrease to $0.36 \pm (\mu\text{g} \cdot \text{mL}^{-1})$ in 1.5 mM during 1st day, while




Fig. 1: Growth rate of *S. quadricauda* at concentration of benzo (a) pyrene.




Fig. 2: Doubling time of *S. quadricauda* at various concentrations of benzo (a) pyrene.




Fig. 3: Chlorophyll an amount of *S. quadricauda* at various concentrations of Benzo[a]pyrene .




Fig. 4: Chlorophyll b amount of *S. quadricauda* at various concentrations of Benzo [a] pyrene.

the high value was $9.19 \pm (\mu\text{g} \cdot \text{mL}^{-1})$ in 1.5 mM during 15th day compared to control (Fig. 3,4) The synthesis of pigments may be inhibited by the lipophilic nature of aromatic hydrocarbons, which can alter the permeability and fluidity of the cell membrane. This alteration could disrupt the lipid bilayer and reduce energy transduction, affecting the activity of membrane-associated proteins (Heipieper et al. 1995). Mohapatra & Schiewer (2000) suggested that the interaction of toxicants with the membrane and dispersion of lipoprotein membranes are responsible for changes in behavior and pigment content in *Synechocystis*. Mostafa & Helling (2002) proposed that the decrease in chlorophyll, carotenoid, and phycobiliprotein contents might be a result of direct inhibition of pigment synthesis by the organic compound or accelerated degradation of pigments since increased reactive oxygen species (ROS) and degradation of the thylakoid membrane by polycyclic aromatic hydrocarbons (PAHs).

Reactive Oxygen Species (ROS)

Benzo [a] pyrene-induced oxidative stress for cell algae in

treated cultures, leading high amount of ROS. where the highest value of ROS is 0.028 % throughout the 15th day with 1.5 mM of Benzo[a]pyrene compared to the control (Fig. 5). A notable disparity was observed between the treatments and control culture. The hydrophobic feature of Benzo[a]pyrene leads to the overproduction of ROS, which in turn causes alterations in the structure of biomembranes, resulting in an elevated permeability (Sikkema et al. 1994, Solomon 2000). As a result, significant disruptions in the electron transport chains and/or the process of phosphorylation uncoupling may occur (Aksmann & Tukaj 2008, Aksmann et al. 2011). An abundance of intracellular ROS can result in the oxidation of macromolecules, such as lipids, proteins, and DNA bases (Meewes et al. 2001). This oxidative process can lead to damage in cell membranes, mitochondria, and chloroplasts, ultimately inhibiting photosynthesis, physiological activities, and cell growth. In response to the assault of ROS, cells have created enzyme systems that act as antioxidant defense mechanisms to eliminate ROS (Takáčová et al. 2014).




Fig. 5: ROS amount of *S. quadricauda* at various concentrations of Benzo[a]pyrene.

Carbohydrate

Carbohydrate is synthesized by photosynthetic organisms like algae and are necessary to obtain energy. All Environmental toxicants have effects on the content of carbohydrates in algae by reducing carbohydrate synthesis; perhaps stress also increases respiration, which then uses up the carbohydrates that were synthesized previously (Mustafa 2013). The carbohydrate amount of $12.25 \mu\text{m.mL}^{-1}$ is recorded at 1.5 mM on the 15th day. The amount of Benzo[a]pyrene is $15.52 \mu\text{m.mL}^{-1}$ and $16.30 \mu\text{m.mL}^{-1}$ for 0.5 mM and 1 mM, respectively. The greatest amount of $35.13 \mu\text{m.mL}^{-1}$ is seen on the 1st day with 1 mM (Fig. 6). Both Nagajyoti et al. (2009) and Abdul Razak (1985) demonstrated that the decrease in carbs may be attributed to the suppression of Ribulose Biphosphate carboxylase (RUBP) activity, leading to a fall in carbohydrate levels. RUBP carboxylase is the predominant enzyme in photosynthesis and plays a crucial role in assimilating carbohydrates and regulating carbohydrate metabolism. It is involved in important

metabolic processes, including fixation of carbon and glycolysis (Barsanti & Gualtieri 2006). The toxicant can hinder the process of glucose oxidation metabolism, which is necessary for obtaining ATP and reducing metabolic activity inside the cell (Greenhaff et al. 2004). From the findings of this study, the treatment of *S. quadricauda* with various Benzo[a]pyrene concentrations caused a significant decrease in total carbohydrate content with increasing concentrations of Benzo [a] pyrene.

Total Protein

Benzo[a]pyrene stress has an effect on the content of proteins in all treatments, and it causes a decrease in total protein with increasing amounts of Benzo [a] pyrene. The max protein content of $2.14 \mu\text{g.mL}^{-1}$ throughout the 8th day is recorded at 1mM, while the min protein content of 1.83 mM is recorded at 1.5 mM amount throughout the 8th day (Fig. 7). The reduction in protein synthesis can be attributed to the inhibition of enzymes and structural proteins necessary




Fig. 6: carbohydrate amount of *S. quadricauda* at various concentrations of Benzo [a] pyrene.




Fig. 7: Total protein amount of *S. quadricauda* at various concentrations of Benzo [a] pyrene.

for the organism's growth (Kapoor & Arora 1996). Additionally, Carfagna et al. (2013) suggest that the decrease in protein content may also be caused by a shortage of carbon skeleton resulting from a low photosynthetic rate. Sub-lethal doses of PAHs may disrupt cell division by inhibiting DNA synthesis, lowering the proportion of cells entering mitosis, and subsequently limiting population expansion and protein production (Carfagna et al. 2013, Chia et al. 2015).

Superoxide Dismutase (SOD)

In the present study, it can be observed clear effect of various Benzo[a]pyrene amount in the activity of the enzyme (SOD) in alga *S. quadricauda* indicate a significant variance ($p \leq 0.05$) between the treatments and control. There was increasing in content SOD at Benzo[a]pyrene 60 µg.mL⁻¹ in the amount of 1.5 mM throughout the 8th day, compared with control (26.66 µg.mL⁻¹), where the lowest magnitude at (0.5 mM) was 22.22 µg.mL⁻¹ throughout 1st day (Fig. 8). When the quantity of reactive oxygen species (ROS) in microalgal cells becomes very high, the cells'

capacity to eliminate ROS by creating sufficient amounts of oxidative response kinases, such as SOD, is compromised. Prior research by Papadimitriou and Loumbourdis (2002) indicated that some species may counteract oxidative stress by boosting the synthesis of SOD to remove excess ROS. SOD is a crucial defensive mechanism that regulates the levels of free radicals. It is recognized as the primary defense against ROS in live cells. The elevated level of SOD activity leads to the increased breakdown of harmful ROS, such as the superoxide radical (O_2^-), into a less toxic ROS, such as hydrogen peroxide (H_2O_2). This hydrogen peroxide may then be further reduced by the heightened activity of glutathione reductase (GR), as seen in the study by Samanta et al. (2019). However, the decreased functioning of SOD in the presence of high amounts of PAHs is probably due to the excessive accumulation of H_2O_2 (Binark et al. 2000).

Catalase (CAT)

In the current research, it can be observed clear effect of various Benzo[a]pyrene amount in the activity of the enzyme




Fig. 8: SOD amount of *S. quadricauda* at various concentrations of Benzo [a] pyrene.

(CAT) in alga *S. quadricauda* and indicate the significant variance ($p \leq 0.05$) between the control and treatments. There was increasing in the content of catalase at Benzo[a] pyrene $73.33 \mu\text{g.mL}^{-1}$ in concentration 1 mM during the 15th day, compared with the control ($10 \mu\text{g.mL}^{-1}$). Where the lowest magnitude at (1 mM) was $13.33 \mu\text{g.mL}^{-1}$ throughout the 8th and 15th day (Fig. 9). Like other living creatures, algae create several ROS as a result of oxidative metabolism, such as H_2O_2 , which may be harmful at high quantities (Torres et al. 2010). The formation of these ROS is sluggish under typical circumstances. Nevertheless, contamination, especially PAHs, enhances their creation (Binark et al. 2000, Torres et al. 2010). CAT is an important antioxidant enzyme that plays a crucial role in regulating the redox state inside cells. Its primary function is to protect cells from damage caused by peroxidation. This has been demonstrated in studies conducted by Cheng et al. (2016) and Hassan et al. (2017, 2018). The enzyme catalase (CAT) may mitigate the harmful effects of hydrogen peroxide (H_2O_2) by facilitating its transformation into water (H_2O) (Torres et al. 2010). The

excessive generation of ROS may diminish the functioning of CAT, leading to oxidative stress (El Maghraby & Hassan 2021).

Lipid Peroxidation (Malondialdehyde MDA)

Lipids, being crucial constituents of the membrane, are the main target of ROS and result in lipid peroxidation by the extraction of hydrogen from the unsaturated fatty acid chain. Thus, lipid peroxidation plays a crucial role in causing cellular damage in living organisms experiencing oxidative stress. Lipid peroxidation generates many harmful compounds, including malondialdehyde (MDA) and aldehydes, from polyunsaturated fatty acids. MDA is a marker for oxidative damage. Algae possess chloroplasts that include a complex network of membranes rich in polyunsaturated fatty acids. These fatty acids are particularly susceptible to peroxidation (Garg & Manchanda 2009, Halliwell & Chirico 1993).

The findings study demonstrates a significant variance ($p \leq 0.05$) in the content of the MDA for the alga




Fig. 9: CAT amount of *S. quadricauda* at various amounts of Benzo [a] pyrene.




Fig. 10: MDA amount of *S. quadricauda* at various amounts of Benzo [a] pyrene.

S. quadricauda after treatment; the content of MDA decreased at various amounts Benzo[a]pyrene compared with control ($34.14 \mu\text{mol.L}^{-1}$), the minimum value is 36.58, 39.02 and $42.50 \mu\text{mol.L}^{-1}$ at concentration 1, 0.5 and 1.5 mM respectively, throughout 15th day (Fig. 10). This may be due to stimulated antioxidant defense in algae was efficient in eliminating ROS, and inhibiting the algal cells from oxidative damage (Zhang et al. 2018).

Bioremediation of Benzo[a]pyrene

Researchers have extensively investigated the capacity of certain types of algae to break down PAH (polycyclic aromatic hydrocarbon) compounds found in water due to their toxicity and carcinogenic properties. In a study conducted by Kumar et al. (2018), it was discovered that PAHs interact with cytochrome P450 monooxygenase CYP in the active sites of algae through intermolecular hydrogen bonding, hydrophobic bonding π - π interactions, and van der Waals interactions. These interactions enable algae to remediate PAHs more effectively than other microorganisms. The results showed the effect of different amounts of benzo[a]pyrene on the degradation of *S. quadricauda*; the highest value was 40.13 throughout the 15th with 0.5 mM, while the lowest value was 0 throughout 1st day with 0.5 and 1 mM (Fig. 11), in the present study demonstrated *S. quadeicauda* have high ability to removed Benzo[a]pyrene after 1, 8 and 15 days of treatments. Ei-Sheekh et al. (2012) suggested that the chemical structure of the molecule and the physiological metabolism of the algae appear to determine how polycyclic and heterocyclic aromatic compounds break down under them. Apart from being ingested by cells or absorbed on the wall, the results of the study imply that algae broke down the compound's high removal efficiencies by several processes, including a reduction, oxidation, and the activation

of enzymes, including cytochrome P450, diphenol oxidase, and Peroxidase, Detoxifying B[a]P and breaking down these toxic chemicals depend critically on these enzyme systems (Takáčová et al. 2014).

CONCLUSIONS

Benzo[a]pyrene, a form of polycyclic aromatic hydrocarbon (PAH), may be efficiently removed from polluted water by use of the green alga *S. quadricauda* in the process of bioremediation. This results from harmless substances being produced all along the algal breakdown. The results of the investigation show that the alga shows great ability for BaP removal at low concentrations and that the pollution is cleared in a short period. Algae showed biological reactions in some results of exposure to different biological parameters with different amounts of B[a]P such as decreased amount, total protein, chlorophyll a, and b, increased oxidative stress ROS and carbohydrates and enzyme activity, SOD, MDA, and catalase.

ACKNOWLEDGMENT

The authors extend their thanks and appreciation to the Department of Biology, College of Science at the University of Babylon, for their assistance in supporting this work.

REFERENCES

- Abdul Razak, V., 1985. Physiological and Biochemical Aspects of Metal Tolerance in *Arachis hypogaea*. LM Phil. Dissertation, SV University, Tirupati. Abstr. No. XIV-25.
- Aksmann, A., Shutova, T., Samuelsson, G. and Tukaj, Z., 2011. The mechanism of anthracene interaction with photosynthetic apparatus: A study using intact cells, thylakoid membranes, and PS II complexes isolated from *Chlamydomonas reinhardtii*. *Aquatic Toxicology*, 104(3-4), pp.205-210.




Fig. 11: HPLC Chromatogram for Benzo[a]pyrene standard at various concentrations.

- Aksmann, A. and Tukaj, Z., 2004. The effect of anthracene and phenanthrene on the growth, photosynthesis, and SOD activity of the green alga *Scenedesmus armatus* depends on the PAR irradiance and CO₂ level. *Archives of Environmental Contamination and Toxicology*, 47, pp.177-184.
- Aksmann, A. and Tukaj, Z., 2008. Intact anthracene inhibits photosynthesis in algal cells: A fluorescence induction study on *Chlamydomonas reinhardtii* cw92 strain. *Chemosphere*, 74(1), pp.26-32.
- Alegbeleye, O.O., Opeolu, B.O. and Jackson, V.A., 2017. Polycyclic aromatic hydrocarbons: A critical review of environmental occurrence and bioremediation. *Environmental Management*, 60, pp.758-783.
- Alexander, M., 1999. *Biodegradation and Bioremediation*. 2nd ed. Academic Press.
- Allen, J.F., 2003. State transitions—A question of balance. *Science*, 299(5612), pp.1530-1532.
- Amran, R.H., Jamal, M.T., Pugazhendhi, A., Al-Harbi, M., Ghandourah, M., Al-Otaibi, A. and Haque, M.F., 2022. Biodegradation and bioremediation of petroleum hydrocarbons in marine ecosystems by microorganisms: A review. *Nature Environment & Pollution Technology*, 21(3).
- Aust, S.D., 1985. *Handbook of Methods for Oxygen Radical Research*. Springer, pp.203-207.
- Barsanti, L. and Gualtieri, P., 2022. *Algae: Anatomy, Biochemistry, and Biotechnology*. CRC Press.
- Bhatt, K.K., Lily, M.K., Joshi, G. and Dangwal, K., 2018. Benzo(a)pyrene degradation pathway in *Bacillus subtilis* BMT4i (MTCC 9447). *Turkish Journal of Biochemistry*, 43(6), pp.693-701.
- Binark, N., Güven, K.C., Gezgin, T. and Ünlü, S., 2000. Oil pollution of marine algae. *Bulletin of Environmental Contamination & Toxicology*, 64(6).
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), pp.248-254.
- Burtis, C.A. and Ashwood, E.R., 1999. *Tietz Textbook of Clinical Chemistry*. Philadelphia, pp.1654-1655.
- Calow, P. and Sibly, R.M., 1990. A physiological basis of population processes: Ecotoxicological implications. *Functional Ecology*, 18, pp.283-288.
- Carfagna, S., Lanza, N., Salbitani, G., Basile, A., Sorbo, S. and Vona, V., 2013. Physiological and morphological responses of lead- or cadmium-exposed *Chlorella sorokiniana* 211-8K (*Chlorophyceae*). *SpringerPlus*, 2, pp.1-7.
- Cheng, J., Qiu, H., Chang, Z., Jiang, Z. and Yin, W., 2016. The effect of cadmium on the growth and antioxidant response for freshwater algae *Chlorella vulgaris*. *SpringerPlus*, 5, pp.1-7.
- Chia, M.A., Chimdirim, P.K. and Japhet, W.S., 2015. Lead-induced antioxidant response and phenotypic plasticity of *Scenedesmus quadricauda* (Turp.) de Brébisson under different nitrogen concentrations. *Journal of Applied Phycology*, 27, pp.293-302.
- Chia, M.A., Lombardi, A.T. and Melao, M.D.G.G., 2013. Growth and biochemical composition of *Chlorella vulgaris* in different growth media. *Anais da Academia Brasileira de Ciências*, 85(4), pp.1427-1438.
- Echeveste, P., Agustí, S. and Dachs, J., 2010. Cell size-dependent toxicity thresholds of polycyclic aromatic hydrocarbons to natural and cultured phytoplankton populations. *Environmental Pollution*, 158(1), pp.299-307.
- El Maghraby, D. and Hassan, I., 2021. Photosynthetic and biochemical response of *Ulva lactuca* to marine pollution by polyaromatic hydrocarbons (PAHs) collected from different regions in Alexandria City, Egypt. *Egyptian Journal of Botany*, 61(2), pp.467-478.
- Ei-Sheekh, M.M., Ghareib, M.M. and El-Souod, G.W., 2012. Biodegradation of phenolic and polycyclic aromatic compounds by some algae and cyanobacteria. *Journal of Bioremediation & Biodegradation*, 3(133), p.2.
- Ermis, H., Guven-Gulhan, U., Cakir, T. and Altinbas, M., 2020. Effect of iron and magnesium addition on population dynamics and high-value product of microalgae grown in the anaerobic liquid digestate. *Scientific Reports*, 10(1), p.3510.
- Frary, A., Göll, D., Keleş, D., Ökmen, B., Pınar, H., Şığva, H.Ö., Yemencioğlu, A. and Doğanlar, S., 2010. Salt tolerance in *Solanum pennellii*: Antioxidant response and related QTL. *BMC Plant Biology*, 10, pp.1-16.
- García de Llasera, M.P., Olmos-Espejel, J.D.J., Díaz-Flores, G. and Montaña-Montiel, A., 2016. Biodegradation of benzo(a)pyrene by two freshwater microalgae *Selenastrum capricornutum* and *Scenedesmus acutus*: A comparative study useful for bioremediation. *Environmental Science and Pollution Research*, 23, pp.3365-3375.
- Garg, N. and Manchanda, G., 2009. ROS generation in plants: Boon or bane? *Plant Biosystems*, 143(1), pp.81-96.
- Gour, R.S., Kant, A. and Chauhan, R.S., 2014. Screening of microalgae for growth and lipid accumulation properties. *Journal of Algal Biomass Utilization*, 5, pp.38-46.
- Halliwell, B. and Chirico, S., 1993. Lipid peroxidation: Its mechanism, measurement, and significance. *The American Journal of Clinical Nutrition*, 57(5), pp.715S-725S.
- Haritash, A.K. and Kaushik, C.P., 2009. Biodegradation aspects of polycyclic aromatic hydrocarbons (PAHs): A review. *Journal of Hazardous Materials*, 169(1-3), pp.1-15.
- Greenhaff, P.L., Hultman, E. and Harris, R.C., 2004. Carbohydrate metabolism. *Medicine and Sport Science*, 46, pp.108-151.
- Hassan, I.A., Bell, N.B., Ashmore, M.R., Cotrozzi, L., Haiba, N.S., Basahi, J.M., Summan, A., Almeelbi, T. and Ismail, I.M., 2018. Radish (*Raphanus sativus* L.) cultivar-specific response to O₃: Patterns of biochemical and plant growth characteristics. *Clean-Soil, Air, Water*, 46(6), p.1800124.
- Hassan, I.A., Haiba, N.S., Badr, R.H., Basahi, J.M., Almeelbi, T., Ismail, I.M. and Taia, W.K., 2017. Effects of ambient ozone on reactive oxygen species and antioxidant metabolites in leaves of pea (*Pisum sativum* L.) plants. *Pakistani Journal of Botany*, 49(1), pp.47-55.
- Heipieper, H.J., Löffel, B., Keweloh, H. and de Bont, J.A., 1995. The *cis/trans* isomerization of unsaturated fatty acids in *Pseudomonas putida* S12: an indicator for environmental stress due to organic compounds. *Chemosphere*, 30(6), pp.1041-1051.
- Herbert, D., Phipps, P.J. and Strange, R.E., 1971. *Methods in Microbiology*. Academic Press, pp. 209-344.
- Hwang, J.H., Church, J., Lee, S.J., Park, J. and Lee, W.H., 2016. Use of microalgae for advanced wastewater treatment and sustainable bioenergy generation. *Environmental Engineering Science*, 33(11), pp.882-897.
- Juda, S.A., Salah, M.M. and Salman, J.M., 2019. Efficiency of green algae *Chlorella vulgaris* in remediation of polycyclic aromatic hydrocarbon (anthracene) from culture media. *Baghdad Science Journal*, 16(3), pp.543-549.
- Juneja, A., Ceballos, R.M. and Murthy, G.S., 2013. Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: a review. *Energies*, 6(9), pp.4607-4638.
- Kapoor, K. and Arora, L., 1996. Observations on growth responses of cyanobacteria under the influence of herbicides. *Pollution Research*, 15, pp.343-351.
- Keith, L.H., 2015. The source of US EPA's sixteen PAH priority pollutants. *Polycyclic Aromatic Compounds*, 35(2-4), pp.147-160.
- Kumar, P.S., Thomas, J. and Poornima, V., 2018. Structural insights on bioremediation of polycyclic aromatic hydrocarbons using microalgae: A modeling-based computational study. *Environmental Monitoring and Assessment*, 190(92), pp.1-16.
- Lawal, A.T., 2017. Polycyclic aromatic hydrocarbons: A review. *Cogent Environmental Science*, 3(1), p.1339841.

- Lin, Y.H., Pan, K.Y., Hung, C.H., Huang, H.E., Chen, C.L., Feng, T.Y. and Huang, L.F., 2013. Overexpression of ferredoxin, PETF, enhances tolerance to heat stress in *Chlamydomonas reinhardtii*. *International Journal of Molecular Sciences*, 14(10), pp.20913-20929.
- Liu, H., Yin, H., Tang, S., Wei, K., Peng, H., Lu, G. and Dang, Z., 2019. Effects of benzo[a]pyrene (BaP) on the composting and microbial community of sewage sludge. *Chemosphere*, 222, pp.517-526.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J., 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193(1), pp.265-275.
- Luka, Y., Highina, B.K. and Zubairu, A., 2018. Bioremediation: A solution to environmental pollution: A review. *American Journal of Engineering Research*, 7(2), pp.101-109.
- McCann, J.H. and Solomon, K.R., 2000. The effect of creosote on membrane ion leakage in *Myriophyllum spicatum* L. *Aquatic Toxicology*, 50(3), pp.275-284.
- Meewes, C., Brenneisen, P., Wenk, J., Kuhr, L., Ma, W., Alikoski, J., Poswig, A., Krieg, T. and Scharffetter-Kochanek, K., 2001. Adaptive antioxidant response protects dermal fibroblasts from UVA-induced phototoxicity. *Free Radical Biology and Medicine*, 30(3), pp.238-247.
- Miller, K.P. and Ramos, K.S., 2001. Impact of cellular metabolism on the biological effects of benzo[a]pyrene and related hydrocarbons. *Drug Metabolism Reviews*, 33(1), pp.1-35.
- Mohapatra, P.K. and Schiewer, U., 2000. Dimethoate and quinalphos toxicity: Pattern of photosynthetic pigment degradation and recovery in *Synechocystis* sp. PCC 6803. *Archiv für Hydrobiologie. Supplementband, Algological Studies*, 134, pp.79-94.
- Mostafa, F.I. and Helling, C.S., 2002. Impact of four pesticides on the growth and metabolic activities of two photosynthetic algae. *Journal of Environmental Science and Health, Part B*, 37(5), pp.417-444.
- Mustafa, E.M., 2013. *Effect of Selected Chemical Contaminants on Growth, Biochemical Composition, DNA Damage, and Superoxide Dismutase Activity in Four Marine Algae*. University of Malaya (Malaysia).
- Nagajyoti, P.C., Dinakar, N., Suresh, S., Udaykiran, Y., Suresh, C., Prasad, T.N.V.K.V. and Damodharam, T., 2009. Biomass industrial effluent effect on carbohydrates, amino acids, nitrite, and nitrite enzyme activities of *Arachis hypogaea* L. *Agricultural Sciences in China*, 8(2), pp.203-215.
- Nandal, M., Solanki, P., Rastogi, M. and Hooda, R., 2015. Bioremediation: A sustainable tool for environmental management of oily sludge. *Nature Environment and Pollution Technology*, 14(1), p.181.
- Okpashi, V.E., Ogugua, V.N., Ubani, S.C., Ujah, I.I. and Ozioko, J.N., 2017. Estimation of residual polycyclic aromatic hydrocarbons concentration in fish species: Implication in reciprocal corollary. *Cogent Environmental Science*, 3(1), p.1303979.
- Pandya, D.K. and Kumar, M.A., 2021. Chemo-metric engineering designs for deciphering the biodegradation of polycyclic aromatic hydrocarbons. *Journal of Hazardous Materials*, 411, p.125154.
- Papadimitriou, E. and Loumbourdis, N.S., 2002. Exposure of the frog *Rana ridibunda* to copper: Impact on two biomarkers, lipid peroxidation and glutathione. *Bulletin of Environmental Contamination and Toxicology*, 69(6).
- Patel, A.K., Suseela, M.R., Singh, M. and Nayaka, S., 2015. Application of response surface methodology for optimization of biomass, carbohydrate, and lipid production in BG11+ by *Scenedesmus quadricauda*. *International Journal of Research in Engineering and Applied Sciences*, 5, pp.199-215.
- Richmond, A. ed., 2004. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology* (Vol. 577). Oxford: Blackwell Science
- Salman, J.M., Grmasha, R.A., Stenger-Kovács, C., Lengyel, E., Al-Sareji, O.J., Al-Cheban, A.M.A. and Meiczing, M., 2023. Influence of magnesium concentrations on the biomass and biochemical variations in the freshwater algae *Chlorella vulgaris*. *Heliyon*, 9(1).
- Samanta, P., Shin, S., Jang, S. and Kim, J.K., 2019. Comparative assessment of salinity tolerance based on physiological and biochemical performances in *Ulva australis* and *Pyropia yezoensis*. *Algal Research*, 42, p.101590.
- Sánchez-Bayo, A., Morales, V., Rodríguez, R., Vicente, G. and Bautista, L.F., 2020. Cultivation of microalgae and cyanobacteria: Effect of operating conditions on growth and biomass composition. *Molecules*, 25(12), p.2834.
- Selvan, B.K., Revathi, M., Piriya, P.S., Vasan, P.T., Prabhu, D. and Vennison, S.J., 2013. Biodiesel production from marine cyanobacteria cultured in plate and tubular photobioreactors. *Journal of Bioenergy and Algal Research*, 15(2), pp.45-53.
- Shukla, K.P., Singh, N.K. and Sharma, S., 2010. Bioremediation: Developments, current practices and perspectives. *Genetic Engineering and Biotechnology Journal*, 3, pp.1-20.
- Sikkema, J., de Bont, J.A. and Poolman, B., 1994. Interactions of cyclic hydrocarbons with biological membranes. *Journal of Biological Chemistry*, 269(11), pp.8022-8028.
- Singh, O.V., 2006. Proteomics and metabolomics: The molecular make-up of toxic aromatic pollutant bioremediation. *Proteomics*, 6(20), pp.5481-5492.
- Sinha, R., Kulldorff, M., Gunter, M.J., Strickland, P. and Rothman, N., 2005. Dietary benzo[a]pyrene intake and risk of colorectal adenoma. *Cancer Epidemiology Biomarkers & Prevention*, 14(8), pp.2030-2034.
- Sun, Y., Xu, Y., Zhou, Q., Wang, L., Lin, D. and Liang, X., 2013. The potential of gibberellic acid 3 (GA3) and Tween-80 induced phytoremediation of co-contamination of Cd and benzo[a]pyrene (B[a]P) using *Tagetes patula*. *Journal of Environmental Management*, 114, pp.202-208.
- Takáčová, A., Smolinská, M., Ryba, J., Mackul'ak, T., Jokrllová, J., Hronec, P. and Čík, G., 2014. Biodegradation of benzo[a]pyrene through the use of algae. *Central European Journal of Chemistry*, 12, pp.1133-1143.
- Tewari, A., Bhutada, D.S. and Wadgaonkar, V., 2023. Heavy metal remediation from water/wastewater using bioadsorbents—A review. *Nature Environment & Pollution Technology*, 22(4), p.11.
- Torres, M.A., Barros, M.P., Campos, S.C., Pinto, E., Rajamani, S., Sayre, R.T. and Colepicolo, P., 2008. Biochemical biomarkers in algae and marine pollution: A review. *Ecotoxicology and Environmental Safety*, 71(1), pp.1-15.
- Tripranthakam, S., Duxbury, C.L., Babu, T.S. and Greenberg, B.M., 1999. *Environmental Toxicology and Risk Assessment: Standardization of Biomarkers for Endocrine Disruption and Environmental Assessment*. ASTM International.
- Wiśniewska, K.A., Lewandowska, A.U., Śliwińska-Wilczewska, S., Staniszevska, M. and Budzalek, G., 2023. The ability of airborne microalgae and cyanobacteria to survive and transfer the carcinogenic benzo[a]pyrene in coastal regions. *Cells*, 12(7), p.1073.
- Yang, B., Xue, Q.H., Qu, C.T., Lu, C., Liu, F.F., Zhang, H., Ma, L.T., Qi, L. and Wang, Y.T., 2024. Research progress on in-situ remediation of typical heavy metals in petroleum hydrocarbon-contaminated soil enrichment by plants. *Nature Environment & Pollution Technology*, 23(1), p.65.
- Zhang, M., Wang, X., Tao, J., Li, S., Hao, S., Zhu, X. and Hong, Y., 2018. PAHs would alter cyanobacterial blooms by affecting the microcystin production and physiological characteristics of *Microcystis aeruginosa*. *Ecotoxicology and Environmental Safety*, 157, pp.134-142.
- Zhao, L., 2018. Measurement of superoxide dismutase-like activity in peel and pulp of apple from Anshan acres. *AIP Conference Proceedings*, 1956(1), p.7521.