

RESEARCH ARTICLE

<https://doi.org/10.46488/NEPT.2026.v25i04.D1905>

OPEN ACCESS

Evaluation of Iron Ore Dust Emission Rates and Associated Bacterial Load

Melainine Aillal^{1†}, **Olfa Ben Said^{2,3}**, **Hamouda Beyrem²** and **Zeinebou Sidoumou¹**¹University of Nouakchott, Faculty of Science and Technology, New University Campus, 5026, Nouakchott, Mauritania²Laboratory of Environment Biomonitoring, Coastal Ecology Unit, Faculty of Sciences of Bizerte, University of Carthage, Zarzouna, Tunisia³Laboratory of Biotechnology and Nuclear Technology, National Centre of Nuclear Science and Technology, Sidi Thabet Technopark, 2020 Ariana, Tunisia†Corresponding author: Melainine Aillal; melainineailal@gmail.com

ARTICLE HISTORY

Received: 22-01-2026

Revised: 09-03-2026

Accepted: 19-03-2026

KEYWORDS

Mining activities
Dust emissions
Heavy metals
Environmental pollution
Staphylococcus
Occupational health

CITE THIS ARTICLE

Aillal, M., Ben Said, O., Beyrem, H. and Sidoumou, Z., 2026. Evaluation of iron ore dust emission rates and associated bacterial load. *Nature Environment and Pollution Technology*, 25(4), D1905. <https://doi.org/10.46488/NEPT.2026.v25i04.D1905>

COPYRIGHT & LICENCE

© 2026 The Authors
Open Access under CC BY 4.0.

ABSTRACT

The National Industrial and Mining Company (SNIM) is facing significant dust emissions, raising serious environmental and public health concerns. This study aimed to characterize airborne dust emissions from mining activities and to assess their physicochemical and microbiological composition. Dust samples were analyzed and found to contain hazardous heavy metals, including lead, copper, nickel, and cobalt. A standardized measurement procedure was developed and applied in accordance with French Standard X 43-014 to quantify dust emission rates.

The potential sources and causes of dust emissions were identified, and immediate preventive and corrective actions were proposed using the Failure Modes, Effects, and Criticality Analysis (FMECA) method. The results revealed exceptionally high dust emission levels at the Guelb El Rhein mine, exceeding the thresholds associated with highly polluted environments. In contrast, emissions at the Kedia d'Idjil mine remained close to regulatory limits and were comparable to levels observed in relatively livable environments.

Microbiological analysis of dust particles revealed the presence of potentially pathogenic staphylococcal species, including *Staphylococcus epidermidis*, *S. hominis*, *S. capitis*, *S. haemolyticus*, *S. lentus*, *S. auricularis*, *S. sciuri*, and *S. cohnii* subsp. *urealyticum*. These findings highlight the combined chemical and biological risks associated with mining dust emissions and underscore the need for improved dust management strategies in mining environments.

1. INTRODUCTION

Atmospheric pollution has existed for centuries, but has intensified considerably with the expansion of human activities. The large-scale production of energy through combustion processes and the rapid development of industrial operations have profoundly disrupted natural atmospheric balances, not only at local scales but also globally. These changes have led to significant degradation of air quality and have contributed to climate change (Boussahel 2007). Atmospheric emissions include not only gases and fine particulate matter but also heavy metals and various chemical compounds that can exert toxic effects on ecosystems and human health.

The mechanisms governing environmental contamination through atmospheric deposition remain insufficiently understood. During cloud formation, pollutants are partitioned among different atmospheric phases, namely gaseous, liquid, and particulate. While gas-liquid equilibrium processes have been extensively studied,

interactions between particulate and liquid phases are still poorly documented. Recent studies have demonstrated that the pollutant load varies according to particle size, which strongly influences particle transport, deposition processes, and environmental fate (Ma et al. 2014, Hassanvand et al. 2015). Fine particles are capable of long-range transport, whereas coarse particles tend to deposit more rapidly, leading to pronounced local impacts on air, soil, and water quality.

Air pollution remains one of the most critical global environmental challenges and continues to intensify due to rapid urbanization and the expansion of industrial and mining activities. This situation underscores the urgent need to control emissions to protect both public health and the environment (Elodie et al. 2020). In addition to chemical contamination, air, particularly indoor air in industrial settings, can serve as a favorable medium for the development and persistence of pathogenic microorganisms associated with dust and aerosols (Viegas et al. 2020, Yan et al. 2021). Previous research has shown that airborne dust and bioaerosols can significantly increase occupational health risks, including infections, toxic effects, and allergic responses (Eduard et al. 2012, Heederik & Von Mutius 2012).

In Mauritania, the discovery and exploitation of mineral resources over the past two decades have led to the rapid expansion of mining projects, predominantly using open-pit extraction methods. One of the most significant environmental threats associated with these operations is the generation of large quantities of suspended particulate matter, which can adversely affect soils, vegetation, and nearby populations (Antoine 2012). Consequently, mining companies operating in the country are confronted not only with production-related challenges but also with increasing concerns regarding the environmental and health impacts of dust emissions, which have become a major issue for both employers and local authorities.

The National Industrial and Mining Company (SNIM), which specializes in iron ore production, operates mainly in the regions of Tiris Zemmour (Zouerate) and Dakhlet Nouadhibou in Mauritania (Fig. 1). Ore extraction and processing activities are primarily conducted near the city of Zouerate, after which the products are transported to Nouadhibou for export. The ore is conveyed over a distance of approximately 700 km by a 2.5 km-long ore train with an average capacity of 17,000 tons per convoy. From Nouadhibou, where SNIM operates two mineral ports, iron ore products are shipped to several Western European countries and China (SNIM 2022).

An Environmental and Social Impact Assessment conducted in 2009 for SNIM mining operations indicated that air pollution associated with the Guelbs 1 facilities

posed significant risks to human health, particularly with respect to respiratory diseases. The assessment further predicted that the development of the Guelbs 2 project would increase dust emissions by 10–15% (Tecsult International Limited 2009). According to this study, workers involved in the Guelbs 2 project would be exposed to concentrations of total dust, quartz dust containing silica, iron dust, and welding fumes exceeding applicable regulatory standards, thereby necessitating the implementation of corrective and preventive measures.

In this context, the present study focuses on dust emissions from SNIM's Guelb El Rhein (Guelbs 2) and Kedia d'Idjil (TO14, Rouessa) mining sites, which were selected due to their substantial dust-generating activities. A corrective approach was implemented alongside a detailed assessment of dust emission levels and the associated bacterial load in airborne particles. The objective of this work is to improve understanding of the combined environmental and health impacts of mining-related dust emissions and to contribute to the development of effective mitigation strategies for mining environments.

2. MATERIALS AND METHODS

Dust assessment can be conducted using three main approaches: (i) measurement of airborne particulate concentrations (TSP, PM₁₀, PM_{2.5}), expressed in $\mu\text{g m}^{-3}$ (Winberry 1999); (ii) quantification of deposited dust expressed in $\text{mg m}^{-2} \text{day}^{-1}$, which is the method adopted in this study; and (iii) estimation of total annual emissions expressed in tons per year (t yr^{-1}), according to particle size fractions.

The quantification method applied in this study is based on dust sampling over a defined exposure period. From a qualitative perspective, several analytical techniques were employed: X-ray fluorescence (XRF) for elemental composition, X-ray diffraction (XRD) for mineralogical characterization and silica content, atomic absorption spectrometry for heavy metal determination, the Most Probable Number (MPN) method for bacterial quantification, and the API gallery system for bacterial species identification (NIOSH 1998).

2.1. Quantification of Sedimented Dust

The procedure used to measure deposited dust followed the French Standard NF X 43-014 (AFNOR 2003), as recommended by INERIS (2004). Dust was collected using Bergerhoff or Owen deposit gauges, which were weighed before and after the exposure period. Sampling devices were installed in accordance with standard recommendations, away from high-risk zones and major traffic sources.




Fig. 1: Location of sampling stations within the National Industrial and Mining Company (SNIM) in the Tiris Zemmour region, Mauritania. Stations A, B, and C are located in the Guelb El Rhein mine along the Guelb 2 production circuit. Station D (Rouessa) and stations E and F (TO14 area) are situated in the Kédia d'Idjil mine. Stations G, H, and I are located in downtown Zouerate.

The dust monitoring was conducted using Owen gauges, which were strategically positioned at each sampling point at a standardized height of 1.5 m above ground level to minimize the influence of saltating particles from the soil surface. To quantify the atmospheric dust fallout, the total mass of the collected particles was related to the sampling surface and the exposure time. The dust deposition rate (T) was calculated according to the following formula:

$$T = \frac{P}{S \times t}$$

Where:

- T : dust deposition rate ($\text{mg m}^{-2} \text{day}^{-1}$)
- P : total mass of dust collected in the gauge (mg)
- S : surface area of the gauge opening (m^2)
- t : duration of the exposure period (days)

This calculation procedure follows the rigorous recommendations of the NF X 43-014 standard for the measurement of atmospheric dust fallout. This standardized approach ensures that the exceptionally high values recorded at the source-point stations are a precise reflection of the intense industrial activity at the mining sites during the sampling window.

Dust pollution levels were classified as follows:

- Very low: $< 150 \text{ mg m}^{-2} \text{day}^{-1}$ (low pollution area)
- Low: $150\text{--}350 \text{ mg m}^{-2} \text{day}^{-1}$ (moderately polluted area)
- High: $350\text{--}500 \text{ mg m}^{-2} \text{day}^{-1}$ (polluted area)
- Very high: $500\text{--}1,000 \text{ mg m}^{-2} \text{day}^{-1}$ (highly polluted area)

Extremely high: $> 1,000 \text{ mg m}^{-2} \text{day}^{-1}$ (very heavily polluted area) (Budillon Rabatel 2009)

2.2. Study Area and Sampling Strategy

The study was conducted at two major mining sites located near Zouerate, northern Mauritania: the Guelb El Rhein magnetite mine and the Kédia d'Idjil hematite mine. Sampling was carried out over two weeks from August 2 to August 16, 2019.

Dust deposition was monitored at nine sampling stations (Fig. 1). Six stations were located within the mining areas: three along the Guelb 2 production circuit and three at Kédia d'Idjil. The remaining three stations (G, H, and I), located in central Zouerate, served as urban control sites.

Climatic conditions during the monitoring period were characterized by temperatures ranging from $22\text{--}43^\circ\text{C}$, no precipitation, and wind speeds ranging from $12\text{--}52 \text{ km h}^{-1}$.

(Blanchon & Penven 2012). While the two-week monitoring period (August 2–16, 2019) captures a specific temporal window, it coincides with extreme summer conditions in the Saharan climate, characterized by peak temperatures and significant wind speeds conducive to maximum dust suspension. We acknowledge that this short duration does not account for annual seasonal variability; however, it provides a critical baseline for peak emission periods during the dry season.

2.3. Chemical and Mineralogical Analyses

Elemental composition of dust samples was determined using X-ray fluorescence (XRF), allowing quantification of major oxides including CaO, SiO₂, MgO, Fe₂O₃, Al₂O₃, SO₃, K₂O, and Na₂O. XRF analysis enabled continuous monitoring of dust composition throughout the sampling period (Jensen et al. 2006).

Mineralogical composition and crystalline structure were analyzed by X-ray diffractometry (XRD) (Ron 2006).

The exploitation of hematite (Fe₂O₃) and magnetite (Fe₃O₄) ores constitutes a significant source of heavy metals such as Ni, Pb, Cu, Co, Cd, and Zn. These metals are transported by dust particles and deposited on surrounding surfaces at distances influenced by wind speed and particle size (Simpson & Spadaro 2016, Embile Jr. et al. 2018). Elevated iron concentrations commonly observed in polluted mining environments may induce analytical interferences, notably Fe/Cu and Co/Fe interactions (Adamson et al. 2000, Manceau et al. 2002, Sammut 2007, Aïllal et al. 2023).

To minimize such interferences, heavy metal analyses were performed on the fine fraction of iron ore dust using Flame Atomic Absorption Spectroscopy (FAAS) following acid digestion with HCl, HF, and HNO₃ (López-García et al. 1995). The metals quantified included Co, Cu, Ni, and Pb (Fig. 5).

2.4. Microbiological Analyses

2.4.1. Total Cultivable Bacterial Quantification

For inoculum preparation, dust samples from each sampling site were homogenized, weighed, and stored in sterile screw-capped tubes. For each dust type (magnetite from Guelb El Rhein and hematite from Kédia d'Idjil), 1.5 g of dust was suspended in 5 mL of sterile physiological saline solution. Serial dilutions were prepared using physiological saline (9 g L⁻¹ NaCl) according to the suspension–dilution technique described by Rapilly (1968).

Bacterial enumeration was performed using the Most Probable Number (MPN) method with a five-tube series.

Tubes were incubated at 37°C for 48 h. Nutrient broth, thioglycollate medium, and LB broth were used for bacterial growth and quantification (Chandrapati & Williams 2014).

2.4.2. Isolation and Characterization of Anaerobic Bacterial Strains

Three grams of hematite dust and three grams of magnetite dust were suspended separately in 5 mL of sterile physiological water and subjected to thermal shock at 63°C to eliminate vegetative cells while preserving sporulated forms (Cammalleri et al. 2015). Serial dilutions were prepared, and selected dilutions were inoculated onto iron sulfite agar under anaerobic conditions and incubated at 37°C.

Dominant strains were purified through successive subcultures. Preliminary identification was based on growth on selective media, Gram staining, spore staining, and macroscopic and microscopic observations using standard taxonomic keys.

2.4.3. Identification Using the API Staph System

Bacterial identification was further performed using the API Staph system, which consists of 19 microtubes containing dehydrated substrates for the differentiation of *Staphylococcus*, *Kocuria*, and *Micrococcus* species (Aouissi et al. 2007, Rouaiguia 2010). The galleries were incubated at 37°C for 24 h, and biochemical profiles were interpreted according to the manufacturer's instructions (Brown & Smith 2015). Final identification was confirmed using dedicated microbial identification software (Debabza 2015).

2.5. Failure Modes, Effects, and Criticality Analysis (FMECA)

The Failure Modes, Effects, and Criticality Analysis (FMECA), also referred to as AMDEC, is a preventive and inductive method used to identify potential failures and implement corrective actions at an early stage. This approach applies to products, processes, and production systems and is widely used in industrial risk assessment and environmental management (Hariz 2009).

2.6. Statistical Analysis

Dust emission data were expressed as mean ± standard deviation (SD). Statistical analyses were performed using STATISTICA version 8 software. Differences between dust emission rates were evaluated using one-way analysis of variance (ANOVA), followed by Tukey's HSD post hoc test when significant differences were observed. Statistical significance was set at $p < 0.05$.

Table 1: Results of dust emission monitoring at the National Industrial and Mining Company (SNIM) during August 2–16, 2019.

Sites	Gauge location	Owen gauge	Emission rate ($\text{mg m}^{-2} \text{ day}^{-1}$)
Guelb El Rhein	Guelb 2	A	38045
		B	27901
		C	50480
Kedia d'Idjil	Rouessa	D	19007
		E	12076
		F	15146
Zouerate	Downtown	G	594
		H	596
		I	592

3. RESULTS AND DISCUSSION

3.1. Dust Emission Rates

Dust emissions occur at different levels of the National Industrial and Mining Company (SNIM). However, this

work focuses only on the iron ore production sites of Guelb El Rhein (Guelb 2), Kedia d'Idjil (TO14 and Rouessa), as well as downtown Zouerate. The results of this monitoring and the measurements carried out during August 2–16, 2019, are presented in Table 1.

To illustrate variation in these emissions, the mean emission rates at each collection site are shown in Fig. 2.

A one-way ANOVA revealed a significant difference in mean dust deposition rates among the three locations, $F(2, 6) = 23.87$, $p = 0.0014$. Tukey's HSD post hoc test showed that the mean dust deposition rate at Guelb El Rhein was significantly higher than those recorded at Kedia d'Idjil ($p = 0.013$) and downtown Zouerate ($p = 0.001$). However, the difference between Kedia d'Idjil and downtown Zouerate was not statistically significant ($p = 0.084$; Fig. 2).

3.2. Chemical Analysis

Dust collected from the two sites (Guelb El Rhein and Kedia d'Idjil) was subjected to XRF, XRD, and AAS analyses.




Fig. 2: Mean $\pm$ standard deviation of dust emission rates. Data indicated by different letters are significantly different (Tukey's HSD test, $p < 0.05$).




Fig. 3: Mean $\pm$ standard deviation of percentage composition determined by X-ray fluorescence (XRF) analysis.




Fig. 4: X-ray diffraction diagrams for dust samples from Guelb El Rhein and Kedia d'Idjil. H, hematite; M, magnetite; Q, quartz.

Fig. 3 shows the results of the X-ray fluorescence (XRF) analysis.

The results showed that dust emitted at Guelb El Rhein contained $38.1 \pm 2.2\%$ iron oxide, indicating magnetite-type ore (37% iron content); however, this ore must undergo an enrichment process to reach the commercial value set by the company at 66% iron. Dust from Kedia d'Idjil contained $65.7 \pm 1.9\%$ iron, consistent with hematite-type ore (66% iron content).

Based on these results, an X-ray diffractometer was used to determine the type of silica in the iron oxide-silica mixture in a fraction approaching the alveolar fraction (Fig. 4).

Mineralogical analysis indicated that the iron ore dust from the National Industrial and Mining Company was composed mainly of hematite (Fe_2O_3), quartz (SiO_2), and magnetite (Fe_3O_4).

The diffractogram of dust emitted from Guelb El Rhein revealed three main intense peaks: a quartz peak at $2\theta = 27^\circ$, a magnetite peak at $2\theta = 36^\circ$, and a hematite peak at $2\theta = 35^\circ$. The diffractogram of dust emitted from Kedia d'Idjil showed mainly hematite at five intense peaks, magnetite at weaker peaks, and quartz with a prominent peak at 2θ

$= 35^\circ$. The high iron levels in the dust, as shown in Fig. 3, may cause significant analytical interference with heavy metals. Dust samples from Kedia d'Idjil and Guelb El Rhein were analyzed using flame atomic absorption spectroscopy (FAAS) (Fig. 5).

The primary focus of the study was to assess dust emissions from iron ore extraction, with a particular emphasis on heavy metal concentrations. To this end, representative samples of iron ore dust from Guelb El Rhein and Kedia d'Idjil were collected and subjected to detailed analysis (Fig. 5). The findings reveal distinct variation in mean levels of heavy metals. The highest peaks belong to cobalt (Co) and the lowest to copper (Cu). Kedia d'Idjil dust samples have average values of 62.42 mg kg^{-1} of Co, 16.41 mg kg^{-1} of Cu, 29.08 mg kg^{-1} of Ni, and 25.71 mg kg^{-1} of Pb, respectively. However, the dust samples from Guelb El Rhein have Co, Cu, Ni, and Pb concentrations of 71.62, 22.34, 42.04, and 42.62 mg kg^{-1} , respectively. The concentration order for the Kedia d'Idjil samples is $\text{Co} > \text{Ni} > \text{Pb} > \text{Cu}$, with values of 62.42 mg kg^{-1} , 29.08 mg kg^{-1} , 25.71 mg kg^{-1} , and 16.41 mg kg^{-1} . For the Guelb El Rhein samples, the order is $\text{Co} > \text{Pb} > \text{Ni} > \text{Cu}$, with corresponding values of 71.62 mg kg^{-1} , 42.62 mg kg^{-1} , 42.04 mg kg^{-1} , and 22.34 mg kg^{-1} .




Fig. 5: Heavy metal (Co, Cu, Ni, and Pb) contents measured in the two iron ore dust types, Guelb El Rhein dust and Kedia d'Idjil dust.

Notably, the concentrations of heavy metals in dust samples collected from the city center of Zouerate were below the detection threshold. The concentration of Co (71.62 mg kg^{-1}) and Ni (42.04 mg kg^{-1}) in Guelb El Rhein dust is notably high when compared to the urban control (Zouerate), where these metals were undetectable. These levels approach or exceed common soil quality guidelines for industrial zones, suggesting that long-term deposition could lead to significant soil enrichment and potential ecological transfer, particularly given the fine fraction's capacity for transport. This study also shows that heavy metal content in the Guelb El Rhein samples of dust is higher than that in samples from Kedia d'Idjil. This conclusion indicates a potentially higher environmental impact of dust emissions in Guelb El Rhein when compared with Kedia d'Idjil. In brief, the present investigation offers comparisons of heavy metals contained in dust emissions as produced by iron ore mining. Results highlight the need for monitoring and controlling these emissions to reduce their environmental and human health implications.

3.3. Bacterial Analysis

3.3.1. Bacterial Quantification

The five-tube most probable number (MPN) technique was used to obtain quantitative estimates of the bacteria in each sample. Quantification was achieved by serial dilutions of a bacterial culture, dividing the sample into replicates, followed by incubation and subsequent visual examination of each sample to determine growth. Table 2 presents the abundance of bacterial groups.

Although direct enumeration by the MPN method yielded no detectable bacteria in the Guelb El Rhein dust sample (Table 2), this result indicates that the concentration of culturable bacteria in the untreated 1.5 g sample was below the detection limit of the method rather than confirming their complete absence. The *Staphylococcus* isolates reported in Table 3 were recovered through selective enrichment and culture on staphylococci-specific media, followed by identification using the API Staph system. They were not obtained through the heat-shock procedure. The heat treatment at 63°C was performed separately to select heat-resistant or spore-forming bacteria and was not used for the isolation of *Staphylococcus* species.

Table 2: Bacterial quantification results.

Bacterial group	Types of samples	MPN g^{-1}
Total anaerobic and microaerophilic bacteria	Guelb El Rhein	0
	Kedia d'Idjil	2100
Total aerobic bacteria	Guelb El Rhein	0
	Kedia d'Idjil	75

3.3.2. Bacterial Strain Isolation and Characterization

After subculturing the bacterial strains isolated from several prepared selective media and incubated at 37°C , the macroscopic study of the colonies obtained made it possible to differentiate several microbial strains according to their appearance (size, color, etc.). All strains were subcultured on nutrient agar and incubated at 37°C . Several subcultures were subsequently carried out to ensure their purity (by careful observation of the colonies formed after re-isolation and differential Gram staining). Fifteen pure Gram-positive bacterial strains were selected for subsequent analysis.

3.3.3. Results of API Staph Identification

The appropriate API gallery for isolate identification was selected based on macroscopic, microscopic, and respiratory enzyme results, which guided identification toward specific bacterial genera.

After 24 h of incubation at 37°C , the API Staph gallery results (Table 3) were read and interpreted according to the manufacturer's technical sheet.

MPN testing confirmed the presence of cultivable bacteria in the dust emitted by SNIM.

Through an examination of the API Staph gallery, species were identified in the dust. Notably, several of the isolated commensal staphylococci may also act as opportunistic pathogens. The dust sample from Kedia d'Idjil, rich in hematite, also has a distinct bacterial community from the magnetite-rich Guelb El Rhein. The percentage composition of the different staphylococcal strains differed between the two samples, indicating differences in the bacterial composition of the two dust types. The predominant microorganism in Kedia d'Idjil dust is *Staphylococcus cohnii* subsp. *urealyticum* (37.51%), followed by *Staphylococcus capitis* (24.38%). The dust from Guelb El Rhein is characterized by a higher percentage of *Staphylococcus lentus* (22.76%) and

Table 3: Relative abundance (%) of bacterial species identified using the API Staph system in the two iron ore dust types.

Species	Kedia d'Idjil dust (hematite) (%)	Guelb El Rhein dust (magnetite) (%)
<i>Staphylococcus hominis</i>	11.92	10.96
<i>Staphylococcus capitis</i>	24.38	20.77
<i>Staphylococcus lentus</i>	0.00	22.76
<i>Staphylococcus epidermidis</i>	0.00	12.10
<i>Staphylococcus auricularis</i>	12.43	5.71
<i>Staphylococcus haemolyticus</i>	13.76	5.71
<i>Staphylococcus cohnii</i> subsp. <i>urealyticum</i>	37.51	0.00
<i>Staphylococcus sciuri</i>	0.00	22.00

Staphylococcus sciuri (22.00%). This variation in dominant bacteria indicates a site-specific diversity. Furthermore, Kedia d'Idjil dust is relatively evenly distributed and rich in *Staphylococcus capitis*, *Staphylococcus auricularis*, and *Staphylococcus haemolyticus*. By contrast, the bacterial community in Guelb El Rhein dust is more diverse, with multiple strains such as *Staphylococcus epidermidis* and *Staphylococcus auricularis*. Another interesting feature is the relatively abundant occurrence of *Staphylococcus cohnii* subsp. *urealyticum* in Kedia d'Idjil dust, but it was absent from Guelb El Rhein dust.

This difference raises questions about site-specific environmental factors, such as geology and climate, that may influence bacterial composition.

Comparing our results with similar studies could deepen our understanding of these findings. Future research might investigate whether these differences match patterns observed in other regions with different geological or climatic traits. Additionally, looking into the potential ecological and health effects of such variations in bacterial composition could help assess the environmental impact of these iron-rich dust types more thoroughly.

Table 4: Causes of dust emissions at SNIM.

Material	Equipment	Method	Middle	Workforce
Production quantity and delivery	Workshop maintenance	Accidental walking	Climatic condition	Lack of discipline Lack of training
Granulometry	Lack of means of recovery	Inefficient air filtration system	Air Currents	Lack of awareness
Humidity	Lack of staff	Competence management	Lack of cleaning	Lack of motivation
	Drop points	Corrective maintenance	Clutter	

Table 5: FMECA (AMDEC) rating grid.

Severity (G)	
Note	Criteria
1	Indirect effect
2	Effect causing emission
3	Origin of the problem
4	Direct effect (dust emitter)
Frequency (F)	
Note	Criteria
1	Once a year
2	1 time/month
3	1 time/week
4	1 time/day
Detectability (D)	
Note	Criteria
1	Detection with appropriate indicator
2	Detection by an indirect indicator
3	Visual
4	Not detectable

3.4. Recommendations to Reduce Dust Emissions at SNIM

3.4.1. Causes of Failure

This study aims to analyze the causes of the significant increase in dust emissions within the SNIM company and to propose appropriate solutions. Based on this analysis, it seeks to identify ways to reintroduce dust fallout into the production circuits.

These dust fallouts mainly correspond to iron production losses occurring at the final stage of processing. Their recovery and reintegration into the production cycle offer both economic benefits and a reduction in waste.

Table 4 presents the potential causes of dust emissions and their classification. These causes are grouped into five categories according to their origin within SNIM facilities.

3.4.2. Criticality Scale

The criticality index is calculated for each failure from the combination of three criteria (Table 5), by multiplying their respective scores:

$$C = G \times F \times D$$

Table 6: Criticality results and analysis.

Causes (aspects)	Analyses	G	F	D	C
Lack of staff, training, awareness, and motivation	A direct effect on production, daily, and not detectable.	4	4	4	64
Accidental walk	With an indirect effect on production, annually, and can be detected visually.	1	1	3	3
Lack of workshop maintenance	A direct effect on production, daily and non-detectable.	4	4	4	64
Lack of cleaning	With a direct effect on daily production, and detectable with an appropriate indicator.	4	4	1	16
Inefficient air filtration system	With a direct effect on daily production, and detectable with an appropriate indicator.	4	4	1	16
Footprint	A direct effect on production, daily and non-detectable.	4	4	4	64
The drop points	With a direct effect on production, annually, and detectable with an appropriate indicator.	4	1	1	4
Absence of means of recovery in time	A direct effect on production, daily, and non-detectable.	4	4	4	64
Presence of draft	With a direct effect on daily production, and detectable with an appropriate indicator.	4	4	1	16

Where:

G: severity of the consequences generated by the failure (severity index);

F: frequency of occurrence of the failure (frequency index); and

D: detectability of the failure (detectability index).

Each of these criteria will be evaluated with a rating table established at four levels. Table 5 shows the criticality rating scale used.

Any criticality value greater than 24 will be classified as a significant aspect whose occurrence will be addressed by the team.

The first level of criticality is based on the severity rating (assessment of the magnitude of the impact and its reversibility), while the second is based on repetition within a specified time interval.

The last level of criticality is based on dust emission indicators. Thus, for each criterion, a score ranging from 1 to 4 will be assigned (the weighting factor for each criterion is 4).

The causes will be classified according to the scale of criticality. Thus, the causes and their criticality analyses will be presented at the beginning. Any cause with a criticality rating greater than 24 will be considered significant.

The values presented in Table 6 will make it possible to measure the causes of dust emission problems based on criticality.

Due to the lack of history of these measurements, the threshold value of criticality was arbitrarily set at 24, which is a value resulting from a very serious cause, with an intermediate frequency and detected by an indirect indicator.

$$C = 4 \times 3 \times 2 = 24$$

To use the results of the FMECA analysis, technical solutions were proposed to reduce the criticality of the most consequential failure modes and improve operational availability.

A maintenance policy based on the improvement actions of the AMDEC analysis is recommended, allowing the implementation of a preventive maintenance plan (Table 7).

3.5. Discussion

Dust deposition rates exhibited marked spatial variability across the sampling sites (Fig. 2), reflecting heterogeneous emission dynamics within and between the mining areas. During the exposure period, the Owens gauges collected 37,500 mg, 27,500 mg, and 49,700 mg of dust at stations a, b, and c, respectively, at the Guelb El Rhein mine, compared to 18,600 mg, 11,800 mg, and 14,800 mg at stations d, e, and f at the Kedia d'Idjil mine. The exceptionally high deposition

Table 7: Significant criticalities and proposed remedies.

Causes (aspects)	Criticality	Remedy
Lack of staff, training, awareness, and motivation	64	Plan, implement, and follow a training plan for employees
Lack of workshop maintenance		Revise the maintenance plan to promote preventive maintenance
Footprint		Expansion or investment of another line
Absence of means of recovery in time		Installation of a mining dome and acquisition of an industrial vacuum cleaner

rates recorded at Guelb El Rhein ($38,045\text{--}50,480\text{ mg m}^{-2}\text{ day}^{-1}$) significantly exceed the threshold for extremely polluted environments ($>1000\text{ mg m}^{-2}\text{ day}^{-1}$). These values reflect the strategic positioning of Owen gauges (Stations a, b, and c) in immediate proximity to the primary crushing and enrichment units of the Guelb 2 production circuit. These measurements represent source-point emissions under peak operational conditions rather than ambient background levels, explaining the several orders of magnitude difference from standard polluted environment thresholds.

These variations indicate that dust emissions are strongly influenced by local working conditions, equipment operation and maintenance, climatic parameters, and human activities.

Dust deposition rates differed markedly among the three study areas. Guelb El Rhein had the highest mean deposition rate ($38,808.7 \pm 11,308.9\text{ mg m}^{-2}\text{ day}^{-1}$), followed by Kédia d'Idjil ($15,409.7 \pm 3,473.0\text{ mg m}^{-2}\text{ day}^{-1}$), whereas the lowest rate was recorded in downtown Zouérate ($594.0 \pm 2.0\text{ mg m}^{-2}\text{ day}^{-1}$). Thus, mean deposition at Guelb El Rhein and Kédia d'Idjil was approximately 65 and 26 times higher, respectively, than at the urban control site. These findings indicate substantially greater dust deposition within the mining areas, particularly at Guelb El Rhein. The NF X 43-014 standard was used as a measurement protocol and not as a source of regulatory or occupational exposure limits.

X-ray fluorescence analysis confirmed the presence of silicon dioxide (silica) in dust samples from both mining sites, with notably higher concentrations at Guelb El Rhein (Fig. 3). The toxicity of mineral dust, particularly crystalline silica, has been extensively documented (Fubini et al. 1999). Inhalation of silica-containing dust represents a major occupational health risk, as respirable particles can penetrate deep into the pulmonary region where gas exchange occurs (Roberge et al. 2004). The International Agency for Research on Cancer has classified inhaled crystalline silica (quartz and cristobalite) as a Group 1 human carcinogen (IARC 1997). However, its carcinogenic potential may vary depending on mineralogical characteristics, surface properties, particle morphology, and associated impurities (IARC 2012, UNICEM 2014).

Beyond chemical hazards, the microbiological analysis revealed a substantial presence of *Staphylococcus* species in the dust, highlighting an additional biological risk associated with mining emissions. Several studies have demonstrated that *Staphylococcus* spp. can exhibit high resistance to heavy metals such as copper, nickel, cobalt, and lead (Nieto et al. 1989, Sabry et al. 1997, Avilés et al. 1993), potentially due to selective pressure exerted by metal-rich environments. This resistance may be further enhanced by the widespread use

of metal-based compounds in industrial and environmental applications (Hiramatsu et al. 1997).

Notably, *Staphylococcus* spp. have also been reported to possess biosorption capacities for heavy metals. Kumar et al. (2010) demonstrated up to 93% lead removal by *Staphylococcus* strains, while Rahman et al. (2019) reported strong lead-binding capacity in *Staphylococcus hominis* strain AMB-2. These findings suggest a dual role of *Staphylococcus* spp. as both indicators of environmental contamination and potential agents of bioremediation.

Marked differences in *Staphylococcus* community composition were observed between the two mining sites, reflecting site-specific environmental conditions. At Kédia d'Idjil, *Staphylococcus capitis* (24.38%) and *Staphylococcus cohnii* subsp. *urealyticum* (37.51%) predominated, whereas dust from Guelb El Rhein was characterized by higher proportions of *Staphylococcus lentus* (22.76%) and *Staphylococcus sciuri* (22.00%). Several of these species are known opportunistic pathogens associated with bacteremia, endocarditis, urinary tract infections, and other severe clinical conditions (Wieser & Busse 2000, Meservey et al. 2020, Fan et al. 2020, Koksai et al. 2009).

The relatively homogeneous distribution of species in Kédia d'Idjil dust contrasts with the higher microbial diversity observed at Guelb El Rhein, suggesting that geological context, mineral composition, and operational practices may exert strong selective pressure on microbial communities. Furthermore, resistance to heavy metals and antibiotics has been reported in species such as *Staphylococcus epidermidis* and *Staphylococcus haemolyticus*, with co-selection mechanisms driven by environmental contamination (Sitthisak et al. 2005, Xue et al. 2015).

We hypothesize that the distinct bacterial communities observed, dominated by *S. capitis* in the hematite-rich Kédia d'Idjil site and *S. lentus* in the magnetite-rich Guelb El Rhein site, reflect selective pressures exerted by the specific mineralogical composition and associated heavy metal profiles of each environment. In metal-enriched mining settings, microorganisms often develop adaptive mechanisms that allow them to tolerate or transform toxic elements, including metal efflux systems and intracellular sequestration processes, as highlighted by Rajput et al. (2025). Although further mechanistic and statistical analyses are required to confirm this relationship, the observed site-specific microbial diversity suggests that the mineralogical characteristics of iron ores may act as a key driver shaping microbial assemblage structure in these mining environments.

Finally, the Failure Modes, Effects, and Criticality Analysis (FMECA) highlighted poor workshop maintenance, operational congestion, and delays in corrective measures as

major contributing factors to dust emissions, with criticality indices exceeding 24. Preventive actions targeting these factors were therefore recommended to improve production efficiency, reduce emissions, and extend equipment lifespan.

4. CONCLUSIONS

Dust emissions along the production circuit of the Guelb El Rhein mine operated by the National Industrial and Mining Company have reached critical levels, posing serious environmental and occupational health risks. This study developed and implemented a comprehensive monitoring framework to quantify dust emissions, identify their sources, and assess both their chemical composition and associated bacterial load.

The results demonstrated that dust emission levels at Guelb El Rhein substantially exceed the stated dust-pollution threshold ($>1,000 \text{ mg m}^{-2} \text{ day}^{-1}$), rendering certain areas unsafe for personnel involved in production monitoring. Emissions at Kedia d'Idjil were lower than those at Guelb El Rhein but still substantial, highlighting the influence of site-specific operational and geological factors.

Chemical analyses confirmed the presence of silica and heavy metals in emitted dust, while microbiological investigations revealed a diverse assemblage of potentially pathogenic *Staphylococcus* species, underscoring the combined chemical and biological hazards associated with mining dust. These findings emphasize the need for integrated mitigation strategies addressing both particulate matter and microbial exposure.

To reduce dust emissions and associated risks, the implementation of preventive measures is strongly recommended, including:

- installation of enclosed mining domes to limit dust dispersion;
- deployment of conveyor belt recovery systems to recapture iron-rich dust;
- integration of air filtration and ventilation systems in enclosed workshops;
- installation of localized extraction hoods at material drop points to direct dust toward filtration units.

In addition to occupational exposure, gravity-driven deposition of particulate matter can lead to contamination of soils, vegetation, and water bodies, potentially causing habitat degradation and biodiversity loss if emissions are widespread and persistent. Overall, this study highlights the urgency of adopting sustainable dust management practices in mining environments to protect workers, surrounding ecosystems, and public health.

REFERENCES

- Adamson, I.Y., Prieditis, H., Hedgecock, C. and Vincent, R., 2000. Zinc is the toxic factor in the lung response to an atmospheric particulate sample. *Toxicology and Applied Pharmacology*, 166(2), pp. 111–119. [DOI]
- AFNOR, 2003. *NF X43-014: Détermination des retombées atmosphériques totales – Échantillonnage, préparation des échantillons avant analyses*. 32 p.
- Aillal, M., Khazri, A., Al-Hoshani, N., Boufahja, F., Beyrem, H. and Lafdal, M.Y., 2023. Are iron ore microparticles toxic for the European clam *Ruditapes decussatus*? Response elements from biomarker activities and in silico modeling. *Saudi Journal of Biological Sciences*, 30(8), p. 103718. [DOI]
- Antoine, C., 2012. *Exposition aux poussières provenant d'une mine à ciel ouvert: évaluation des risques et biodisponibilité des métaux*. Essai présenté au Centre Universitaire de Formation en Environnement. 87 p.
- Aouissi, A., Fouzari, A. and Meziane, N., 2007. *Qualité bactériologique de l'eau d'Oued Seybouse*. Mémoire d'ingénieur, Université 8 Mai 1945 Guelma. 57 p.
- Avilés, M., Codina, J.C., Pérez-García, A., Cazorla, F., Romero, P. and De Vicente, A., 1993. Occurrence of resistance to antibiotics and metals and of plasmids in bacterial strains isolated from marine environments. *Water Science and Technology*, 27, pp. 475–478.
- Blanchon, V. and Penven, M., 2012. *Jean Lefebvre Méditerranée Fos sur Mer (13): Compte rendu des mesures des retombées atmosphériques de poussières dans l'environnement autour de la future plate-forme multimodale*. 19 p.
- Boussahel, S., 2007. *Analyse des retombées particulaires aux alentours de la cimenterie d'Ain-El-Kebira*. Mémoire de fin d'étude en vue de l'obtention du diplôme d'ingénieur d'état en écologie végétale et environnement. 88 p.
- Brown, A.E. and Smith, H., 2015. *Benson's Microbiological Applications: Laboratory Manual in General Microbiology*, 13th ed. McGraw-Hill Higher Education, New York. 5 p.
- Budillon Rabatel, 2009. *Carrière de Saint-Paul-lès-Romans: Règles techniques concernant les poussières*. Document technique, section 9.2.4. 15 p.
- Cammalleri, M., Pipitone, E., Rubino, T., Saporito, L. and Mammina, C., 2015. Experimental evaluation of a new thermal process for microorganisms inactivation. *Journal of Food Process Engineering*, 38(5), pp. 445–451. [DOI]
- Chandrapati, S. and Williams, M.G., 2014. Total viable counts: Most probable number (MPN). In: Batt, C.A. and Tortorello, M.L. (eds.), *Encyclopedia of Food Microbiology*, 2nd ed., pp. 621–624. Academic Press. [DOI]
- Debabza, M., 2015. *Emergence en milieu hospitalier des bacilles Gram négatifs multirésistants aux antibiotiques: étude bactériologique et moléculaire*. Thèse de doctorat d'état, Université Badji Mokhtar-Annaba, pp. 5–65.
- Eduard, W., Heederik, D., Duchaine, C. and Green, B.J., 2012. Bioaerosol exposure assessment in the workplace: the past, present and recent advances. *Journal of Environmental Monitoring*, 14(2), pp. 334–339. [DOI]
- Elodie, G., Diane, L.B. and Fabrice, A., 2020. Rôle des particules dans la contamination en micropolluants des retombées atmosphériques. *PIREN-Seine Phase 8 - Rapport 2020*, p. 9.
- Emble Jr., R.F., Walder, I.F., Schuh, C. and Donatelli, J.L., 2018. Cu, Pb and Fe release from sulfide-containing tailings in seawater: results from laboratory simulation of submarine tailings disposal. *Marine Pollution Bulletin*, 137, pp. 582–592. [DOI]
- Fan, Z., Yang, Y., Li, D. and Fei, Q., 2020. A rare lumbar pyogenic spondylodiscitis caused by *Staphylococcus caprae* with initial misdiagnosis: case report and literature review. *BMC Surgery*, 20(1), p. 200. [DOI]

- Fubini, B., Zanetti, G., Altília, S., Tiozzo, R., Lison, D. and Saffiotti, U., 1999. Relationship between surface properties and cellular responses to silica: studies with heat-treated cristobalite. *Chemical Research in Toxicology*, 12(8), pp. 737–745. [DOI]
- Hariz, S., 2009. *Etude critique du système de management environnemental au niveau des entreprises algériennes*. Mémoire master en Hygiène et Sécurité Industrielle. 161 p.
- Hassanvand, M.S., Naddafi, K., Faridi, S., Nabizadeh, R., Sowlat, M.H., Momeni, F., Gholampour, A., Arhami, M., Kashani, H., Zare, A., Niazi, S., Rastkari, N., Nazmara, S., Ghani, M. and Yunesian, M., 2015. Characterization of PAHs and metals in indoor/outdoor PM₁₀/PM_{2.5}/PM₁ in a retirement home and a school dormitory. *Science of the Total Environment*, 527–528, pp. 100–110. [DOI]
- Heederik, D. and von Mutius, E., 2012. Does diversity of environmental microbial exposure matter for the occurrence of allergy and asthma? *Journal of Allergy and Clinical Immunology*, 130(1), pp. 44–50. [DOI]
- Hiramatsu, K., Hanaki, H., Ino, T., Yabuta, K., Oguri, T. and Tenover, F.C., 1997. Methicillin-resistant *Staphylococcus aureus* clinical strain with reduced vancomycin susceptibility. *Journal of Antimicrobial Chemotherapy*, 40(1), pp. 135–136. [DOI]
- INERIS, 2004. *Rapport annuel 2003*. Institut National de l'Environnement Industriel et des Risques, Verneuil-en-Halatte, France. 52 p.
- International Agency for Research on Cancer (IARC), 1997. *Silica, Some Silicates, Coal Dust and para-Aramid Fibrils*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Volume 68. Lyon: IARC. ISBN 978-92-832-1268-3.
- International Agency for Research on Cancer (IARC), 2012. *Arsenic, Metals, Fibres, and Dusts*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Volume 100C. Lyon: IARC. ISBN 978-92-832-1320-8.
- Jensen, H., Pedersen, J.H., Jørgensen, J.-E., Pedersen, J.S., Joensen, K.D., Iversen, S.B. and Søgaard, E.G., 2006. Determination of size distributions in nanosized powders by TEM, XRD, and SAXS. *Journal of Experimental Nanoscience*, 1(3), pp. 355–373. [DOI]
- Koksal, F., Yasar, H. and Samasti, M., 2009. Antibiotic resistance patterns of coagulase-negative staphylococcus strains isolated from blood cultures of septicemic patients in Turkey. *Microbiological Research*, 164(4), pp. 404–410. [DOI]
- Kumar, A., Bisht, B.S. and Joshi, V.D., 2010. Biosorption of heavy metals by four acclimated microbial species, *Bacillus* spp., *Pseudomonas* spp., *Staphylococcus* spp. and *Aspergillus niger*. *Journal of Biological and Environmental Sciences*, 4(12), pp. 97–108.
- López-García, I., Viñas, P., Campillo, N. and Hernández-Córdoba, M., 1995. Use of submicroliter-volume samples for extending the dynamic range of flow-injection flame atomic absorption spectrometry. *Analytica Chimica Acta*, 308(1–3), pp. 85–95. [DOI]
- Ma, J., Chen, L., Guo, Y., Wu, Q., Yang, M., Wu, M. and Kannan, K., 2014. Phthalate diesters in airborne PM_{2.5} and PM₁₀ in a suburban area of Shanghai: seasonal distribution and risk assessment. *Science of the Total Environment*, 497–498, pp. 467–474. [DOI]
- Manceau, A., Lanson, B. and Drits, V.A., 2002. Structure of heavy metal sorbed birnessite. Part III: results from powder and polarized extended X-ray absorption fine structure spectroscopy. *Geochimica et Cosmochimica Acta*, 66(15), pp. 2639–2663. [DOI]
- Meservey, A., Sullivan, A., Wu, C. and Lantos, P.M., 2020. *Staphylococcus sciuri* peritonitis in a patient on peritoneal dialysis. *Zoonoses and Public Health*, 67(1), pp. 93–95. [DOI]
- National Institute for Occupational Safety and Health (NIOSH), 1998. Childhood agricultural safety and health research: Notice of availability of funds for fiscal year 1998. *Federal Register*, 63(13), pp. 3134–3138.
- Nieto, J.J., Fernández-Castillo, R., Márquez, M.C., Ventosa, A., Quesada, E. and Ruiz-Berruero, F., 1989. Survey of metal tolerance in moderately halophilic eubacteria. *Applied and Environmental Microbiology*, 55(9), pp. 2385–2390. [DOI]
- Rahman, Z., Thomas, L. and Singh, V.P., 2019. Biosorption of heavy metals by a lead (Pb) resistant bacterium, *Staphylococcus hominis* strain AMB-2. *Journal of Basic Microbiology*, 59(5), pp. 477–486. [DOI]
- Rajput, P., Benjiwal, S. and Pandey, R., 2025. A comprehensive review on the role of bioremediation in heavy metal contamination. *Nature Environment & Pollution Technology*, 24(S1), pp. 235–245. [DOI]
- Rapilly, F., 1968. *Les techniques de mycologie en pathologie végétale*. *Annales des Épiphyties*, 19 (hors série). 97 p.
- Roberge, B., Ménard, L., Baril, M., Legris, J. and Deadman, J.-É., 2004. *Manuel d'hygiène du travail: du diagnostic à la maîtrise des facteurs de risque*. Mont-Royal, Québec: Modulo-Griffon. xiv + 738 p. ISBN 2-89443-210-0.
- Ron, J., 2006. X-ray techniques: overview. *Encyclopedia of Analytical Chemistry*, p. 20.
- Rouaiguia, M., 2010. *Qualité microbiologique de l'eau de Oued Messida*. Mémoire de master 2, Université 8 Mai 1945 Guelma. 78 p.
- Sabry, S.A., Ghazlan, H.A. and Abou-Zeid, D.M., 1997. Metal tolerance and antibiotic resistance patterns of a bacterial population isolated from sea water. *Journal of Applied Microbiology*, 82(2), pp. 245–252. [DOI]
- Sammut, M., 2007. *Spéciation du cadmium, du plomb et du zinc dans les poussières d'émissions atmosphériques d'origine sidérurgique: approche de l'impact toxicologique des poussières*. Thèse, Université Paul Cézanne. 254 p.
- Simpson, S.L. and Spadaro, D.A., 2016. Bioavailability and chronic toxicity of metal sulfide minerals to benthic marine invertebrates: implications for deep sea exploration, mining and tailings disposal. *Environmental Science & Technology*, 50(7), pp. 4061–4070. [DOI]
- Sitthisak, S., Howieson, K., Amezola, C. and Jayaswal, R.K., 2005. Characterization of a multicopper oxidase gene from *Staphylococcus aureus*. *Applied and Environmental Microbiology*, 71(9), pp. 5650–5653. [DOI]
- SNIM, 2022. *Rapport annuel: un engagement durable et responsable*. Société Nationale Industrielle et Minière, p. 98.
- Tecslut International Limited, 2009. *SNIM's Guelbs II Project, Mauritania – Environmental and Social Impact Assessment – Summary*. Prepared for Société Nationale Industrielle et Minière de Mauritanie, Nouadhibou: SMIM. 636 p.
- UNICEM, 2014. *Prévention des risques liés à l'exposition aux poussières: guide méthodologique pour les industries extractives*. Union nationale des industries de carrières et matériaux de construction, Paris, France.
- Viegas, C., Fleming, G.T.A., Kadir, A., Almeida, B., Caetano, L.A., Gomes, A.Q., Twarużek, M., Kosicki, R., Viegas, S. and Coggins, A.M., 2020. Occupational exposures to organic dust in Irish bakeries and a pizzeria restaurant. *Microorganisms*, 8(1), p. 118. [DOI]
- Wieser, M. and Busse, H.J., 2000. Rapid identification of *Staphylococcus epidermidis*. *International Journal of Systematic and Evolutionary Microbiology*, 50(3), pp. 1087–1093. [DOI]
- Winberry, W., 1999. *Method IO-2.1 - Sampling of ambient air for total suspended particulate matter (SPM) and PM₁₀ using high volume (HV) sampler (EPA/625/R-96/010a)*. Office of Research and Development, U.S. Environmental Protection Agency, Cincinnati. 74 p.
- Xue, H., Wu, Z., Li, L., Li, F., Wang, Y. and Zhao, Z., 2015. Coexistence of heavy metal and antibiotic resistance within a novel composite staphylococcal cassette chromosome in a *Staphylococcus haemolyticus* isolate from bovine mastitis milk. *Antimicrobial Agents and Chemotherapy*, 59(9), pp. 5788–5792. [DOI]
- Yan, C., Leng, Y.L. and Wu, J.T., 2021. Quantitative microbial risk assessment for occupational health of temporary entrants and staff equipped with various-grade PPE and exposed to microbial bioaerosols in two WWTPs. *International Archives of Occupational and Environmental Health*, 94(6), pp. 1327–1343. [DOI]