



Environmental Impact Assessment of Air Quality Issues Caused by the Granite Quarrying and Stone Processing Industry in Ramanagara District, Karnataka State, India

V. Nagaraja[†], C. J. Rakesh, H. N. Sindhu and N. Harishnaika

Department of PG Studies and Research in Applied Geology, Kuvempu University, Jnanasahyadri, Shankaraghatta-577451, Shivamogga District, Karnataka State, India

[†]Corresponding author: V. Nagaraja; nagarajavr02@gmail.com

Nat. Env. & Poll. Tech.
Website: www.neptjournal.com

Received: 30-04-2024

Revised: 06-06-2024

Accepted: 19-06-2024

Key Words:

Environmental Impact Assessment
Granite quarrying
Stone processing
Air quality
Pollutants
Health

ABSTRACT

The environmental impacts of mining, quarrying, and the stone processing industry are significant, affecting air quality, health, and the socioeconomic status of communities worldwide. Key contributors to air pollution include the waste of raw materials from quarrying, non-compliance with scientific protocols, and the extraction of natural mineral resources. The rapid increase in pollution sources, such as dust, water, and noise, has led to the release of various pollutants into the atmosphere, degrading local air quality. This study conducted sampling at twelve sites, adhering to the Central Pollution Control Board's (CPCB) monitoring guidelines. Twelve metrics, including PM₁₀, PM_{2.5}, SO₂, NO_x, CO, O₃, Pb, NH₃, C₆H₆, C₂OH₁₂, As, and Ni, were measured twice a week over a three-month period (January 2024 to March 2024) by the National Ambient Air Quality Standards (NAAQS) in the research area. The results indicated that while SO₂ and NO_x levels were within permissible limits at all monitored locations, Suspended Particulate Matter (SPM) levels were high at every station. The average baseline levels of PM₁₀ (37.17 µg/m³ to 70.52 µg/m³), PM_{2.5} (16.98 µg/m³ to 39.85 µg/m³), SO₂ (5.29 µg/m³ to 13.91 µg/m³), NO_x (9.8 µg/m³ to 29.71 µg/m³), CO (0.15 mg/m³ to 0.32 mg/m³), O₃ (6.9 µg/m³ to 15.37 µg/m³), and NH₃, Pb, Ni, As, C₂OH₁₂, and C₆H₆ were below the detection levels (BDL) and limits of quantification (LOQ), all within the National Ambient Air Quality Standards for commercial, industrial, and residential areas during the study period. This research highlights the urgent need for effective pollution control measures to mitigate the adverse environmental and health impacts of these industries.

INTRODUCTION

Quarrying is the process of extracting minerals from the upper crust of the earth. The natural resources of the world, including land, soil, air, dust, noise, water, the biosphere, human health, society, and the socioeconomic status of the populace, are severely being harmed by the quarrying and stone processing industries (Gbeve 2013, Emmanuel 2018, Umar et al. 2023). This is because there are no scientific techniques for managing granite reserves and resources, and they are overexploited by humans (Vallack et al. 1998, Ukpogong 2012, Sreekala et al. 2023). In addition to air, noise, dust, and water pollution, land degradation, biodiversity loss, topographical relief, flooding, extremely cold temperatures, forest fires, changes in the climate and seasonal pattern variation, greenhouse effects, and health risks from drilling, blasting, and ground vibration, we are currently dealing with serious environmental issues (Singh et al. 2010, Jahed Armaghani et al. 2015, Kittipongvises 2017).

For four or five centuries, natural resources have been overexploited without regard for the consequences to the ecosystem, which has led to these problems (Sheikh et al. 2011, Melodi 2017). The most damaging human activities are those that harm the ecosystem both immediately and over time (Mwangi 2014, Ming'ate et al. 2016). The environment is seriously threatened by overexploitation and negligent practices in the mining and stone processing industries (Odewumi et al. 2015, Okafor et al. 2023).

The primary environmental effects of granite quarries were the creation of artificial ponds, isolated artificial reliefs, regular-shaped depressions, erosion and instability of quarry scarps, partial or total destruction of fluvial terraces, piezometric surface depression, altered groundwater flow direction, formation of periodically flooded areas, permanent removal of areas from agricultural use, changes to farming practices, and pedological changes in soil (Oyinloye et al. 2015, Kofi-Boye 2017, Leon-Kabamba et al. 2020,

Rathore 2020, Roja 2022). To ensure that these regions can eventually be exploited again, people should work toward degraded environment regeneration at every stage of quarrying activities (Saha et al. 2011, Santizo 2022).

The surrounding landscapes of quarrying places are severely impacted by the difficulties in disposing of waste rock (Singh et al. 2010, Sayara 2016). A significant global concern, aside from the granite industry, is human-caused air pollution (Singhal 2018, Salem 2021, Sreekala et al. 2023). Understanding the ways that metropolitan areas impact the chemistry, composition, and life cycles of the atmosphere in several hundred-kilometer downwind regimes can be achieved by considering them as concentrated sources of massive anthropogenic emissions of pollutants (Yeh et al. 1980, Ajibade et al. 2022). Additionally, research on the global epidemiological impacts of air pollution has demonstrated that particulate matter and gaseous pollutants can seriously impair human health, leading to respiratory disorders, heart disease, and cardiopulmonary death (Abdul-Wahab et al. 2015 & 2022, Zimwara et al. 2021).

Fossil fuels and their derivatives are being used more frequently as a result of emerging countries' increasing industry, urbanization, technological innovation, and modernization (Ajah et al. 2018, Zimwara et al. 2021). Thus, the primary challenge facing emerging nations is reducing air pollution, particularly in quickly expanding megacities (Daspan et al. 2018). Estimates from the World Health Organization indicate that air pollution causes nearly two million premature deaths annually, including lung infections, heart attacks, respiratory diseases, and even cancer (Haseeb 2018, Werner et al. 2019). The use of fossil fuels with poor environmental performance, improper land use patterns, industrialization, fast population growth, an increase in vehicles, inadequate transportation infrastructure, improper land use patterns, and-most importantly-ineffective environmental regulations are some of the factors contributing to India's rising air pollution levels (Chattopadhyay et al. 2010, Henry et al. 2017).

The World Health Organization's (WHO) recommended health thresholds for air pollution are exceeded in the majority of Indian cities with populations greater than two million people (Halwenge 2015, Martins et al. 2017). With the aid of remote sensing and GIS software, monitoring the concentrations of various ambient air quality indicators not only prevents detrimental health effects but also provides policymakers with a foundation for successful environmental regulations (Moeletsi 2018, Werner et al. 2019). SPM (suspended particulate matter), SO₂ (sulfur dioxide), and NO_x (nitrogen oxide), together with other chemicals generated by numerous businesses, particularly the granite

industry, are the main pollutants causing environmental destruction in the area. Reduced plant photosynthesis, noise and dust pollution, structural flaws, biodiversity loss, and nuisance dust are only a few of the problems that quarrying is frequently linked to (Nandan et al. 2017, Omeiza et al. 2022).

People's property and health are being significantly impacted by industrial site noise, ground vibrations, drainage, and particulate matter (Sunyer 2001, Stieb et al. 2005). Potential effects of these include decreased agricultural productivity, significant harm to the region's road network, and environmental issues such as soil erosion, vegetation loss, building fissures, biodiversity loss, and topographical changes (Sun et al. 2016, Sayara et al. 2016). Because it generates a variety of contaminants, the granite quarrying and processing industry is dangerous (Tsiouri et al. 2015).

Today, those who live near quarries have extremely terrible lives as a result of pollution. Health problems can affect anyone at any moment; respiratory and skin conditions are two examples (Zanobetti et al. 2009, Yakovleva 2017). The term "AQI" (air quality index) is used because it is challenging to provide high-quality data regarding the concentration of several environmental indicators (Wright 2009, Trivedi et al. 2009 & 2010). The common spaces and woodlands of the communities are covered in waste material, often known as by-products (Yasobant et al. 2017, Abeya 2023).

A significant amount of inert dust is discharged into the sky during the quarrying process, and depending on the wind direction, this dust can travel up to 4 or even 5 kilometers (Chowdhury 2004, Aarthy et al. 2018). This causes cloudbursts, which in turn cause downpours and flash floods by preventing the formation of rain clouds (Ekpa et al. 2022 & 2023). Because it entails the direct exploitation of resources, quarrying is distinct. It supplies the building sector with raw materials. The construction sector needs a lot of stone for its projects. The increasing need for building materials, including aggregate, sand, gravel, and rocks, as well as industrial manufacturing, is driving the market for quarry commodities. This leads to the mismanagement of natural resources, which is detrimental to the ecology and economy.

The objective of the current study is to assess how the granite quarrying, stone-cutting, crushing, and stone-processing activities that are conducted in the study region affect the local environment. The state of Karnataka is home to most of the rich mineral and ore resources, such as bauxite, iron, manganese, gold, copper, and some significant rocks like gneiss, granite, granodiorites, and doleritic dykes.

The Ramanagara region of Karnataka State's granite quarries are the subject of this study's environmental effects. To develop mitigation and control techniques, the research




Fig. 1: Study region with data on elevation.

will gather and analyze air samples, calculate their impact on air quality in residential, industrial, and quarry contexts, and use the results. This research will help future researchers and add to the body of information about the quarrying business. In this district, the local inhabitants and the environment are being negatively impacted by the stone processing and quarrying businesses.

STUDY AREA

On August 23, 2007, the former Bengaluru Rural District—which included the taluks of Ramanagara, Channapatna, Kanakapura, and Magadi—was divided into the Ramanagara district. According to Bhat et al. 1991 & 1994, and Swamy 1998, its largest lengths are 102.25 kilometers in the north-south direction and 62.08 kilometers in the east-west direction. The district is composed of 823 settlements spread over a 3516 km² physical area, of which 699.46 km²—or around 17.21% of the district's total area—are forest. The district's headquarters are in the town of Ramanagara, which is situated roughly 50 miles southwest of Bengaluru, the state capital.

The district of Ramanagara is located between latitudes 12.72°N and 77.27°E. Fig. 1 shows the contour lines that delineate the district's most notable high and low elevation zones, which are located at elevations of 1196 and 289 meters

above mean sea level, respectively. There are 742.50 meters above sea level on average. The district borders Mandya on the west, Bangalore Urban on the northeast, and Tumakuru, Bangalore Rural, and Bangalore Urban on the north.

MATERIALS AND METHODS

Materials

The toposheet was used in this study to derive the morphological area. Toposheet numbers are compiled by the Indian government's Survey of India (SOI). D43R4_57G4, D43R8_57G8, D43X1_57H1, D43X2_57H2, D43X3_57H3, D43X5_57H5, D43X6_57H6, D43X7_57H7, D43X8_57H8, D43X9_57H9, D43X10_57H10, and D43X11_57H11 are some instances of toposheet numbers.

Elevation and land cover data are obtained from DEM Cartosat-1 and LISS satellite photos on the ISRO Bhuvan website. The meteorological and air quality data sets come from a field investigation, a few official technical publications, and the Open Government Data (OGD) platform in India. Several government bodies, such as the Department of Mines and Geology (DMG) and the Ramanagara district mineral survey report, have provided reports and information about quarrying. Table 1 indicates these data sets.

Table 1: Datasets utilized in the field investigation

Data name	Resolution	website
DEM	10m	Indian Geo platform (Bhuvan)
LISS 3	30m	Indian Geo platform (Bhuvan)
Landsat 8	30m	USGS
Toposheets	1:50000	SOI(survey of India)
Rainfall and temperature	0.5*0.5 km	IMD(India meteorological department)
Quarry and rocks	-	DMG(department of Mines and Geology)
Air quality	-	open governmental data (OGD) platform, India

Methodology

Compiling information from many sources: Primary and secondary sources served as the foundation for this investigation. Field and market surveys, maps, questionnaires, interviews, and laboratory tests are examples of primary sources. Books, journals, published and unpublished reports, and official government documents are examples of secondary sources. The effects of air pollution on the local ecology, people, flora and wildlife, crops, and surroundings were evaluated using field surveys and general observations. The Ramanagara district of the Indian state of Karnataka, which is renowned for its rich granite deposits, diversity, and amount of forest cover, is the only area covered by the study. Using a systematic and random selection process, a sample of the homes at the randomly selected quarry sites was created, resulting in the sample households.

The sample consisted of only one or two hundred adjacent towns or residences that are both directly and indirectly affected by the quarrying activity. Field research and a standardized questionnaire were used to collect data from the families. A research study report was prepared for the analysis in the Ramanagara district locations using baseline data collected at many sites throughout a single season (January 2024 to March 2024). The data are shown in Table 2. These figures were used to assess the possible harm

Table 2: Parameter derived from multiple sources

Sl. No.	Attribute	Parameter	Source of Data
1.	Climatology and Meteorology	Wind speed, Wind direction, Dry bulb temperature, Wet bulb temperature, Relative humidity, Rainfall, Solar radiation, Cloud cover and Environmental Lapse.	Field research, collecting data from a variety of locations, including industrial, residential, and quarry sites.
2.	Geology, rock and Soil	Geological history	Field research, Primary and secondary sources, satellite photos, topographical maps, etc.
3.	Ambient Air Quality	PM ₁₀ , PM _{2.5} , SO ₂ , NO _x , Pb, CO, O ₃ , C ₆ H ₆ , As, Ni, NH ₃ , C ₂ OH ₁₂ , Free Silica etc.	Monitored Data (12 locations)

caused by the stone-processing and quarrying industries. They provided guidance on the best course of action for the quarry site and industry, as well as resources for management, government organizations, and mitigation strategies. For this research project, the findings of the analysis report and study summary were carefully considered.

Choosing study sites for the sampling process: Table 3 shows the location, with sampling locations labeled in different environmental settings. Climate data, topography and terrain, human settlements, health status, accessibility to the monitoring site, resource availability, representativeness of the region for establishing baseline status, and representativeness about likely impact areas were taken into account when choosing the locations for ambient air quality monitoring.

Guidelines for collecting ambient air quality samples: For three months, ambient air quality was monitored twice a week in each region. The following parameters were used to create baseline data for the air environment: lead (Pb), arsenic (As), nickel (Ni), free silica, benzene (C₆H₆), Benzo (a) pyrene C₂OH₁₂, ammonia (NH₃), ozone (O₃), nitrogen dioxide (NO_x), sulfur dioxide (SO₂), carbon monoxide (CO); particulate matter (PM_{2.5}), and (PM₁₀) size less than 10 µm. A comparison has been conducted between the national ambient air quality guidelines and the measured concentrations of pollutant parameters.

Equipment for gathering samples: Particulate matter (PM₁₀ and PM_{2.5}) was detected using Envirotech 271; combo PM₁₀ and PM_{2.5} samplers; breathable dust samplers (Envirotech APM 460 BL); and fine particulate samplers (APM 550). Gaseous pollutants, such as SO₂, NO_x, O₃, and NH₃, were sampled using Ecotech's AAS 109, Envirotech's APM 411, and Ecotech's APM 460 gaseous pollution samplers. Electrochemical sensor technology was used for carbon monoxide monitoring. Every year, the monitoring instruments need repairs and calibrations.

Techniques for analysis and sampling: Table 4 enumerates the sampling and analytical methods utilized to track ambient

Table 3: Location and the gathering of samples in various environments

Location Code	Sample code	Locations	Latitude	Longitude	Distance (km) from Ramanagara town	Azimuth Directions	Environmental setting
QL 1	AAQ1	Doddamudawadi quarry site	12°37'38.08" N	77°22'47.27" E	15.74	SE	Commercial
QL 2	AAQ2	Yadamaranahalli quarry site	12°24'54.09"N	77°22'56.53"E	35.54	S	Commercial
QL 3	AAQ3	Hanakadaburu, kodihalli hobali quarry site	12°24'17.24"N	77°32'50.28"E	45.4	SE	Commercial
QL 4	AAQ4	Hanakadaburu, kodihalli hobali quarry site	12°23'49.59"N	77°32'39.51"E	46.58	SE	Commercial
QL 5	AAQ5	Achalu quarry site	12°27'55.43"N	77°21'18.06"E	30.37	S	Commercial
QLV 6	AAQ6	Hanakadaburu village	12°24'1.12"N	77°32'51.18"E	46.5	SE	Residential
QLV 7	AAQ7	Kodahalli village	12°25'9.41"N	77°18'34.70"E	33.66	SW	Residential
QLV 8	AAQ8	Achalu village	12°28'30.11"N	77°21'47.08" E	28.55	S	Residential
QLV 9	AAQ9	Harohalli village	12°40'53.59"N	77°28'26.33"E	21.3	SE	Residential
QLV 10	AAQ10	Bidadi village	12°47'56.48"N	77°23'9.77"E	15.67	NE	Residential
QLIN 11	AAQ11	Harohalli Industrial area	12°41'0.05"N	77°26'43.60"E	18.27	SE	Industrial
QLIN 12	AAQ12	Bidadi industrial area	12°47'54.84"N	77°23'0.33"E	14.51	NE	Industrial

Table 4: Techniques for data analysis for ambient air quality parameter Assessment (NAAQS).

Sl. No.	Parameters	Analytical method	NAAQ standards: 2009	
1.	Sulphur Dioxide (SO ₂), µg/m ³	IS: 11255 (Part 2)/USEPA Method 6	50 (Annual)	80 (24 Hours)
2.	Nitrogen Dioxide (NO ₂), µg/m ³	IS: 5182 (Part - guidelines Volume1 6): 2006 /CPCB	40 (Annual)	80 (24 Hours)
3.	Particulate Matter (PM _{2.5}), µg/m ³	In house method (Gravimetric method) based on CPCB guidelines Volume1	40(Annual)	60 (24hours)
4.	Particulate Matter (PM ₁₀), µg/m ³	IS:5182 (Part-guidelines Volume1 23): 2006/CPCB	60 (Annual)	100 (24 hours)
5.	CO, mg/m ³	IS:5182(Part-10):1999(Reaff:2006) CPCB guidelines Volume1	2 (8 hours)	4 (1hour)
6.	Pb, µg/m ³	IS:5182(Part-22):2004(Reaff:2006) CPCB guidelines Volume1	0.5 (Annual)	1 (24 hours)
7.	O ₃ , µg/m ³	In house method (Spectrophotometric method) based on CPCB guidelines Volume1	100 (8hours)	180 (1hour)
8.	NH ₃ , µg/m ³	In house method (Spectrophotometric method) based on CPCB guidelines Volume1	100 (Annual)	400 (24 hours)
9.	Benzene, µg/m ³	GC FID/ GC MS based on IS 5182 (Part:12) / CPCB guidelines Volume1	5 (Annual)	5 (Annual)
10.	Benzo (a) pyrene, ng/m ³	In House Validated Method By HPCL, UV and GC MS Based on IS:5182(Part- 12) CPCB guidelines Volume1	1 (Annual)	1 (Annual)
11.	Arsenic, ng/m ³	In house method (AAS method) Based on CPCB guidelines Volume 1	6 (Annual)	6 (Annual)
12.	Nickel, ng/m ³	In house method (AAS method) Based on CPCB guidelines Volume 1	20 (Annual)	20 (Annual)

air quality. The closest power sources, which included homes at each AAQ station or governmental entities including Panchayat offices, schools, and temples, were used to power the AAQ equipment.

Table 5 displays the frequency and techniques of monitoring. The analysis report and summary of the air quality survey results were carefully considered for this study project.

Table 5: The frequency and monitoring techniques.

Attributes	Sampling		Measurement Method	Remarks
	Network	Frequency		
Meteorology				
Wind speed, Wind direction, Dry bulb temperature, Wet bulb temperature, Relative humidity, Rainfall, Solar radiation, Cloud cover and Environmental Laps	Selected locations	Continuous for 3 Months	Weather monitors with the database	As per Meteorological department standard. Primary or secondary data
Air Environment				
Particulate Matter (PM ₁₀)	Selected locations	24 hourly-Twice a week for 3 months in non-Monsoon season	Gravimetric (High-Volume with Cyclone)	As per CPCB standards under 18 th November 2009 Notification for National Ambient Air Quality Standards (NAAQS).
Particulate Matter (PM _{2.5})			Gravimetric (High-Volume with Cyclone)	Field survey, Primary or secondary data
Oxides of Sulphur (SO ₂)			EPA Modified West and Gaeke method	
Oxides of Nitrogen (NO _x)			Arsenite Modified Jacob and Hochheiser	
Carbon Monoxide (CO)			Gas Analyzer (NDIR)	
Ozone (O ₃)			UV photometric	
Ammonia (NH ₃)			Indophenol Blue Method	
Lead (Pb)			Atmospheric Absorption Spectrometer	
Arsenic (As)				
Nickel (Ni)				
Benzene			GC-MS/MS	
Benzo Alpha Pyrene			GC-MS/MS	

RESULTS AND DISCUSSION

The Geological Setting of the Granite Quarry in the Study Area

The Ramanagara district's geology: According to Bhat et al. (1994), Syed Abrar et al. (2000 & 2005), Swamy (1998), the rocks in the district come from the Sargur group, the Peninsular Gneissic Complex (PGC), the Charnockite group, the Closepet granite, and the basic and more recent intrusive. The Charnockite group's representative is Charnockite. Lenses found in gneisses, migmatite, amphibolite, and banded magnetite quartzite, which appears as tiny bands—are all part of the Sargur group. Granites, gneisses, and migmatite make up the PGC, which is situated east and west of the Closepet granite (Bhat et al. 1991 & 1994, Simha et al. 2015, Venkatesha et al. 2015). Within the gneisses, the intrusive bodies containing the Closepet granite stretch 50 km in length and 15-20 km in width, with a near-N-S trend. There are reported to be enclaves of migmatite, gneisses, quartzite, and amphibolites among the various compositions found in the Closepet granite. The common intrusives are dolerite, gabbro, pyroxenite, and rarely norite. Dolerite is the most common of the fundamental dikes (Paranthaman

et al. 1995, Swamy 1998, Syed Abrar et al. 2000 & 2005). Three main lineaments in the district go to NNW-SSE. These lineaments range in length from 45 to 70 kilometers. According to the interpretation of this data, a deep-seated fault that is trending NNW-SSE crosses the Closepet granites (Suresh Babu 1995).

The main land uses in the areas covered by quarry leases are plan land, hills, vegetation, various rock formations, and soil cover or overburden. Pebbles are most common in low-lying, loose soils, and most of the leased areas have thin patches of dirt covering them that are between 0.5 and 10 meters thick. Both on higher ground and in deposits at the surface, there are rock outcrops. Reddish-brown soil capping covers the majority of the hill's northern, western, and eastern sections at the lower elevation of the quarry lease area.

The field investigation's findings indicate that the geological setting of the granite quarry area is influenced by several variables, such as the quarry's topography, size, potential deposit formation type, material quality, mineral structure and texture, color, and chemical makeup, as well as the availability of resources, the company's financial standing, the intended output, market, price, supply and demand, and so forth.

Table 6: Lithostratigraphy and rock types in Ramanagara District

Soil Clover	
I. Younger intrusive (More recent invasion)	Extremely potassic rocks; dolerite/gabber/diorite/norite dykes; felsite/felsite porphyry dykes; coarse-grained pink/grey porphyritic granite.
II. Closepet granite (2400–2100 Million years)	Pink/grey porphyritic granite with coarse grain; pink granites (less mafic); grey granites; pink hornblende granite; pink equigranular/porphyrite migmatites.
III. Peninsular Gneissic Complex (3000 Million years)	Grey migmatites, biotite gneiss, leucogneiss and homophonous gneiss, charnockite and migmatite.
IV. Sargur group (> 3000 Million years)	Garnet-sillimanite gneiss, quartzite (fuchsite and BMQ), meta-ultramafites, amphibolites (both massive and schistose), and banded magnetite quartzite.
Not visible is the base	

With an average thickness of fewer than 0.5 meters, the dirt in the middle of the ridge is only present in the spaces between the stones. The established stratigraphic succession of the research region is displayed here, based on fieldwork and observations. The rock types and lithostratigraphy of the Ramanagara District are depicted in Table 6.

The study area is situated in Seismic Zone III, which is characterized as having a very high potential for earthquake shocks of 5 or 6 on the Richter scale, and Seismic Zone IV, commonly known as the high damage risk zone.

In this district, sand mining and defined and non-specified minor mineral leases in Kanakapura, Ramanagara, Channapatna, and Magadi Taluk were conducted in parallel with taluk-wise quarrying operations for dimension stone, construction stone, brick earth, crushers, and stone cutting facilities. Moreover, the area has more crushing factories




Fig. 2: Lithology and its percentage of Ramanagara district.

per square mile than other districts, and four to five decades ago; it was home to several granite quarries and crushing businesses.

The environment is contaminated by the stone-cutting and quarrying processes' noise, dust, water, and other factors. The associated radioactivity varies depending on the surrounding environment due to dust emissions from quarrying and crushing processes. Fig. 2 displays the lithology and area of the Ramanagara district.

Within the research region of the Ramanagara district, which spans a total geographical area of 3516.21 sq.

km, various types of rocks are dispersed as follows: The region encompassed 173.6 sq. km of acid to intermediate charnockite; 3.12 sq. km of amphibolite; 4.24 sq. km of banded iron formation; 1.2 sq. km of banded magnetite quartzite; 0.3 sq. km of cordierite-sillimanite gneiss; 0.96 sq. km of diorite porphyry; 12.4 sq. km of dolerite; 3.26 sq. km of felsite; 12.82 sq. km of gabbro; 0.54 sq. km of granite; 29.36 sq. km of granitoid; 3.23 sq. km of granodiorite; 1522.9 sq. km of grey Granite; 545.83 sq. km of grey hornblende biotite gneiss; 0.69 sq. km of hornblende biotite granite gneiss; 0.59 sq. km of marble; 1080.64 sq. km of migmatite




Fig. 3: Conditions of the soil and its percentage in the study area.

gneiss; 3.37 sq. km of pegmatoidal granite; 12.03 sq. km of pink biotite granite; 8.07 sq. km of pink granite; 6.92 sq. km of pink migmatite; 66.46 sq. km of pink porphyritic granite; 4.66 sq. km of porphyritic granite; 2 sq. km of pyroxene granulite; 0.23 sq. km of quartz vein/reef; 16.19 sq. km of quartzite; and 0.57 sq. km of syenite. The Ramanagara district is entirely covered in rocks. Gray granite and migmatite gneiss are the two main lithologies in this area. The region is roughly 1552 km² and 1080 km².

The Ramanagara District’s Soil Type and Proportion Covered

Although dirt is essential to air pollution, it usually serves no purpose in the quarrying process or other associated operations. About 60% of the Ramanagara district is covered with red, sandy soil (Hema et al. 2012). The remaining soil is loamy and reddish-brown. Geology and soil are closely related in a pedogenic system because of the combined effects of parent material, climate, biosphere, terrain, relief, stage, age, maturity, and time of rock weathering conversion, rainfall, solar radiation, and management (enhancement and

degradation/erosion). The main regions with red sandy soil include the taluks of Channapatna, Kanakapura, Magadi, and Ramanagara due to their varied topographies (Caruthers 1984, Hema et al. 2013, Ganesha et al. 2017). Fig. 3 list every type of soil.

With a total land area of 3516.21 sq. km, the Ramanagara district contains a variety of soil types that are scattered throughout the research region, including the following: 8.59.7 sq.km of skeletal clayey; 17.09 sq.km of dyke ridges; 1634.47 sq.km of fine; 54.79 sq.km of fine loamy; 81.91 sq.km of habitation mask; 12.23 sq.km of loamy; 192.06 sq.km of skeletal loamy; 98.34 sq.km of rock outcrops; 445.7 sq.km of sandy skeletal; 117.54 sq.km of water body mask of habitation.

The Methods Used in the Ramanagara District for Granite Quarrying Operations

In the Ramanagara district region, the majority of stone quarrying activities employ opencast extraction techniques. In open-cast quarrying, significant overburdens such as organic plant cover and topsoil must be removed to reach the

Table 7: Quarrying activities in the Ramanagara district areas leased by the government as of 2022-23.

Sl. No	Taluk	Building stone	Ornamental or dimension stone	Brick earth	Crusher Units	M-sand units
1	Ramanagara	38	02	-	29	08
2	Kanakapura	25	52	-	03	02
3	Magadi	19	04	02	10	02
4	Channapatna	-	02	-	-	-
	Total	82	60	02	42	12




Fig. 4: Quarrying areas spreading in the Ramanagara district.

mineral reserves. The following procedures are involved in extracting granite through open-cast quarrying for dimension or ornamental stone: clearing the area, loading waste or well-dressed materials, transportation, processing raw materials, quality control, shifting, management of stockyards, and recycling waste materials as khandas or brick materials (Rajan Babu et al. 2000 & 2002 & 2003, Drew et al. 2002, Gbeve 2013, Eshiwani 2014,). Stone quarrying, stone crusher buildings, and sand mining are subject to additional operational limitations. There is evidence that these actions have a detrimental effect on the ecosystem.

There are various phases of quarrying operations, and each one has a unique impact on the environment. These phases typically entail deposit exploration and prospecting, the development and preparation of quarries, the use of materials from the quarries, and the processing of the minerals collected at the various installations to create products that may be marketed (Chalekode 1978, Kuzu et al. 2005, Ibrahim et al. 2018, Chaanda et al. 2019). However, there are numerous other environmental risks associated with open-cast quarrying practices. As of 2022–23, Table 7 indicates that the following Ramanagara district lands have been leased by the government (department of mines and geology) for quarrying purposes. Fig. 4 shows the quarrying areas spreading in the Ramanagara district.

Quarry Rock has the third and fourth positions in the world's non-fuel mineral commodities value and volume rankings. However, the quarry's operations deteriorate the land's condition and raise air pollution levels, which negatively impacts the neighborhood's residents and workers (Bada et al. 2013, Bassey et al. 2021).

According to Amankwah et al. (2019) and Bakamwesiga et al. (2021), and the outcome shows that the dust produced during the operation is carried by the atmosphere and travels significant distances from the crusher and quarrying sites to the nearby communities. Table 8 shows that the list of equipment needed to operate a quarry varies according to production capacity.

The amount of air pollution depends on the local microclimate, the size, composition, and volume of dust in the surrounding air, and one of the main causes of air pollution is noise from quarry sites. For example, the mining and processing of granite produce silica dust (Loveson et al. 1997, Esguerra 2008, Doley et al. 2010, Eshiwani 2014, Amankwah et al. 2019).

Dust not only tends to accumulate on surfaces, making it unsightly, but it can also be harmful, especially to people who have respiratory problems. Dust can physically injure plants nearby by obstructing their internal organs and abrading their

Table 8: The minimum machinery and equipment required for quarrying operation (minimum production of 100 cubic meter) and impact on natural environment

Sl. No.	Type	Nos.	Size/ capacity	Make	Motive Power	Impact on natural environment
1.	Compressor	2	440 cfm	Atlas Copco	Diesel/ Electricity/DG	Air, noise, emission pollution
2.	Jack hammer	10	33mm dia.	Atlas Copco	-	Air, noise, dust pollution
3.	Hydraulic rock breaker/Excavator	2	1.2 cu. M or 380 HP	Sany – SY210C-9 or Tata Hitachi	Diesel	Air, noise, dust, emission pollution, ground vibration
4.	Tipper	2	20 tons	Tata	Diesel	Air, noise, dust, emission pollution, ground vibration
5.	Water tanker	1	5000 liter	Mahindra	Diesel	Air, noise, dust, emission pollution
6.	Tractor	1	04 tons	Mahindra	Diesel	Air, noise, dust, emission pollution
7.	JCB	1	---	JCB	Diesel	Air, noise, dust, emission pollution
8.	Diamond wire saw machine	2	40 and 60 hp.	---	Electricity	Air, water, dust pollution
9.	Quarry core drilling machine	1	---	---	Diesel/ Electricity/DG	Air, dust, noise, water pollution
10.	Diesel generator	2	320 KVA and 26 KVA	Cummins	Diesel/Electricity	Air, emission, noise pollution
11.	Accessories	---	All size drilling rods and other machine spare parts	---	---	---

leaves and cuticles. Dust may also have chemical effects that jeopardize the plants' long-term survival (Tiba 2017, Ogbonna et al. 2018, Timofeeva et al. 2022, Abdul-Wahab et al. 2022).

Quarry dust is linked to a multitude of leaf diseases, including abscission (early leaf fall), epinasty (bending of the leaf downward due to a faster rate of growth on the upper surface), necrosis (death of a portion of the leaf), and chlorosis (yellowing of the leaf due to loss or decrease of chlorophyll). Quarry dust is also a major source of air pollution (Sanjay VEDIYA 2014, Sayara et al. 2016, Sairanen et al. 2018 & 2019, Ekpa et al. 2022 & 2023, Opondo et al. 2023). According to Abayomi et al. 2014, Hamzart-Giwa et al. 2023, the quantity and quality of resource discovery, extraction, stonecutting procedures, and transportation to the final destination determine the impact on the air environment. According to Busuyi (2008), Olusegun et al. (2009), Akanwa et al. (2016 & 2017), Abhishek Pandey (2018) the rate of production and transportation also affects the intensity of these processes.

In the regions where stone crushing, quarrying, and stone processing are conducted, some of the sources of emissions or dust creation are as follows: dust from drill bits and blasting; dust from stone bodies and objects dug out and extracted. Transporting waste and dressing materials or processing materials from quarries to stockyards and disposal sites; loading, unloading, and moving finished items; and movement of vehicles on haul highways (Thompson et al. 1984, Iqbal et al. 2001, Moibi 2007, Ojeaga et al. 2023). A region's wind direction and speed have a direct bearing on the probability of dust pollution spreading in the study area.

The research area's major wind direction is west, and during the AAQ monitoring period, data on wind speed and direction were concurrently gathered. The ambient air quality of the research region has been evaluated using a network of twelve ambient air quality stations. The study region's meteorological characteristics were taken into consideration when building these stations, along with several other elements such as a densely populated area, industrial and residential development, and environmental sensitivity. It was noted that there are no ecosystems surrounding the intended exploratory well placements, and all of the wells are situated inside open spaces.

The principal inhabited the region and the predominant wind directions were considered when choosing the AAQ sites. Baseline ambient air quality evaluations provide information about the site's surroundings and are an essential component of research on environmental impact assessments. In addition to the impact of the surrounding topography, notable variations in the predominant winds and weather

patterns are seen in the winter, summer, and post-monsoon seasons. A methodical program of air quality surveillance is used to evaluate the baseline air quality in the research region.

The quarrying site reported temperatures as high as 33°C and as low as 19°C during the research period. It was found that the quarry zone had an average temperature of 26°C. Throughout the trial, the average relative humidity ranged from 34% to 63%. In the Ramanagara District, there was some rainfall during the study period. There were five days of rain on average. The most rain that fell in March was 12 mm, while the least was 8 mm. The district receives roughly 931.58 mm of rain annually on average. During the winter monsoon, the majority of winds are north-easterly, whereas during the summer monsoon, they are south-westerly. During the three-month investigation, monthly wind speeds in the quarry zones ranged from 4.9 m/s to 5.6 m/s.

The AERMET data includes daily records for temperature, relative humidity, air pressure, solar radiation, precipitation, wind speed, and wind direction. AERMET reformats meteorological data to make it suitable for use as input in the AERMOD model (Neshuku 2012). The stone quarrying and stone-processing sectors won't alter the climate. When evaluating and keeping track of ambient air quality, soil and geological features are crucial.

Air Quality Index (AQI) Quantification

Monitoring sites have been reporting the ambient air pollution they have observed within a specific time frame (e.g., daily) by utilizing a grading system based on the AQI universal standard. The AQI is used to compare current pollution levels to ensure that impact reduction requirements are being followed and to increase public awareness of the risks associated with regular exposure to pollution. Even though the AQI is just a number that represents some aspects of air quality, to make its meaning clear to the general public, it is combined with various messaging, color schemes, graphics, and labels for the different categories of air quality (such as "good," "moderate," or "hazardous" as per CPCB (200) guidelines).

Based on a set of standards, the Air Quality Index (AQI) is an environmental indicator that provides information on the general condition and trend of the ambient air in a given area. This tool uses the weighted values of different air pollution parameters to create a single or multiple-number representation. The AQI is a useful tool for assessing and measuring the overall ambient air quality of a particular area since it accounts for the combined impact of all contaminants. The AQI can also be used to control equipment that, for example, raises the concentrations of some pollutants while reducing

Table 9: Air Quality Index grading system

AQI Value	Remarks	Health Concern
00 – 25	Clean air (CA)	None/minimal health effect
26 – 50	Light air pollution (LAP)	Possible respiratory or cardiac effect for most sensitive group, skin allergies, cough etc.
51 – 75	Moderate Air Pollution (MAP)	Increasing symptoms of respiratory and cardiovascular illness, skin allergies, cough, asthma, etc.
76 – 100	Heavy Air Pollution (HAP)	Aggravation of heart and lung diseases, asthma, eye irritation, skin diseases, etc.
> 100	Severe Air Pollution (SAP)	Serious aggravation of heart and lung Diseases, risk of death in children, asthma, cough, eye diseases, mental disorder etc.

others, or to create alternate strategies for preventing air pollution.

The AQI can be calculated using a variety of formulas and methods. Nonetheless, the following calculation has been used in this instance to determine the AQI value:

$$AQI = \frac{1}{4} \times (ISPM/SSPM + ISO_2/SSO_2 + INO_x/SNO_x) \times 100$$

Individual measurements obtained during sampling that correlate to ISPM, IRSPM, ISO₂, and INO_x are respirable particulate matter, suspended particulate matter, sulfur dioxide, and oxides of nitrogen, respectively. For ambient air quality, the Central Pollution Control Board of India (CPCB) has set SSPM, SRSPM, SSO₂, and SNO_x criteria. Higher air pollution levels and associated health risks are correlated with higher AQI numbers. Table 9 lists the five categories that comprise the AQI scale. It outlines the range of air quality as well as any possible health hazards.

Examination of the Surrounding Air Quality in the Study Area

Sampling was done after GPS and remote sensing pinpointed the precise locations. Samples and measurements of the chemical components of atmospheric pollutants were taken at twelve different locations using an air particle counter and a fuel gas analyzer. Every location had an air particle counter accessible to measure SPM. The SPM concentration was printed and recorded by the instrument. A fuel gas analyzer was used to determine the extra air pollutants (NO_x,




Fig. 5: Sites of ambient air quality monitoring.

SO_x, H₂S, and C_xH_x). The equipment was calibrated and constructed correctly, and additional equipment was utilized to print the read-off of the atmospheric concentration of each gas on a screen.

The direction and speed of the wind were measured using an electrically powered digital anemometer. The selection criteria for monitoring locations include the following: topography and terrain; meteorological circumstances (both upwind and downwind locations); residential and sensitive areas within the study area; regional baseline air quality and pollution levels; and representation of possibly impacted areas.

Throughout one (1) season or three months (January 2024 to March 2024), the ambient air quality in the research region was measured twice a week by the NAAQS for twelve parameters (PM₁₀, PM_{2.5}, SO₂, NO_x, CO, O₃, Pb, NH₃, C₆H₆, C₂OH₁₂, As, and Ni). The Central Pollution Control Board's (CPCB) monitoring criteria were followed when sampling was done at each location. PM₁₀, PM_{2.5}, SO₂, NO_x, CO, Pb, O₃, NH₃, C₆H₆, C₂OH₁₂, As, and Ni, maximum values at all study locations are well under the National Ambient Air Quality Standards for residential, commercial, and industrial areas (Tiwari et al. 2012).

Throughout the study, the average baseline levels of PM₁₀ (37.17 to 70.52 µg/m³), PM_{2.5} (16.98 to 39.85 µg/m³), SO₂ (5.29 to 13.91 µg/m³), NO_x (9.8 to 29.71 µg/m³), CO (0.15–0.32 mg/m³), O₃ (6.9 to 15.37 µg/m³), and some BDL and LOQ were found to be well within the National Ambient Air Quality Standards for both commercial, residential and industrial areas. Places to monitor ambient air quality are depicted in Fig. 5. The observed data for each location was used to calculate

several measures, including the maximum, minimum, and average.

Findings from the Analysis of Ambient Air Quality

The quarrying and stone processing industries are known to produce air pollution issues due to many reasons, such as truck traffic, blasting, and drilling. Small-scale quarrying is unlikely to significantly impact the current air quality in the core zone. Using environmental controls in a way that reduces the likelihood of contamination is necessary to reduce air pollution. The test results that are enclosed concern the surrounding area and ambient air quality at the quarry. Using the observed data for each location, several metrics were computed, including the maximum, minimum, and average. The overall summary of the ambient air quality test findings is shown in Table 10. Fig. 6 shows that the graphical representation of air quality monitoring results.

Observations in the Sampling Areas

PM₁₀ concentrations were measured to be at a maximum of 70.52 µg/m³ and a minimum of 37.17 µg/m³ respectively. The village of Harohalli had the highest concentration, while the village of Achalu recorded the lowest. The range of average values was 43.225 µg/m³ to 60.00µg/m³. The PM_{2.5} values were measured to be 39.85 µg/m³ at the maximum and 16.98 µg/m³ at the minimum. The village of Harohalli had the highest concentration, while the village of Achalu recorded the lowest. The measured average values fell between 20.655 µg/m³ to 33.905 µg/m³. The measurements of SO₂ concentrations at the maximum and minimum were 13.91µg/m³ and 5.29µg/m³ respectively. The




Fig. 6: Graphical representation of air quality monitoring results.

Bidadi industrial sector recorded the highest concentration, while the Yadamaranahalli quarry site recorded the lowest. The measured average values fell between the ranges of $6.415 \mu\text{g}/\text{m}^3$ to $12.735 \mu\text{g}/\text{m}^3$. There was a maximum of $29.71 \mu\text{g}/\text{m}^3$ and a low of $9.8 \mu\text{g}/\text{m}^3$ for the concentration of NO_x . The settlement of Achalu village recorded the lowest concentration, while the village of Harohalli recorded the largest concentration. From $11.85 \mu\text{g}/\text{m}^3$ to $25.28 \mu\text{g}/\text{m}^3$, the average levels were found to be present.

The observed CO values were $0.32 \text{ mg}/\text{m}^3$ at the maximum and $0.15 \text{ mg}/\text{m}^3$ at the minimum. The Doddamudawadi quarry site had the lowest concentration, whereas the Bidadi industrial area had the highest concentration. It was noted that the average values fell between $0.13 \text{ mg}/\text{m}^3$ and $0.245 \text{ mg}/\text{m}^3$. The reported values for O_3 concentrations were $15.37 \mu\text{g}/\text{m}^3$ for the maximum and $6.9 \mu\text{g}/\text{m}^3$ for the minimum. The Bidadi industrial sector had the highest concentration, while the Yadamaranahalli quarry site recorded the lowest concentration. The observed average values fell between $8.5 \mu\text{g}/\text{m}^3$ and $14.08 \mu\text{g}/\text{m}^3$. The Central Pollution Control Board's (CPCB) guidelines for industrial, rural, residential, commercial, and other areas are well under the limits in the research sites, while the concentrations of NH_3 , Pb, Ni, As, C_6H_6 and C_2OH_{12} were all below detectable levels at all locations.

Numerous factors and parameters, such as the climate, topography, extent and method of quarry operation, use of equipment and machinery, population density, density of highly polluted industrial areas, density of road vehicles, density of forest fires, density of vegetation, and so on, affect the distribution of the aforementioned chemical elements in different quarry locations, local areas, residential areas, and industrial areas. Air pollution levels are higher throughout the entire area as a result of the previously described factors. The observation detailed the areas where the lower and higher levels of air pollution were spreading at the sampling locations.

Air Pollution Effects on Human Health

Quarrying is viewed as essential to human survival on a global scale. This is because it has contributed to the growth and development of every country, region, and sub-region. As a result, it is easier to build concrete structures, and the state's road and bridge infrastructure is developed, both of which have a significant positive economic impact on the nation (William et al. 2006, Sahu et al. 2018, William 2020). Furthermore, it produced jobs that were both direct and indirect, which made it easier for individuals to find work and support themselves (Peter et al. 2018, Samba et al. 2022). The study found that because it gave individuals economic

power and maintained their well-being, quarrying was an especially sustainable industry (Ako et al. 2015, Ugbogu et al. 2009). This implies that, in addition to agriculture, quarrying provides a sizeable portion of the local population's income (Fugiel et al. 2017, Yasobant et al. 2017).

The local people, the environment, wildlife, workers, the population's socioeconomic situation, health, and society are all impacted by the various barriers, problems, and challenges that come with stone quarrying (Nartey et al. 2012, Turyahabwe et al. 2021). These problems include changes in topography, degradation, and abandonment of land, loss of ecosystems and biodiversity, emissions, noise and dust pollution, air pollution, and water pollution, including the release of ionizing radiation into the atmosphere (Sanjay Vediya 2014, Pal et al. 2019). These results indicate that places that were productive in the past have become unproductive, which is upsetting the ecosystem as a whole.

Deforestation and land degradation therefore affect biodiversity (Rani et al. 2017). Due to their role as mosquito habitats and rain-retaining materials, open quarries and deforestation have been associated with an increase in non-communicable diseases, including malaria. This also affects the cost of buying and selling real estate (Owolabi et al. 2020).

The results of the study show how much dust is produced throughout several stages of the quarrying process, including drilling, stone blasting and crushing, loading, and transporting the finished products. The larger dust particles settle close to one another, while the thinner ones spread out widely, depending on the direction and speed of the wind. Different levels of particulate matter (PM) are produced based on its size, which also affects how long and how far it travels when suspended in the air (Romo-Kroger et al. 1989, Pradhananga et al. 2020).

Based on their aerodynamic diameter, the sizes are divided into $\text{PM}_{2.5}$ (equal to or less than 2.5 micrometers) and PM_{10} (equivalent to or fewer than 10 micrometers) particles. The average diameter of human hair is between 50 and 70 μm ; therefore, these sizes are less than that (Ogbonna et al. 2020, Chamdimba et al. 2023). Dispersed particulate matter can cause several issues, depending on its size. According to William (2020) and Ekpa et al. (2022 & 2023), for example, it can lead to poor visibility near quarry and crusher units, reduced agricultural yields because dust cover on plant surfaces blocks light needed for photosynthesis, ophthalmic disease because particulates carry pathogens, and respiratory disorders in humans and animals.

It is common knowledge that employees in quarries disobey health and safety; laws. People often forget to wear safety gear, like dust masks, helmets, and appropriately



Fig. 7: The safety and health conditions of the workers at the quarry sites and field visit images from several quarries and crusher unit.

protected clothing (Sanjay VEDIYA 2014, Sayara et al. 2016, Hassan 2023). As a result, they come into contact with small dust particles. Their health is consequently adversely affected (Vandana et al. 2020, William et al. 2006). Numerous health problems, including lung infections, skin and eye infections, respiratory and pulmonary disorders, and lung collapse, can be brought on by dust exposure (Ndinwa et al. 2014, Mbandi 2017, Nemer et al. 2020).

The Ramanagara district quarry workers found that their lack of protective gear and the dust from the quarries had caused respiratory difficulties. In addition to dyspnea, these symptoms before and after the quarry operation include coughing, wheezing, asthma, headaches, eye issues, chest discomfort, heart problems, mental stress, throat infections, allergies, and skin disorders (Sinha et al. 2000, Singhal et al. 2022, Abeya 2023). Based on the previously indicated facts—that dust is irksome, that dust has an impact on health, and

Table 11: The summary of air pollution impacts on human health

Pollutants	Environmental impacts on human health
Ozone (O ₃)	Ozone is linked to brief effects on the respiratory system in humans especially asthma, including reductions in pulmonary function in those engaging in mild to moderate exertion., various diseases and early mortality. Ozone depletion damages vegetation and reduces visibility, among other negative effects.
Particulate matters (PM _{2.5} & PM ₁₀)	Particulate matter has varying effects on health, and as particle size drops, so does the likelihood that the particle will have an impact on human health. Particles smaller than 1 µm entered the lower parts of the lungs, while larger particles lodged in the throat and nose, measuring more than 10 µm. Death, both acute and chronic bronchitis.
Sulphate (SO ₄)	Chest troubles, respiratory disorders, mild RADs, days missed from work, and moderate to severe asthma symptoms are all examples of lower and upper respiratory illnesses.
Carbon monoxide (CO)	As a result of its interaction with hemoglobin in human blood, it lowers the blood's transport of oxygen. When CO levels are high, those who have cardiovascular illness or chronic heart disease may feel pain in their chest. Reduced delay to angina onset, hospital admissions, mortality, and congestive heart failure. CO can cause unconsciousness and death, damage manual dexterity and vision.
Nitrogen oxides (NO ₂)	Short- and long-term impacts of nitrogen dioxide on human health are evident; children's respiratory disorders are more common in the former case, while the latter reduces immunity to respiratory infections. In addition to lowering auxiliary beat frequency and increasing permeability of cell membranes, NO ₂ exposure also makes asthmatics more sensitive to respiratory infections and inhaled allergens.
Sulphur dioxide (SO ₂)	Another gas that negatively affects human health is sulfur dioxide, since it can have a severe effect on the respiratory system. Asthma sufferers' lung function is altered by the emission, which is directly correlated with fuel's sulfur content. Sensitive people's respiratory symptoms are made worse. Respiratory symptoms, alterations in pulmonary function, and morbidity among asthmatics who exercise.
Lead (Pb)	Deaths, high blood pressure, non-fatal heart attacks, non-fatal strokes, and declines in IQ.

that rain-covered roofs are unclean-the impact of quarrying on human health is examined. The safety and health conditions of the workers at the quarry sites are depicted in the Fig. 7, which also include field visit images from several quarries and crusher unit.

The results demonstrated that all respondents (100%) considered noise and dust to be health hazards and a major source of annoyance. Dust almost always contaminates rain that falls from roofs. Additionally, every respondent mentioned that noise, dust, water, and floodwaters could be connected to their on-going health issues. Numerous illnesses and health problems are signs of the detrimental impacts of stone processing and quarrying, particularly for the villages close to crushers. There is also a connection between acid rain and quarrying (Maduka et al. 2014, Boutemedjet et al. 2019).

According to the study, acid rain is known to damage plants, worsen air pollution, and sicken people and animals brought on by quarrying. There are several quarrying-related health risks. An analysis showed a connection between the vibrations caused by quarrying and fractures. This could lead to building collapses, which would cause fatalities and injuries. When asked how they felt about the quality of the air right now, respondents said that 24% thought it was poor and 76% said the stone-cutting and quarrying sectors should be held accountable. Since the quarry owner is providing benefits, the majority of respondents chose not to react. Thus, it seems logical that stone-cutting and quarrying have an impact on air quality. Table 11 provides a summary of how typical air contaminants affect human health.

TECHNIQUES FOR MITIGATION AND CONTROL MEASURES

The following mitigation and control techniques, corrective measures, and remedies have been suggested for the granite quarrying and stone processing industries: a sizable buffer zone, or "green belt," ought to be created surrounding the industrial area. Providing dust masks for laborers. Allowing the staff to don the appropriate personal protective equipment. Avoid overcharging blast holes by using controlled blasting techniques. It is advisable to regularly monitor the quality of the surrounding water, air, and noise. It is also suggested that you use effective, quiet equipment. Hazardous places should be avoided, and signboards displaying instructions should be posted.

Dust pollution will be managed with the use of electrostatic precipitators. By using silica in water quarrying and groundwater dust control systems, silica-related problems can be minimized. At sensitive locations, such as haul roads, crusher and screening plants, mineral handling

facilities, quarrying facilities, and stone processing plants, dust suppression is enforced using heavy dust sprinklers and road watering trucks. Mineral handling, HEMM, crushing, and screening equipment must be provided in dust extraction facilities. Hoods, water sprays, and dust collectors are helpful pieces of equipment for controlling drilling dust. Moreover, suitable chute design, vulcanizing conveyor belt joints, under-belt cleaning equipment, hoods, transfer points, and dust suppression and/or extraction systems for conveyors are frequently employed as dust pollution management techniques. Enclosures used in mineral handling plants must be well secured.

All vehicles, including trucks, dumpers, and transport vehicles, must be leak-proof. Use the proper components when spraying to stop dust from flying. Highways and other roads should be adequately coated with chemical additives to successfully limit dust emissions. Haul roads and service roads should be graded regularly. Haul roads should also have any loose dirt and collected dust removed. Appropriate maintenance, such as checking exhaust emissions and establishing speed limitations on the cars and earthmoving equipment. When transporting stone products, make sure tippers and dumpers are not overloaded, and cover heavy tippers with tarpaulins.

Utilizing vegetation to stabilize significant regions that generate dust and waste yards. Reduce the number of open regions and seal off the area used for stone quarrying and processing to reduce the amount of dust produced. Ideal enclosed areas for equipment and supply storage for building projects. Utilizing and maintaining machinery and equipment efficiently to lower air emissions, noise pollution, and energy use. The exhaust vent of the DG set will have enough stack height to ensure that gaseous emissions diffuse rapidly. Vehicles must receive routine maintenance and oversight to guarantee that their fuel usage and emissions are within permitted limits. Inspecting construction sites, stone processing plants, and quarry activities regularly to ensure that waste debris is promptly removed and disposed of in landfills, recycled, or used in other ways. Grading transport and service roads and removing accumulated dusty material regularly.

CONCLUSIONS AND RECOMMENDATIONS

Conclusions

We start with an analysis of the field work, laboratory and desk work data in order to track the quality of the air in the Ramanagara district. Twelve metrics, including PM_{2.5}, PM₁₀, SO₂, NO_x, CO, O₃, Pb, NH₃, C₆H₆, C₂OH₁₂, As, and Ni, were monitored twice a week for three months (January

2024 to March 2024) by the National Ambient Air Quality Standards (NAAQS) in the study area. This indicates the greater distance that the air quality covers. According to the Central Pollution Control Board (CPCB) monitoring recommendations, sampling was carried out at each location. The NAAQS in the research area measured twelve metrics twice a week for one (01) season, or three months (January 2024 to March 2024), including PM₁₀, PM_{2.5}, SO₂, NO_x, CO, O₃, Pb, NH₃, C₆H₆, C₂OH₁₂, As, and Ni. The findings showed that SPM is high in every station, even if SO₂ and NO_x levels are within allowable bounds in all measured locations. The study found that the average baseline levels of PM₁₀ (37.17 µg/m³ to 70.52 µg/m³), PM_{2.5} (16.98 µg/m³ to 39.85 µg/m³), SO₂ (5.29 µg/m³ to 13.91 µg/m³), NO_x (9.8 µg/m³ to 29.71 µg/m³), CO (0.15 mg/m³ to 0.32 mg/m³), O₃ (6.9 µg/m³ to 15.37 µg/m³), and NH₃, Pb, Ni, As, C₆H₆ and C₂OH₁₂ were below detectable limit (BDL) and limit of quantification (LOQ) are all well within the National Ambient Air Quality Standards for commercial, industrial, and residential areas at all monitoring locations for the duration of the study. Apart from the environment, they can also have a major detrimental impact on infrastructure, transportation, agriculture, population density, mining, quarrying, human health, etc. The research areas' stone-processing and granite-quarrying businesses don't significantly harm the air quality. The distribution of air quality source potential in Ramanagara district was evaluated in this research work.

Recommendations

1. Environmental preservation is of utmost importance to the stone quarrying and processing industry. The business will abide by all environmental regulations. A well-manicured greenbelt will be preserved by the quarry operation. Furthermore, all environmental statute standards shall be continuously implemented and upheld.
2. By introducing both natural and artificial techniques to stop the spread of dust in the neighborhood, air pollution caused by billowing dust can be avoided. One way to lessen the harm that the dust from the quarries causes to the environment is to plant densely packed, quickly growing trees around the quarries and on any reserved area they own. The same quickly growing trees with dense foliage should also be added to the proposed buffer zone to increase its density. Rather than using corrugated sheets attached to the side of crushing plants in certain quarries, man-made and natural solutions can be implemented to reduce the amount of dust that blows into the area, thereby mitigating the problem of air pollution caused by billowing dust.
3. In order to improve the green canopy, prevent the spread of dust and other airborne pollutants, stop soil erosion, stop land degradation, and address other issues, reforestation is mandated by the quarrying and stone processing industry plan.
4. There should be frequent environmental audits carried out on all the quarries within the research area. In order to evaluate the effectiveness of the quarries' conservation or anti-pollution program, a methodical analysis of their interactions with the environment is conducted.
5. Regular health awareness efforts should also be conducted to inform the public about the possible dangers of unhygienic surroundings. To help enhance the health status of its residents, the local government should mandate frequent health assessment studies.
6. Local government agencies should be held to a higher standard of accountability and should take an active role in resolving public complaints regarding the detrimental effects of the granite business. Steer clear of quarrying in environmentally vulnerable areas.

REFERENCES

- Aarthy, A., Sudarsan, J.S., Wilfett, J. and Rajan, R.J., 2018. Study on impact of air quality due to quarrying activity in and around Rathinamangalam area of Kancheepuram district, Tamil Nadu, India. *Journal of Mines, Metals, and Fuels*, 66(1).
- Abayomi, D. and Kehinde, I., 2014. Environmental effects of quarrying activities in Oba-Ile, Akure, Ondo State, and South-West Nigeria. *International Journal of Engineering Research & Technology*, 3(1), pp.1103-1113.
- Abdul-Wahab, S., Osman, I. and Al-Shukaili, N., 2022. Air quality assessment for Kunooz Gabbro quarry in Oman: assessment of PM10. *International Journal of Pioneering Technology and Engineering*, 1(02), pp.61-72.
- Abdul-Wahab, S.A., Fgaier, H., Elkamel, A. and Chan, K., 2015. Air quality assessment for the proposed Miller Braeside quarry expansion in Canada: TSP. *Air Quality, Atmosphere & Health*, 8, pp.573-589.
- Abeya, C., 2023. Impacts of stone quarrying on the environment and socio-economy of the local community: the case of Aira District, West Wollega Zone. *Journal of Science, Technology, and Arts Research*, 12(1), pp.90-108.
- Ajah, J.N. and Chukwu, E.N., 2018. Investigative effect of stone crushing dust on the health of workers of quarry industry at Umuoghara community of Ezza-North local government in Ebonyi State. *IOSR Journal of Nursing and Health Science (IOSR-JNHS)*, 7(6), pp.80-91.
- Ajibade, O.M., Olisa, O.G., Oladipupo, S.D., Adegoke, C.B. and Adebayo, O.H., 2022. Health impact assessment of quarrying in some parts of southwestern Nigeria. *Arabian Journal of Geosciences*, 15(8), p.777.
- Akanwa, A., Onwuemesi, F. and Chukwurah, G., 2016. Effects of open-cast quarrying techniques on vegetation cover and the environment in south-eastern Nigeria. *American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS)*, 21(1), pp.227-240.
- Akanwa, A.O., Okeke, F.I., Nnodu, V.C. and Iortyom, E.T., 2017. Quarrying and its effect on vegetation cover for a sustainable development using high-resolution satellite images and GIS. *Environmental Earth Sciences*, 76(14), p.505.

- Ako, T.A., Onoduku, U.S., Waziri, S.H., Adegbe, M., Chukwu, J.N. and Kajena, C.M., 2015. Assessment of the environmental impacts of marble quarrying on surface water at Kwakuti, Niger State, North Central Nigeria. *International Journal of Engineering and Advanced Research Technology*, 1(1), pp.64-70.
- Amankwah, E.O., Owusu-Boateng, G. and Maxwell, A.B., 2019. Trend analysis on air quality and lung function—a case of Consar Stone Quarry Limited, Barekese. *American Journal of Environmental Resource Economics*, 4, pp.44-53.
- Bada, B.S., Olatunde, K.A. and Akande, O.A., 2013. Air quality assessment in the vicinity of the quarry site. *Environment and Natural Resources Research*, 3(2), p.111.
- Bakamwesiga, H., Mugisha, W., Kisira, Y. and Muwanga, A., 2021. An assessment of air and water pollution accrued from stone quarrying in Mukono District, Central Uganda.
- Bassey, P.E. and Okon, N.P., 2021. Perspectives and experiences of the effects of quarrying on the environment and agricultural activities of the quarry communities in Akamkpa Local Government Area of Cross River State, Nigeria. *International Journal of Science and Academic Research*, 2(07), pp.1767-1773.
- Bhat, P.G.K. and Subramani, N., 1990-91. Report on the Second Generation Geological Mapping in Parts of Satnuru—Kadihalli Area, Kanakapura Taluk, Bangalore District, Karnataka. (Unpublished Progress Report for 1990-91).
- Bhat, S.K. and Suresh Babu, V., 1994. Survey for ornamental stones with special reference to pink granite in parts of Kanakapura taluk, Bangalore district. Department of Mines and Geology, Karnataka. (DMG Reports).
- Boutemedjet, A., Bounouala, M., Idres, A. and Benselhou, A., 2019. Assessment of dust pollution related to granite quarry operations in Kef Bouacida, Annaba (Algeria). *Scientific Bulletin of the National Mining University*, (1).
- Busuyi, A.T., 2008. Assessment of the socio-economic impacts of quarrying and processing of limestone at Obajana, Nigeria. *European Journal for Social Sciences*.
- Caruthers, R.M., 1984. Reviews of geophysical techniques for groundwater exploration in crystalline basement Terrain. British Geological Survey Report No. RGRG85/3. Assessment of physico-chemical parameters of groundwater quality in Ramanagaram taluk, Karnataka. (DMG Reports).
- Chaanda, M.S., Onwuteaka, D. and Garba, M.A., 2019. Environmental Impact of Granite Mining in Tattara Area, North Central Nigeria. *IOSR Journal of Applied Geology and Geophysics I*, 7(6), pp. 14-24.
- Chalekode, P.K., 1978. Emissions from the Crushed Granite Industry: State of the Art. (Govt Reports).
- Chamdimba, G., Vunain, E. and Maoni, M., 2023. Assessment of particulate matter exposure to ambient air and its impact on workers at two granite quarry mines at Njuli, Southern Malawi. *Environmental Monitoring and Assessment*, 195(9), p.1069.
- Chattopadhyay, S., Gupta, S. and Saha, R.N., 2010. Spatial and temporal variation of urban air quality: a GIS approach. *Journal of Environmental Protection*, 1(03), pp.264.
- Chowdhury, M.Z., 2004. Characterization of fine particle air pollution in the Indian subcontinent. Georgia Institute of Technology.
- Daspan, R.I., Obadiah, E.G., Lekmang, I.C., Dibal, H.U., Chup, A.S., Daku, S. and Azi, B., 2018. Preliminary Assessment of the Trace Element Composition of Dust from Two Granite Quarries on the Jos Plateau and Their Possible Health Implications.
- Doley, D. and Rossato, L., 2010. Mineral particulates and vegetation: Modelled effects of dust on photosynthesis in plant canopies. *Air Quality and Climate Change*, 44(2), pp.22-27.
- Drew, L.J., Langer, W.H. and Sachs, J.S., 2002. Environmentalism and natural aggregate mining. *Natural Resources Research*, 11, pp.19-28.
- Ekpa, I.D., Laniyan, D.G., Agbor, C.N., Ben, U.C. and Okon, J.E., 2022. Effect of air pollution from quarry activities on agriculture and plant biodiversity in south-eastern Nigeria (Preprint). <https://doi.org/10.21203/rs.3.rs-1819483/v1>
- Ekpa, I.D., Laniyan, D.G., Agbor, C.N., Ben, U.C. and Okon, J.E., 2023. Effect of particulate matter from quarry activities on crops and plant biodiversity in south-eastern Nigeria. *Environmental Monitoring and Assessment*, 195(7), p.837.
- Emmanuel, D.A., 2018. Environmental impact of quarry activities in Supare-Akoko, Ondo State, Nigeria. *Geography Planning Science*, 3(1), pp.101-111.
- Esguerra, N.A., Amistad, F.T. and Rabena, A.R., 2008. Characterizing the environmental effects of the quarrying industry: The case of strategic quarry sites in the Ilocos Region. *The Vector: International Journal of Emerging Science, Technology, and Management (IJESTM)*, 17(1).
- Eshiwani, F., 2014. Effects of quarrying activities on the environment in Nairobi County: A Case Study of Embakasi District. (Doctoral Dissertation, University of Nairobi).
- Fugiel, A., Burchart-Korol, D., Czaplicka-Kolarz, K. and Smolinski, A., 2017. Environmental Impact and Damage Categories Caused by Air Pollution Emissions from the Mining and Quarrying Sectors of European Countries. *Journal of Cleaner Production*, 143(2), pp.159-168.
- Ganesha, A.V., Krishnaiah, C., Krishnegowda and Suresh, S., 2017. Assessment of physico-chemical parameters of groundwater quality in Ramanagaram taluk, Karnataka. *Special Publication Journal of Engineering Geology, EGCON*, pp.255-263.
- Gebe, B., 2013. Environmental Impacts of Construction Aggregate Mining in the Greater Accra Region (A Case Study of Amasaman in the Ga West Municipality). (Doctoral Dissertation).
- Halwenge, J.A., 2015. Dust pollution and its health risks among rock quarry workers in Kajiado County, Kenya. Available at: <https://pdfs.semanticscholar.org/ee32/fb69d83d6d678c07ebdccc590e426a4646.pdf> [Accessed 1 Mar. 2019].
- Hamzart-Giwa, O.G., Odiyi, B.O. and Afolabi, O.J., 2023. Environmental Effects of Quarrying Activities in Akure, Ondo State, Nigeria. *World News of Natural Sciences*, 49, pp.122-136.
- Haseeb, Z., 2018. Monitoring of the environment, health, and socio-economic impacts of limestone quarrying near Chakwal, Pakistan. *Pakistan Journal of Science*, 70(4).
- Hassan, I.A 2023. Socio-economics and health impact of quarry sites on the surrounding localities in Ogun State, Nigeria. *Nig. J. Pure & Appl. Sci.*, 36(2), pp. 4638-4655. : <https://doi.org/10.48198/NJPAS/23.A04>
- Hema, H.C. and Govindaiah, S., 2012. Morphometric analysis using remote sensing and GIS techniques in the sub-watersheds of Kanakapura watershed, Arkavathi river basin, Ramanagara district, Karnataka, India. *Environmental Geochemistry*, 15(2), pp.47-56.
- Hema, H.C., Govindaiah, S., Jayaramaiah, R. and Shahinfar, H., 2013. Remote sensing and GIS application for land use and land cover mapping in the Kanakapura watershed of Ramanagara district, Karnataka, India. *The Indian Mineralogist Spl*, 47(1 & 2), pp.41-57.
- Henry, A.E., Getrude, A.O., Chibuisi, O.F., Shu, E.N., Ignatius, C., Stella, I.A.I., and Halilu, T.B., 2017. Occupational health hazards associated with continuous exposure to quarry activities among quarry workers in Ebonyi State, South East Geopolitical Zone, Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, 11, pp.10-19.
- Ibrahim, W.H.W., Marini, E., Yunus, J., Asra, N. and Sukor, K.M., 2018. Air quality assessment on human well-being in the vicinity of the quarry site. In: *IOP Conference Series: Earth and Environmental Science*, 117(1), p.012010.
- Iqbal, M.Z. and Shafiq, M., 2001. Periodic effect of cement dust pollution on the growth of some plant species. *Turkish Journal of Botany*, 25(1), pp.19-24.
- Jahed Armaghani, D., Hajithassani, M., Monjezi, M., Mohamad, E.T., Marto, A. and Moghaddam, M.R., 2015. Application of two intelligent systems in predicting the environmental impacts of quarry blasting. *Arabian Journal of Geosciences*, 8, pp.9647-9665.

- Kittipongvises, S., 2017. Assessment of the environmental impacts of limestone quarrying operations in Thailand. *Environmental and Climate Technologies*, 20(1), pp.67-83.
- Kofi-Boye, R., 2017. Health Effects of Limestone Quarrying in Yilo Krobo and Lower Many Krobo Municipalities. (Doctoral Dissertation, Ensign Global College).
- Kuzu, C. and Ergin, H., 2005. An assessment of the environmental impacts of quarry-blasting operations: a case study in Istanbul, Turkey. *Environmental Geology*, 48, pp.211-217.
- Leon-Kabamba, N., Ngatu, N.R., Muzembo, B.A., Kakoma, S., Michel-Kabamba, N., Danuser, B. and Hirao, T., 2020. Air quality in the working environment and the respiratory health of female Congolese stone quarry workers. *Tropical Medicine and Infectious Disease*, 5(4), p.171.
- Loveson, V.J., Singh, T.N. and Singh, P.K., 1997. Impact of Noise Pollution on Workers' Health at Cement Plants. *Indian Mining & Engineering Journal*, 36(11), pp.31-34.
- Maduka, S.O., Osim, E.E. and Dimkpa, U., 2014. Respiratory symptoms and blood eosinophil level in workers exposed to quarry dust in south-eastern Nigeria. *Journal of Environmental and Occupational Science*, 3, pp.175-179.
- Martins, I.O., Osu Charles, I. and Iwuoha, G.N., 2017. Impact of stone quarrying on the health of residents in Nigeria. *Life Science Journal*, 14(10).
- Mbandi, I.J., 2017. Assessment of the Environmental Effects of Quarrying in Kitemanga Subcounty of Kajiado in Kenya. (Doctoral dissertation, University of Nairobi).
- Melodi, M.M., 2017. Assessment of the environmental impacts of quarry operations in Ogun State, Nigeria. *FUOYE Journal of Engineering and Technology*, 2(2), pp.2579-0625.
- Ming'ate, F.L.M. and Mohamed, M.Y., 2016. Impact of stone quarrying on the environment and the livelihood of communities in Mandera County, Kenya. *Journal of Scientific Research and Reports*, 10(5), pp.1-9.
- Moeletsi, R.S., 2018. Remote sensing of land cover change within granite quarries. (Master's thesis, University of Johannesburg (South Africa)).
- Moibi, N.M., 2007. Effects of quarrying on the environment: a case study of Tala, Machakos District, Kenya. (Research dissertation)
- Mwangi, S.N., 2014. An assessment of the environmental impacts of quarrying activities in Ndarugo Area, Kiambu County. Bachelor Degree Project, Kenyatta University, Kenya.
- Nandan, A., Tauseef, S.M. and Siddiqui, N.A., 2017. Assessment of ambient air quality parameters in various industries in Uttarakhand, India. In: *Materials, Energy, and Environment Engineering: Select Proceedings of ICACE 2015*. Springer Singapore, pp.279-290.
- Nartey, V.K., Nanor, J.N. and Klake, R.K., 2012. Effects of quarry activities on some selected communities in the lower Many Krobo District of the Eastern Region of Ghana.
- Ndinwa, G.C.C. and Ohwona, C.O., 2014. Environmental and health impact of solid mineral exploration and exploitation in south-northern Nigeria: a case study of IGARRA in EDO STATE. *Review of Environmental and Earth Sciences*, 1, pp.24-36.
- Nemer, M., Giacaman, R. and Husseini, A., 2020. Lung function and respiratory health of populations living close to quarry sites in Palestine: A cross-sectional study. *International Journal of Environmental Research and Public Health*, 17(17), p.6068.
- Neshuku, M.N., 2012. Comparison of the performance of two atmospheric dispersion models (AERMOD and ADMS) for open pit mining sources of air pollution. University of Pretoria (South Africa).
- Odewumi, S.C., Aminu, A.A., Momoh, A. and Bulus, J.A., 2015. Environmental Impact of Mining and Pedogeochemistry in the Agunjin Area, Southwestern Nigeria.
- Ogbonna, P.C., Kalu, E.N. and Nwankwo, O.U., 2018. Determination of heavy metals in sawdust particles, distribution in soil, and accumulation in plants at Abiake timber market in Umuahia, Nigeria. *Nigerian Journal of Environmental Sciences and Technology*, 2(2), p.160.
- Ogbonna, P.C., Ukpai, N.P. and Ubuoh, E.A., 2020. Assessment of particulate matter (PM_{2.5}, PM₁₀) in the air, elemental composition of granite, and weather parameters at a quarry site in Ngwogwo, Ebonyi State, Nigeria. *Nigerian Journal of Environmental Science and Technology*, 4(1), pp.182-196.
- Ojeaga, K. and Atufe, A., 2023. Assessment of PM_{2.5} Around Rock Quarrying Sites and Residential Communities of Iyuku and Ikpeshi, Estako West, Akoko Edo, and Edo State. *Journal of Science and Technology Research*, 5(2).
- Okafor, O.C., Njoku, C. and Akwuebu, A.N., 2023. Environmental impact of quarrying on air quality in Ebonyi State, Nigeria. *Environmental Sciences Europe*, 35(1), p.98.
- Olusegun, O., Adeniyi, A. and Adeola, G.T., 2009. Impact of granite quarrying on the health of workers and nearby residents in Abeokuta, Ogun State, Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 2(1).
- Omeiza, L., Sangari, D. and Ogah, A., 2022. Effects of quarry on the air quality in Mpape, Bwari Area Council, FCT Abuja, Nigeria.
- Oniyide, G.O., Idowu, K.A. and Anikoh, G.A. Investigations of the Environmental Effects of Granite Rock Quarrying (A Case Study of Ebenezer Quarry, Akure, Nigeria).
- Opondo, E.O., Ajayi, D.D. and Makindi, S.M., 2023. Impacts of quarrying activities on the environment and livelihood of people in Border II sub-location, Nyando sub-county, Kisumu County, Kenya. *Environmental Quality Management*, 32(3), pp.147-160.
- Owolabi, A. and Adesida, P., 2020. The Environmental and Health Implications of Quarrying Activities in the Host Community of Oba-Ile in Akure, Nigeria. *Journal of Human Environment and Health Promotion*, 6(1), pp.6-10.
- Oyinloye, M.A. and Ajayi, E.A., 2015. Environmental impact assessment of quarry activities in Oba-Ile, Ondo State, Nigeria. *The Journal of MacroTrends in Energy and Sustainable*, 3(1), pp.32-45.
- Pal, S. and Mandal, I., 2019. Impact of aggregate quarrying and crushing on socio-ecological components of the Chottanagpur plateau fringe area of India. *Environmental Earth Sciences*, 78(23), p.661.
- Pandey, A., 2018. Environmental impact and damage categories caused by air pollution emissions from mining and quarrying sectors of European countries. *Journal of Cleaner Production*, 97, pp.1008-1020.
- Paranthaman, S. and Ahmad, Md., 1994-1995. Resource Survey Of Ornamental Granite And Dike Rocks For Polishing And Dimensional Stone Industry In Parts Of Kanakapura Taluk, Bangalore Rural District, Karnataka. (GSI Reports).
- Peter, M.C., Alozie, C.E. and Azubuine, C.I., 2018. Stone quarrying impacts air and soil water in Ebonyi State, Nigeria. *Journal of Pollution Effects & Control*, 6(2). DOI: 10.4172/2375-4397.1000225.
- Pradhananga, P. and Elzomor, M., 2020. Environmental implications of quarry rock dust: a sustainable alternative material to sand in concrete. In: *Construction Research Congress 2020*, pp.916-924. Reston, VA: American Society of Civil Engineers.
- Rajan Babu, A., Behera, P.K. and Gupta, R.N., 2003. Eco-friendly stone quarrying-some Issues. In: *Seminar on safe & Environmental friendly quarrying*, Mining Engineers Association of India, Bangalore. (National Seminar).
- Rajan Babu, A., Behera, P.K., Subrahmanyam, D.S. and Gupta, R.N., 2002. Development of guidelines for improving recovery in granite mining operation. (Final report 2002. NIRM).
- Rajan Babu, A., Subrahmanyam, D.S. and Gupta, R.N., 2000. Performance improvement in dimensional stone quarrying through affordable scientific approach-A Case Study. In: *Internal conference on Technology Management for Mining, Processing, & Environment at IIT, Kharagpur*, pp.108-115. (Internal conference).
- Rani, H.B., Swamy, S., Bharath, A.L., Dinakar, R.P.H. and Raghu, A.V., 2017. Impact of quarrying and crushing on soil quality: a case study

- in Tumkur district, Karnataka. *International Journal of Research—Granthaalaya*, 5.
- Rathore, K.K., 2020. Environmental impacts of mining and processing of minerals: A review. *SGVU Journal of Climate Change and Water*, 7, pp.85-93.
- Roja, K., 2022. Environmental Impact Assessment of Mining and Quarrying in Madurai District.
- Romo-Kroger, C.M., Morales, R., Llonad, F., Auriold, P. and Wolleter, G.E., 1989. Risks of airborne particulate exposure in a copper mine in Chile. *Industrial Health*, 27(2), pp.95-99.
- Saha, D.C. and Padhy, P.K., 2011. Effects of the stone crushing industry on *Shorea robusta* and *Madhuca indica* foliage in Lalpahari forest. *Atmospheric Pollution Research*, 2(4), pp.463-476.
- Sahu, S.P., Yadav, M., Rani, N. and Das, A.J., 2018. Assessment of occupational health exposure to particulate matter around opencast coal mines in India: a case study. *Arabian Journal of Geosciences*, 11, pp.1-11.
- Sairanen, M. and Rinne, M., 2019. Dust emissions from the crushing of hard rock aggregates. *Atmospheric Pollution Research*, 10(2), pp.656-664.
- Sairanen, M., Rinne, M. and Selonen, O., 2018. A review of dust emission dispersions in rock aggregate and natural stone quarries. *International Journal of Mining, Reclamation, and Environment*, 32(3), pp.196-220.
- Salem, H.S., 2021. Evaluation of the Stone and Marble Industry in Palestine: environmental, geological, health, socioeconomic, cultural, and legal perspectives, in view of sustainable development. *Environmental Science and Pollution Research*, 28(22), pp.28058-28080.
- Samba, G., Fokeng, R.M., Nfor, J.T., Ngwaimbi, R.C. and Youogo, C.M.K., 2022. Effects of quarrying activities on environmental sustainability in Makenene, Centre Region, Cameroon. *Journal of Environmental Earth Science*, 12(1), pp.12-19.
- Sanjay Vediya, S.V., 2014. Dust pollution monitoring by higher groups of plants near the stone quarrying industry—a case study in Vadagam, Arravali District, Gujarat. *Lifesciences Leaflets*, 53, pp.57-60.
- Santizo, I.P., 2022. Impacts of Granite Quarrying: The Case of Subsistence Farmers in the São Pedro River Valley. (Doctoral dissertation, University of the Incarnate Word).
- Sayara, T., 2016. Environmental impact assessment of quarries and stone-cutting industries in Palestine: a case study of Jammain. *Journal of Environment Protection and Sustainable Development*, 2(4), pp.32-38.
- Sayara, T., Hamdan, Y. and Basheer-Salimia, R., 2016. Impact of air pollution from quarrying and stone-cutting industries on agriculture and plant biodiversity. *Resources and Environment*, 6(6), pp.122-126.
- Sheikh, A., Rana, S.V.S. and Pal, A., 2011. Environmental health assessment of stone crushers in and around Jhansi, UP, India. *Journal of Ecophysiology and Occupational Health*, pp.107-115.
- Simha, L. and Babu, M., 2015. Evaluation of Water Quality in and around Byramangala Reservoir, Ramanagaram District, Karnataka, India. *Evaluation*, 8(02), pp.215-221.
- Singh, G., Pal, A., Niranjan, R.K. and Kumar, M., 2010. Assessment of environmental impacts by mining activities: A case study from the Jhansi open-cast mining site in Uttar Pradesh, India. *Journal of Experimental Sciences*, 1(1).
- Singhal, A. and Goel, S., 2022. Impact of Sandstone Quarrying on the Health of Quarry Workers and Local Residents: A Case Study of Keru, Jodhpur, India. In: *Treatment and disposal of solid and hazardous wastes*. Cham: Springer International Publishing, pp.97-118.
- Singhal, A., 2018. Environmental Impacts of Sandstone Quarrying at Keru, Jodhpur, Rajasthan. Indian Institute of Technology, Kharagpur.
- Sinha, R.K., Pandey, D.K. and Sinha, A.K., 2000. Mining and the environment: a case study from the Bijolia quarrying site in Rajasthan, India. *Environmentalist*, 20, pp.195-203.
- Sreekala, S., Geena, P. and Jyothi, S.N., 2023. Environmental impact assessment of a quarry in a taluk in southern Kerala, India. In: *AIP Conference Proceedings*, 2764(1). AIP Publishing.
- Stieb, D.M., Doiron, M.S., Blagden, P. and Burnett, R.T., 2005. Estimating the public health burden attributable to air pollution: an illustration using the development of an alternative air quality index. *Journal of Toxicology and Environmental Health, Part A*, 68(13-14), pp.1275-1288.
- Sun, L., Wei, J., Duan, D.H., Guo, Y.M., Yang, D.X., Jia, C. and Mi, X.T., 2016. Impact of land use and land cover change on urban air quality in representative cities in China. *Journal of Atmospheric and Solar-Terrestrial Physics*, 142, pp.43-54.
- Sunyer, J.J.E.R.J., 2001. Urban air pollution and chronic obstructive pulmonary disease: a review. *European Respiratory Journal*, 17(5), pp.1024-1033.
- Suresh Babu, 1995. Survey and assessment of ornamental granite in parts of Magadi taluk, Bangalore district. Department of Mines and Geology, Karnataka. (DMG Reports).
- Swamy, M.M., 1998. A report on regional assessment and inventory of the resources of dimensional stones, commercial granites, and ornamental stones of closepet granites in parts of Kanakapura, Ramanagaram, and Magadi taluks, Bangalore rural districts, Karnataka. GSI, Southern Region (field season 1994-95), Bangalore. (GSI Unpublished Report).
- Syed Abrar, Mulgundmuth, S.F. and Badami, R.B., 2000. Assessment of Ornamental Stones in Ramanagaram Taluk, Bangalore District. Department of Mines and Geology, Bangalore (Field Season 1992-93). (DMG Reports).
- Syed Abrar, Smeulla, 2005. Ornamental Stone Resources of Channapatna Taluk, Bangalore. DMG, Karnataka. (DMG Reports).
- Thompson, J.R., Mueller, P.W., Flückiger, W. and Rutter, A.J., 1984. The effect of dust on photosynthesis and its significance for roadside plants. *Environmental Pollution Series A, Ecological and Biological*, 34(2), pp.171-190.
- Tiba, M.N., 2017. The environmental impact and risk generated by the extraction of construction rocks in quarries: a case study of Turcoia Quarry. *Research Journal of Agricultural Science*, 49(4).
- Timofeeva, S.S. and Murzin, M.A., 2020. Assessing the environmental risk of mining enterprises by the integral indicator of dust emissions. In: *IOP Conference Series: Earth and Environmental Science*, 408(1), p.012067. IOP Publishing.
- Tiwari, S., Chate, D.M., Srivastava, M.K., Safai, P.D., Srivastava, A.K., Bisht, D.S. and Padmanabhamurty, B., 2012. Statistical evaluation of PM10 and distribution of PM1, PM2.5, and PM10 in ambient air due to extreme fireworks episodes (Deepawali festivals) in megacity Delhi. *Natural Hazards*, 61, pp.521-531.
- Trivedi, R., Chakraborty, M.K. and Tewary, B.K., 2009. Dust dispersion modeling using a fugitive dust model at an opencast coal project of Western Coalfields Limited, India. *Journal of Scientific and Industrial Research*, 8, pp. 71-78.
- Trivedi, R., Mondal, A., Chakraborty, M.K. and Tewary, B.K., 2010. A statistical analysis of ambient air quality around opencast coal projects in Wardha Valley coalfields, Western Coal Fields Limited, India. *Indian Journal of Environmental Protection*, 30(12), pp.969-977.
- Tsiouri, V., Kakosimos, K.E. and Kumar, P., 2015. Concentrations, sources, and exposure risks associated with particulate matter in the Middle East Area—a review. *Air Quality, Atmosphere, and Health*, 8(1), pp.67-80.
- Turyahabwe, R., Asaba, J., Mulabbi, A. and Osuna, C., 2021. Environmental and Socio-economic Impact Assessment of Stone Quarrying in Tororo District, Eastern Uganda. *East African Journal of Environment and Natural Resources*, 4(1), pp.1-14.
- Ugbogu, O.C., Ohakwe, J. and Fotescu, V., 2009. Occurrence of respiratory and skin problems among manual stone-quarrying workers. *African Journal of Respiratory Medicine*, 2009, pp.23-26.
- Ukpong, E.C., 2012. Environmental impact of aggregate mining in the crush rock industry in Akamkpa local government area of Cross River State. *Nigerian Journal of Technology*, 31(2), pp.128-138.

- Umar, J. and Oriri, O., 2023. Environmental Effects of the Quarry Site on the Adjoining Neighborhood in Oluyole Local Government, Oyo State, Nigeria. *Ghana Journal of Geography*, 15(3), pp.223-257.
- Vallack, H.W. and Shillito, D.E., 1998. Suggested guidelines for deposited ambient dust. *Atmospheric Environment*, 32(16), pp.2737-2744.
- Vandana, M., John, S.E., Maya, K. and Padmalal, D., 2020. Environmental impact of quarrying of building stones and laterite blocks: a comparative study of two river basins in the Southern Western Ghats, India. *Environmental Earth Sciences*, 79, pp.1-15.
- Venkatesha, G., Kumar, C. and Gowda, P.R., 2015. Assessment and spatial distribution of groundwater quality in Bidadi industrial area, Ramanagara district, Karnataka, using water quality index (WQI) and GIS.
- Werner, T.T., Bebbington, A. and Gregory, G., 2019. Assessing the impacts of mining: Recent contributions from GIS and remote sensing. *The Extractive Industries and Society*, 6(3), pp.993-1012.
- William, B. and Hugh, D., 2006. Reducing the Environmental Effect of Aggregate Quarrying: Dust, Noise, and Vibration. Edited by Abbie Drew, MIRO.
- William, L., 2020. Assessment of the Effects of Quarrying Activities on the Environment: A Case Study of Lengijave Village, Arusha District. (Doctoral dissertation, The Open University of Tanzania).
- Wright, C., 2009. Air pollution monitoring and evaluation framework for South Africa: prioritizing vulnerable communities. *WIT Transactions on Biomedicine and Health*, 14, pp.55-64.
- Yakovleva, N., 2017. *Corporate Social Responsibility in The Mining Industry*. Routledge. (Edited Book)
- Yasobant, S. and Boondesh, N., 2017. Occupational and Environmental Impacts of Granite Quarry Activities in Chittoor District of Andhra Pradesh, India. India (August 8, 2017).
- Yeh, H.C. and Schum, G.M., 1980. Models of human lung airways and their application to inhaled particle deposition. *Bulletin of Mathematical Biology*, 42(3), pp.461-480.
- Zanobetti, A. and Schwartz, J., 2009. The effect of fine and coarse particulate air pollution on mortality: a national analysis. *Environmental Health Perspectives*, 117(6), pp.898-903.
- Zimwara, D., Maphosa, E., Goriwondo, W.M. and Nyadongo, S.T., 2021. Modelling the environmental impacts of noise and dust from quarry stone mining in Harare—a case study of Pomona Quarry. *Zimbabwe Journal of Science and Technology*, 16(1), pp.25-35.

ORCID DETAILS OF THE AUTHORS

V. Nagaraja: <https://orcid.org/0009-0000-2895-9166>