

REVIEW ARTICLE

<https://doi.org/10.46488/NEPT.2026.v25i04.B4443>

OPEN ACCESS

Groundwater Contamination in India: Regional Variations, Hydrogeochemical Controls, Health Risks, and Integrated Management Strategies

Santosh Sanap¹, **Shrikant Randhavane[†]** and **Narayan Chandak¹**

Department of Civil Engineering, SVKM NMIMS Global University, Dhule 424001, Maharashtra, India

[†]Corresponding author: Shrikant Randhavane; Shrikant.Randhavane@sngu.ac.in; srandhavane@gmail.com

ARTICLE HISTORY

Received: 08-04-2026

Revised: 28-05-2026

Accepted: 05-06-2026

KEYWORDS

Groundwater quality
Fluoride
Arsenic
Nitrate
Physicochemical parameters
Water quality index
Health risk

CITE THIS ARTICLE

Sanap, S., Randhavane, S. and Chandak, N., 2026 Groundwater contamination in India: Regional variations, hydrogeochemical controls, health risks, and integrated management strategies, *Nature Environment and Pollution Technology*, 25(4), B4443. <https://doi.org/10.46488/NEPT.2026.v25i04.B4443>

COPYRIGHT & LICENCE

© 2026 The Authors
Open Access under CC BY 4.0.

ABSTRACT

Groundwater plays a critical role in meeting drinking and agricultural water demands in India. This review focuses on regional variations in groundwater quality, contamination sources, and remediation strategies. The evaluation encompasses pH, electrical conductivity, total dissolved solids, hardness, major ions, and contaminants including fluoride, arsenic, nitrate, and heavy metals. Results reveal significant regional variations, with contamination driven by intensive agriculture, industrial activities, mining, urbanization, and geological factors. Water Quality Index assessments indicate that while many groundwater sources meet drinking standards, numerous sites exceed permissible limits, requiring treatment. Contamination stems from both natural processes (mineral dissolution and rock–water interactions) and anthropogenic activities (fertilizer application, industrial effluents, domestic sewage, and mining operations). Health risk assessments demonstrate elevated noncarcinogenic risks in fluoride- and nitrate-affected areas, causing fluorosis, arsenicosis, methemoglobinemia, and other waterborne diseases. Various remediation approaches suitable for Indian conditions have been investigated, including membrane-based systems, electrochemical methods, adsorption techniques, and nature-based solutions. This review provides an integrated understanding of India's regional groundwater quality challenges and emphasizes the necessity for targeted interventions, enhanced monitoring, and sustainable management strategies to protect groundwater resources. The review further provides an integrated regional comparison framework linking contamination sources, hydrogeochemical processes, health risks, and remediation strategies for groundwater management in India.

1. INTRODUCTION

1.1. Groundwater Significance in India

Water security in India is largely dependent upon groundwater, as it meets more than 85% of rural drinking water demands and nearly half of the urban water demands (Rajendran et al. 2025). With annual extraction exceeding 230 billion cubic meters, India is the largest user of groundwater worldwide (Kaur et al. 2019). The groundwater system is generally distributed across the major hydrogeological zones such as the Indo-Gangetic plains, the Deccan plateau, and the coastal zones. These zones show considerable variations in groundwater quality due to differences in geological environments, climatic conditions, and the extent of human interference (Gayathri et al. 2025, Dhakshinamoorthy & Selvarajan 2025). In the last few decades, the increasing trend of groundwater exploitation, along with the rapid pace of industrialization, intensified agricultural activities, and urbanization, has caused a deterioration in water quality (Pandey et al. 2023, Pattnaik & Bhowmick

2020). Water contamination by fluoride, arsenic, nitrate, metals, and salinity has been reported in a significant number of districts of the country, which is a major threat to public health (Kadam et al. 2021, Yenugu et al. 2020). To overcome these challenges, there is a need for a good understanding of regional patterns of groundwater quality and identification of natural as well as anthropogenic sources of contamination. This knowledge is essential for developing management, protection, and remediation strategies for sustainable use of groundwater in India (Sharma et al. 2021, Kaliyan et al. 2022). This review adopts an integrated approach by linking regional contamination patterns, groundwater chemistry, health risks, and remediation strategies to provide a comprehensive understanding of groundwater quality in India. Unlike earlier review studies that primarily focused on isolated contaminants, localized groundwater quality assessment, or individual hydrogeochemical processes, the present review provides an integrated nationwide synthesis of groundwater contamination in India by combining regional hydrogeochemical variability, contaminant distribution patterns, physicochemical characteristics, human health risks, remediation technologies, and management strategies within a single assessment framework. The review comparatively evaluates contamination trends across major hydrogeological regions of India and incorporates recently published literature up to 2025. In addition, the study emphasizes the interrelationship between contaminant sources, hydrogeochemical controls, health implications, and remediation approaches, thereby providing a broader multidisciplinary perspective useful for researchers, policymakers, and groundwater management agencies.

1.2. Methodology of Review

The present study was conducted as a structured literature review aimed at synthesizing regional groundwater contamination characteristics, hydrogeochemical controls, health risks, and remediation strategies across India. Relevant peer-reviewed literature published between 2000 and 2025 was collected from scientific databases including Scopus, Web of Science, Google Scholar, ScienceDirect, and SpringerLink using keywords such as “groundwater quality India,” “fluoride contamination,” “arsenic groundwater,” “nitrate pollution,” “water quality index,” and “health risk assessment.” Initially, approximately 300 research articles were identified through database searching. After removal of duplicate and irrelevant records, studies were screened based on title, abstract, methodological clarity, and relevance to groundwater quality assessment in India. Studies lacking quantitative groundwater quality data, clear methodological description, or regional relevance were excluded. Finally, 115 studies were selected for detailed review and synthesis.

1.2.1. Inclusion Criteria

- (i) Peer-reviewed articles;
- (ii) Studies focused on groundwater quality in India;
- (iii) Studies reporting physicochemical parameters, contaminant concentrations, health risk assessment, or remediation approaches;
- (iv) Studies containing sufficient methodological and analytical information.

1.2.2. Exclusion Criteria

- (i) Incomplete datasets;
- (ii) Conference abstracts without full text;
- (iii) Studies lacking clear sampling or analytical procedures;
- (iv) Duplicate or non-region-specific reports.

Because of substantial variability in hydrogeological conditions, sampling strategies, analytical methods, temporal scales, and reporting formats among the reviewed studies, the collected information was synthesized qualitatively rather than through formal meta-analysis. The reported concentration ranges and regional trends should therefore be interpreted as indicative comparative assessments rather than statistically aggregated national values. Unlike studies that primarily focused on isolated contaminants, localized case studies, or individual hydrogeochemical processes, the present review attempts to provide a unified nationwide synthesis of groundwater contamination in India by integrating regional hydrogeochemical variability, contaminant sources, health risk implications, remediation technologies, and management strategies within a single assessment framework. The review also incorporates recently published literature up to 2025 and comparatively evaluates contaminant occurrence across major hydrogeological regions of India. This integrated perspective is intended to support researchers, policymakers, and groundwater management agencies in developing region-specific mitigation and sustainable groundwater management strategies. The detailed literature identification, screening, eligibility assessment, and final inclusion process adopted in the present review is illustrated in Fig. 1. The review methodology adopted in the present study improves transparency, reproducibility, and comparability of synthesized groundwater quality information across different hydrogeological regions of India.

To improve methodological transparency and reproducibility, the literature screening and study selection process was performed using a PRISMA-style framework. The inclusion and exclusion criteria adopted for study selection are summarized in Table 1.

Table 1: Inclusion and exclusion criteria used for literature selection.

Criteria Type	Inclusion Criteria	Exclusion Criteria
Publication Type	Peer-reviewed journal articles	Conference abstracts without full text
Study Area	Groundwater quality studies conducted in India	Non-Indian or non-region-specific studies
Data Requirement	Quantitative groundwater quality parameters and contaminant data	Incomplete datasets
Methodology	Studies with clear sampling and analytical procedures	Studies lacking methodological clarity
topic Relevance	Hydrogeochemistry, contamination, health risk, remediation	Studies unrelated to groundwater contamination
Publication Period	2000–2025	Publications outside selected timeframe




Fig. 1: PRISMA-style flow diagram showing literature identification, screening, eligibility assessment, and final study selection process adopted in the present review.

1.3. Research Objectives and Scope

This review article attempts to collate and assess the available literature for the evaluation of groundwater quality in different regions of India. The main objectives of the study are to identify the regional variability in the physicochemical properties and concentration of major pollutants in groundwater. The study also attempts to identify the natural and anthropogenic sources of groundwater pollution, along with the health hazards that may result from the consumption of polluted groundwater. The review also attempts to identify the most commonly used water quality standards, indexing techniques, and treatment options that are applicable in the

context of India. The main advantage of the review article lies in the identification of the available literature and the valuable information that could be obtained for the better management of groundwater resources in India (Shukla & Saxena 2021, Hossain et al. 2021). Although emerging contaminants such as pharmaceuticals, PFAS, pesticides, antibiotic residues, and microplastics are increasingly recognized as potential groundwater pollutants, the present review primarily emphasizes conventional inorganic contaminants because of their widespread occurrence, stronger data availability, and greater public health significance in the Indian context.

2. REGIONAL GROUNDWATER CONTAMINATION PATTERNS

2.1. North India

The groundwater quality problems in North India are diverse, particularly in the Indo-Gangetic Plain. In Punjab and Haryana, intensive agricultural activities have led to groundwater contamination. In these areas, high nitrate concentrations, i.e., above 45 mg L^{-1} , have been recorded in 40–60% of groundwater samples, primarily due to the overuse of chemical fertilizers (Saha & Pal 2025, Raja & Neelakantan 2021). In the Malwa belt of Punjab, uranium contamination is a major issue, with levels up to $644 \text{ } \mu\text{g L}^{-1}$ compared to the WHO guideline value of $30 \text{ } \mu\text{g L}^{-1}$. The contamination is attributed to the use of phosphate fertilizers as well as geological sources (Tanwer et al. 2023). In Rajasthan, the groundwater is significantly affected by arid and semi-arid climatic conditions. High fluoride levels ranging from 1.5 to 8.5 mg L^{-1} and high salinity are commonly reported. These conditions are mainly attributed to high rates of evaporation and dissolution of fluoride minerals and low rates of natural recharge (Kaur & Godara 2025, Maurya et al. 2025, Dhakate et al. 2023). In the case of Delhi and its surrounding National Capital Region, groundwater is characterized by elevated concentrations of total dissolved solids, hardness, and nitrates, with concentrations of 500 to 2500 mg L^{-1} , 300 to 800 mg L^{-1} , and 50 to 150 mg L^{-1} , respectively. These issues are mainly related to sewage water leakage, industrial waste, and agricultural runoff (Maurya et al. 2021, Singh et al. 2024). In Uttar Pradesh, there is a high degree of spatial variability in groundwater quality. In eastern parts of the state, arsenic contamination of groundwater, with concentrations varying between 10 and $500 \text{ } \mu\text{g L}^{-1}$, is common, whereas fluoride contamination, with concentrations varying from 1.5 to 6.0 mg L^{-1} , is common in the western parts of Uttar Pradesh (Adimalla 2020, Karunanidhi et al. 2021). In Jammu and Kashmir, groundwater quality is relatively good, although localized pollution, mainly related to agricultural activities and poor sanitation, has been reported in certain areas of the valleys (Chakraborty et al. 2021). The variability in groundwater contamination in North India is mainly regulated by the interaction of agricultural practices, hydrogeology, and urbanization. The nitrate concentration in the groundwater of Punjab and Haryana is strongly influenced by the excessive application of fertilizers (Saha & Pal 2025, Raja & Neelakantan 2021, Marghade et al. 2021). The presence of uranium in the Malwa region is regulated by both geochemical and fertilizer components (Tanwer et al. 2023). Arsenic in the groundwater of the eastern part of Uttar Pradesh is regulated by reductive dissolution in the

alluvial aquifers (Adimalla 2020, Karunanidhi et al. 2021). It is concluded that groundwater quality is location-specific and is regulated by both geochemical and anthropogenic factors.

2.2. South India

In South India, groundwater is mostly linked with hard rock aquifer systems that are recognized to exhibit specific hydrogeochemical signatures. In Tamil Nadu State, fluoride contamination is widespread, with fluoride levels ranging from 1.5 to 12 mg L^{-1} covering 15 districts. The maximum fluoride levels are reported from Salem, Dharmapuri, and Krishnagiri districts due to the weathering of fluoride-bearing granites and gneisses (Ramalingam et al. 2022, Ramajayam et al. 2024). Seawater intrusion is another major problem reported from the coastal areas of Tamil Nadu State, where the chloride level is above 1000 mg L^{-1} and electrical conductivity is above $3000 \text{ } \mu\text{S cm}^{-1}$, extending up to 5–10 km inland (Ramajayam et al. 2024, Singh et al. 2024). The groundwater quality in Karnataka state has been found to vary in different agro-climatic zones. In the northern districts of the state, the groundwater exhibits elevated concentrations of fluoride (2 – 8 mg L^{-1}) and nitrate (50 – 200 mg L^{-1}), whereas salinity intrusion is a major problem in the coastal districts (Sarma & Singh 2023, Taloor et al. 2023). In the Bangalore urban region, the groundwater is contaminated by elevated concentrations of nitrate (45 – 120 mg L^{-1}), the presence of heavy metals such as iron, manganese, lead, and chromium, and bacterial contamination. The causes of groundwater contamination in this region include industrial activities and the lack of proper sewage treatment systems (Panneerselvam et al. 2021, Biswas et al. 2023). Andhra Pradesh and Telangana states have reported widespread fluoride contamination in various districts, ranging from 1.5 to 9.5 mg L^{-1} . The problem is more acute in districts like Nalgonda, Prakasam, and Guntur (Goyal et al. 2025, Karanveer et al. 2022). In Kerala, coastal groundwater is highly affected by seawater intrusion, with chloride levels above 1500 mg L^{-1} . Approximately 30–40% of groundwater wells located within 2 km of the coastline have elevated salinity and chloride concentrations due to seawater intrusion (CGWB 2017). Inland regions of the state have shown better groundwater quality, although some instances of water pollution by agricultural and domestic waste disposal have been reported (Sane et al. 2025). Puducherry has reported high salinity levels, ranging from 1000 to 3500 mg L^{-1} in terms of TDS, along with nitrate levels ranging from 50 to 180 mg L^{-1} due to the high intensity of agricultural activities in the region (Subramaniyan & Ganesan 2024). The regional variation in groundwater quality in South India is controlled by hard rock aquifer systems, climatic conditions, and coastal processes. For instance, elevated concentrations

of fluoride in groundwater are linked to increased water-rock interaction in granitic rocks (Ramalingam et al. 2022, Singh et al. 2020, Adimalla et al. 2020), whereas salinity in coastal groundwater is controlled by excessive groundwater abstraction and hydraulic imbalance (Singh et al. 2024, Sarma & Singh 2023). In cities such as Bangalore, elevated nitrate and heavy metal concentrations are observed due to poor wastewater management (Panneerselvam et al. 2021, Biswas et al. 2023). These are examples of the effects of geology, urbanization, and coastal processes on groundwater quality.

2.3. East India

Eastern India has a very serious groundwater quality problem, which is mainly in the form of arsenic contamination. This problem is particularly acute in the West Bengal region, where arsenic has been found in 79 blocks in nine districts. The concentration of arsenic has been found to vary from 10 to as high as $3200 \mu\text{g L}^{-1}$, which is well above the WHO guideline value of $10 \mu\text{g L}^{-1}$ (Mukherjee et al. 2015, Sabinaya et al. 2023, Khan et al. 2021). The source of arsenic in the groundwater in the eastern part of India is natural. Geological processes are responsible for the occurrence of arsenic in the groundwater in the eastern part of India. The reducing conditions in the alluvial aquifer allow the release of arsenic from iron oxyhydroxide into the groundwater (Malik et al. 2021). A similar trend has been observed in the Bihar region, where arsenic contamination has been reported in about 15 districts, with a maximum concentration of $1800 \mu\text{g L}^{-1}$ (Sahu et al. 2017). In Jharkhand, the groundwater quality is highly affected by mining activities. In areas where coal mining, iron ore mining, and industries are prevalent, high concentrations of heavy metals such as iron, manganese, lead, cadmium, chromium, copper, and zinc have been recorded (Panneerselvam et al. 2023, Reddy et al. 2022). In addition, fluoride contamination has been recorded in the western districts, where the concentration of fluoride varies from 1.5 to 6.5 mg L^{-1} , primarily due to geological factors (Singh et al. 2022). In Odisha, a significant variation in groundwater quality has been recorded. In the coastal areas, salinity intrusion is a major problem, whereas in the western districts, fluoride concentrations vary from 2 to 7 mg L^{-1} , and areas near mining and industrial activities have high concentrations of heavy metals (Kouser et al. 2022). The quality of groundwater in Assam and other states in the northeastern region tends to be better, considering the high rainfall and good recharge conditions. Yet, there are some reported cases of contamination in the tea estate areas, oil refineries, and in places with poor sanitation facilities (Basha et al. 2022). At some places, the iron and manganese content in the groundwater also exceeds the permissible limits of

0.3 mg L^{-1} and 0.1 mg L^{-1} , respectively, mainly because of the reducing conditions in the aquifers. (Singh et al. 2022) The major processes controlling groundwater contamination in eastern India are geogenic in nature, dominated by arsenic contamination in reducing environments in alluvial aquifers (Mukherjee et al. 2015, Sabinaya et al. 2023, Khan et al. 2021). In contrast to western and southern India, where groundwater contamination is dominated by fluoride, eastern India has its own set of groundwater contamination processes, dominated by sedimentary geology and redox chemistry (Malik et al. 2021, Sahu et al. 2017). In addition, there is heavy metal contamination in Jharkhand due to mining activities, which clearly indicates anthropogenic influences on groundwater chemistry (Panneerselvam et al. 2023, Reddy et al. 2022).

2.4. West India

Significant variations in the quality of groundwater in Western India have been observed. In Gujarat, various problems in the quality of groundwater have been reported. Fluoride contamination in the range of 1.5 to 10 mg L^{-1} has been reported in ten districts. The coastal areas in the Saurashtra and Kutch districts are also affected by salinity intrusion, where the TDS level ranges from 1500 to 5000 mg L^{-1} . Moreover, areas where agricultural activities are concentrated also face problems of nitrate contamination in the range of 50 to 300 mg L^{-1} (Yadav et al. 2018, Dhanya Raj & Shaji 2017, Shaikh et al. 2020). Groundwater contamination by heavy metals like chromium, lead, cadmium, and nickel has been reported in the industrial areas around Ahmedabad, Vadodara, and Surat cities, which is attributed to the effluent discharge from the textile, chemical, and pharmaceutical industries (Srinivas et al. 2013). Groundwater quality in Maharashtra exhibits significant spatial variability influenced by geological and anthropogenic factors. In Vidarbha and Marathwada, the presence of fluoride ranges between 2–8 mg L^{-1} . Nitrate-related problems are present in areas with sugarcane cultivation, ranging between 50–250 mg L^{-1} . (Mhaske et al. 2022, Naskar et al. 2025). Regions with industries have higher levels of heavy metals in groundwater. In Mumbai and other major cities, the issue of total dissolved solids, hardness, and bacterial presence are major problems, and these are mainly caused by leakage in the sewage system and industries (Wagh et al. 2019). In the state of Goa, the presence of seawater in the groundwater of the coastal areas ranges between 40–50% with chloride levels above 2000 mg L^{-1} within 3 km of the shoreline (Sidhu et al. 2024). In the western part of India, the groundwater quality is affected by the arid climatic conditions, the high intensity of agriculture, and industrialization. Elevated fluoride and salinity concentrations in groundwater samples are related

to evaporation and the dissolution of minerals (Yadav et al. 2018, Dhanya Raj & Shaji 2017, Shaikh et al. 2020). Elevated concentrations of nitrate in groundwater samples reflect the impact of agricultural activities (Shaikh et al. 2020). Industrialization has affected the groundwater quality in the form of heavy metals in the urban areas (Srinivas et al. 2013). The impact in the western part of India is a mix of geogenic and anthropogenic factors compared to the eastern part.

2.5. Central India

Groundwater in Central India reflects the fingerprints of nature as well as human activities. In Madhya Pradesh, the presence of fluoride poses a problem in several areas, ranging from 1.5 to 7.5 mg L⁻¹ over 14 districts of the state. The districts of the Malwa Plateau and Vindhyan areas are of particular concern, and the cause of the problem is the weathering of those minerals containing fluoride (Karunanidhi et al. 2021, Vijayakumar et al. 2022). Nitrate contamination of groundwater occurs in several agricultural districts, where the excessive use of chemical fertilizers is the major cause of the problem, with nitrate concentrations ranging from 50 to 180 mg L⁻¹ (Alam et al. 2024). In the state of Chhattisgarh, the groundwater scenario reflects the impact of mining and industries. In the state, heavy metals

like iron, manganese, chromium, lead, and cadmium are present in groundwater at elevated concentrations, often two to ten-fold higher than the permissible limits (Ullengula et al. 2026). In the coal mining areas, sulfate concentrations are significantly elevated, ranging from 200 to 600 mg L⁻¹, and TDS concentrations are also elevated, ranging from 1000 to 3000 mg L⁻¹, mainly because of acid mine drainage (Karunanidhi et al. 2020). In the agricultural areas, nitrate contamination of groundwater occurs, with concentrations ranging from 45 to 150 mg L⁻¹, mainly because of the application of fertilizers and the disposal of organic waste (Ambade et al. 2024). Fig. 2 shows the regional distribution of major groundwater contaminants in India. The quality of groundwater in central India is characterized by various geological formations and anthropogenic activities like mining and agricultural activities. Higher concentrations of fluoride ions are related to the weathering of minerals, whereas heavy metal contamination is mainly related to mining activities. Compared to other regions, central India is characterized by mixed contaminant patterns due to various natural and anthropogenic factors. The map highlights dominant groundwater contamination zones reported in reviewed studies, including arsenic contamination in eastern alluvial aquifers, fluoride enrichment in western and southern alluvial aquifers, fluoride enrichment in western and southern




Fig. 2: Regional distribution of major groundwater contaminants across India showing dominant fluoride contamination in western and southern states, arsenic occurrence in eastern alluvial plains, and nitrate enrichment in agriculturally intensive regions.

hard-rock terrains, nitrate contamination in agriculturally intensive northern regions, salinity intrusion in coastal aquifers, and heavy metal contamination near mining and industrial zones.

2.6. Comparative Regional Analysis

A regional perspective of groundwater contamination in India shows varying degrees of contamination. In the northern part of India, the major groundwater contamination problems are related to nitrates and uranium. These are primarily related to agricultural activities and fertilizer usage. In the western and southern parts of India, fluoride and salinity are major

groundwater contamination problems. These are related to geological and climatic factors. In the eastern part of India, arsenic contamination is a major problem, and this is primarily related to geological factors. In central India, heavy metals are a major groundwater contamination problem, and this is primarily related to mining activities (Saraswat et al. 2023, Khan et al. 2021). A comparative evaluation of groundwater quality in various regions of India has shown specific patterns of groundwater quality in different regions. Northern and western regions exhibit contamination patterns dominated by nitrate and fluoride, respectively, due to agricultural activities and climatic conditions (Yadav et al. 2018, Dhanya

Table 2: Regional groundwater quality parameters across major Indian locations.

State	pH	EC ($\mu\text{S cm}^{-1}$)	TDS (mg L^{-1})	TH (mg L^{-1})	F ⁻ (mg L^{-1})	NO ₃ ⁻ (mg L^{-1})	As ($\mu\text{g L}^{-1}$)	Primary Concerns	Authors
Punjab	7.2–8.5	800–3500	500–2200	300–900	0.5–2.5	45–250	5–644	Nitrate, Uranium	Saha & Pal 2025, Raja & Neelakantan 2021, Marghade et al. 2021
Rajasthan	7.0–8.8	1200–5500	750–3500	400–1500	1.5–8.5	20–120	<10	Fluoride, Salinity	Kaur & Godara 2025, Maurya et al. 2025, Dhakate et al. 2023
Delhi NCR	7.1–8.3	900–4000	600–2500	300–800	0.3–2.0	50–150	<10	TDS, Nitrate, Hardness	Maurya et al. 2021, Singh et al. 2024
Tamil Nadu	6.8–8.6	800–4500	500–3000	250–1200	1.5–12.0	30–180	<10	Fluoride, Salinity	Ramalingam et al. 2022, Singh et al. 2020, Adimalla et al. 2020
Karnataka	6.5–8.4	700–3800	450–2400	200–950	2.0–8.0	50–200	<10	Fluoride, Nitrate	Sarma & Singh 2023, Taloor et al. 2023
West Bengal	6.8–7.8	600–2500	400–1600	200–700	0.2–1.5	20–80	10–3200	Arsenic, Iron	Mukherjee et al. 2015, Sabinaya et al. 2023, Khan et al. 2021
Bihar	6.9–7.9	650–2800	420–1800	220–750	0.3–2.0	25–90	10–1800	Arsenic, Fluoride	Sahu et al. 2017
Gujarat	7.0–8.7	1000–6000	650–4000	300–1800	1.5–10.0	50–300	<10	Fluoride, Salinity, Nitrate	Yadav et al. 2018, Dhanya Raj & Shaji 2017, Shaikh et al. 2020
Maharashtra	6.8–8.5	750–4200	480–2700	250–1100	2.0–8.0	50–250	<10	Fluoride, Nitrate	Mhaske et al. 2022, Naskar et al. 2025
Madhya Pradesh	7.0–8.6	800–3600	520–2300	280–950	1.5–7.5	50–180	<10	Fluoride, Nitrate	Karunanidhi et al. 2021, Vijayakumar et al. 2021, Alam et al. 2024
Chhattisgarh	6.5–8.2	700–3500	450–2200	240–900	0.5–3.0	45–150	<10	Heavy Metals, Nitrate	Ullengula et al. 2026, Karunanidhi et al. 2020
Odisha	6.7–8.3	650–3800	420–2400	220–950	2.0–7.0	30–120	<10	Fluoride, Heavy Metals	G. et al. 2024 Kouser et al. 2022

Note: The values in Table 2 are compiled from multiple studies corresponding to the cited references for each region to provide a literature-supported regional comparison across India.

Raj & Shaji 2017, Shaikh et al. 2020). The eastern parts of India are affected by arsenic contamination (Malik et al. 2021), whereas in southern India, both fluoride and salinity are major groundwater quality problems (Ramalingam et al. 2022). In central India, mining activities are responsible for heavy metal contamination (Ullengula et al. 2026). The results show that groundwater contamination is region-specific. The regional variations in groundwater quality discussed above can be seen in terms of physicochemical properties of groundwater, which are discussed in detail in the next section. The variations discussed here show how geology and human activities affect groundwater chemistry in various parts of India.

The concentration ranges presented in Table 2 represent compiled minimum–maximum ranges reported in the reviewed literature for different groundwater quality parameters across selected regions of India. The values are synthesized from multiple independent studies conducted under varying hydrogeological, seasonal, and analytical conditions and should therefore be interpreted as indicative comparative ranges rather than standardized regional averages.

The contaminant concentration ranges summarized in Table 3 represent indicative ranges compiled from published regional groundwater studies. The values reflect reported concentration intervals from different hydrogeological settings and are intended for comparative regional assessment rather than direct statistical comparison.

The comparative analysis suggests that the arsenic contamination in eastern India poses the highest health risk in terms of toxicity and prevalence, while the fluoride contamination in the western and southern regions has a larger population affected. Nitrate contamination in India has moderate to high spatial variability and is associated with agricultural regions. These comparisons provide a semi-quantitative evaluation of the severity of groundwater contamination in India.

3. HYDROGEOCHEMICAL CHARACTERISTICS AND CONTAMINATION MECHANISMS

3.1. pH

Groundwater pH in India ranges from a neutral to mildly alkaline level, usually between 6.5 and 8.8 (Krishnappa et al. 2024). pH values above 8.0 are commonly reported in arid and semi-arid zones, where evaporation promotes the dissolution of carbonate minerals. If the rainfall is significantly elevated or in areas where mines are present, then the pH level may turn slightly acidic (Shirke et al. 2020). The level of pH is a very significant factor, as it influences the movement of metals, the interaction of minerals, and the treatment process (Saha et al. 2024).

3.2. Electrical Conductivity (EC) and Total Dissolved Solids (TDS)

The range for groundwater conductivity is normally between 600 and 6000 $\mu\text{S cm}^{-1}$, which is equivalent to a TDS of about 400 to 4000 mg L^{-1} (Hanse et al. 2019). Increased levels of electrical conductivity and TDS imply an increased amount of dissolved ions, often resulting from mineral weathering, evaporation, and human activities such as farming and industrial operations (Mukherjee & Singh 2022). Research carried out in various parts of India has indicated that about 40% of the groundwater has TDS levels above the permissible limit of 500 mg L^{-1} (Sahoo et al. 2024, Mukherjee & Singh 2022).

3.3. Total Hardness (TH)

The range of total groundwater hardness is 200 to 1800 mg L^{-1} as CaCO_3 . Reviewed regional studies have reported that approximately 40–60% of groundwater samples in several investigated regions of India fall within the hard to very hard category, with values above 300 mg L^{-1} (Ravindra & Mor 2019). The high level of groundwater

Table 3: Comparative regional summary of groundwater contamination in India.

Region	Dominant Contaminants	Typical Concentration Range	Key Causes	Severity Level	Authors
North India	Nitrate, Uranium	NO_3^- 50–150 mg L^{-1}	Fertilizers, agriculture	High	Saha & Pal 2025, Chakraborty et al. 2021
South India	Fluoride, Salinity	F^- 2–12 mg L^{-1}	rock–water interaction, coastal intrusion	High	Ramalingam et al. 2022, Subramaniyan & Ganesan 2024
East India	Arsenic	As 10–3200 $\mu\text{g L}^{-1}$	Reductive dissolution (geogenic)	Very High	Mukherjee et al. 2015, Singh et al. 2022
West India	Fluoride, Nitrate, Salinity	F^- 2–10 mg L^{-1}	Arid climate, agriculture	High	Yadav et al. 2018, Sidhu et al. 2024
Central India	Fluoride, Heavy Metals	F^- 1.5–7 mg L^{-1}	Mining, geology	Moderate–High	Karunanidhi et al. 2021, Ambade et al. 2024

Note: Table 3 summarizes dominant contaminants, their typical ranges, and associated causes across different regions.

hardness is attributed to the elevated concentrations of calcium and magnesium ions that are dissolved from rocks such as limestone, dolomite, and gypsum (Adimalla et al. 2018). Although the level of water hardness is not a health hazard, high levels can cause pipe scaling and could possibly contribute to heart problems (Ahada & Suthar 2019). Dissimilarities in methodologies for analysis applied in different studies, including differing limits of detection and types of instrumentation (e.g., ICP-MS vs. AAS), as well as sample preservatives applied, may also contribute to differing results. These methodological dissimilarities may add to the uncertainty in results (Hanse et al. 2019, Mukherjee & Singh 2022).

3.4. Major Cations and Anions

3.4.1. Cations

Calcium and magnesium are the major cations present in Indian groundwater. The concentration of calcium ranges from 40–450 mg L⁻¹, while magnesium ranges from 20–280 mg L⁻¹. The major source of calcium and magnesium is mineral weathering, i.e., carbonate and silicate mineral weathering (Mawari et al. 2022). The concentration of sodium ranges from 20–800 mg L⁻¹, while potassium ranges from 2–50 mg L⁻¹. However, in arid areas, coastal areas, and areas of heavy agricultural activity, higher concentrations of sodium and potassium have also been reported, primarily due to evaporation, intrusion of seawater, and extensive use of chemical fertilizers (Mukherjee et al. 2019). Higher concentrations of sodium also pose a major problem, as they reduce soil permeability and can be harmful to human health, especially for people suffering from hypertension (Yadav et al. 2019).

3.4.2. Anions

The most common negatively charged ions present in groundwater are bicarbonate ions. The range of bicarbonate ions is between 150 and 800 mg L⁻¹. Bicarbonate ions are generally derived from the dissolution of carbonate minerals and the action of carbon dioxide (Ahada & Suthar 2018). The level of chloride ions is generally between 20 and 2000 mg L⁻¹. The level of chloride ions is high in coastal areas where seawater intrusion occurs, dry areas where salts are present due to evaporation, and areas where sewage leakage is observed (Hussain et al. 2012). The level of sulfate ions is generally between 20 and 600 mg L⁻¹. The level of sulfate ions is derived from the dissolution of gypsum minerals and industrial waste, as well as acid mine drainage (Madhavan et al. 2023). Around 35–45% of groundwater samples from highly intensive agricultural areas such as Punjab, Haryana, and Gujarat show nitrate levels above the permissible limit of 45 mg L⁻¹. Conversely, in less intensive areas, the levels

of exceedance are low. This may be due to the variation in the application of fertilizers (Chaudhry & Sachdeva 2021).

3.5. Heavy Metals and Trace Elements

3.5.1. Fluoride

There are reports of the presence of fluoride in groundwater in many parts of India, with concentrations ranging from 1.5 to 12 mg L⁻¹. This problem occurs in several districts across several states, affecting millions of people (Das et al. 2009). The primary sources of fluoride in groundwater are the weathering of minerals like fluorite, apatite, and mica, all of which are present in the environment in granitic and gneissic rocks (Ghosh et al. 2024). Groundwater chemistry, like pH, calcium, and bicarbonate, also contributes to the presence of fluoride in groundwater (Gupta et al. 2019).

3.5.2. Arsenic

Among the major groundwater pollutants, arsenic is particularly notable in India, and its concentration in water varies sharply, from a minimum of 10 µg L⁻¹ to a maximum of 3200 µg L⁻¹, with particularly elevated concentrations of contamination detected in states such as West Bengal, Bihar, Uttar Pradesh, Assam, and Chhattisgarh (Gupta et al. 2015). The major source of arsenic entering groundwater is the reductive dissolution of iron oxyhydroxides, particularly in anaerobic and alluvial aquifer systems (Cherukumilli et al. 2018). In groundwater, arsenic is observed in two forms, arsenite (As³⁺) and arsenate (As⁵⁺). Arsenite is more mobile and toxic than arsenate (Inaniyan & Raychoudhury 2019).

3.5.3. Other Heavy Metals

Iron and manganese are often present in groundwater samples in concentrations higher than the safety limits, especially in areas where the environment of the aquifer is reducing. In these areas, iron concentrations vary between 0.1 and 15 mg L⁻¹, and manganese concentrations vary between 0.05 and 5 mg L⁻¹. These two elements are often present together and can cause problems with the color and taste of the water, as well as health risks (Koley 2021). Other heavy metals, like lead (5–150 µg L⁻¹), cadmium (2–50 µg L⁻¹), chromium (10–500 µg L⁻¹), copper (10–200 µg L⁻¹), and zinc (50–1000 µg L⁻¹), are usually reported in areas near industrial centers, mining areas, and areas with intensive agricultural activities, as these are areas with the strongest anthropogenic influence (Singh et al. 2024, Awasthi et al. 2021). Drinking water containing these heavy metals can cause health problems like neurological disorders, kidney problems, and cancer (Khandaker & Rahman 2022). Fig. 3 presents the integrated groundwater quality assessment methodology and its significance.




Fig. 3: Integrated methodological workflow adopted for groundwater quality assessment, hydrogeochemical interpretation, health-risk evaluation, and remediation strategy selection in Indian groundwater studies.

The assessment workflow presented in Fig. 3 demonstrates the importance of integrating field sampling, laboratory analysis, water quality indexing, hydrogeochemical interpretation, and health risk assessment for comprehensive groundwater quality evaluation. The figure indicates that groundwater assessment should not rely solely on physicochemical parameter measurement, but should also include interpretation of contaminant sources, exposure pathways, and associated human health implications. This integrated assessment approach improves the reliability of groundwater quality evaluation and supports region-specific groundwater management strategies.

3.5.4. Nutrients and Emerging Contaminants

One of the most common nutrient pollutants in India's groundwater is nitrate. The primary sources of nitrate contamination are fertilizers, animal waste, and sewage. Nitrate contamination of groundwater is a major issue, as it has been reported that almost 40% of groundwater in highly cultivated areas exceeds permissible limits, although this is not so prevalent in areas with less agricultural activity (Marghade et al. 2023). In India, the normal concentration of phosphates in groundwater is within a range of 0.1 to

5 mg L⁻¹. An increase in phosphates is a sign of organic pollution, which may result in eutrophication in water bodies that are connected to the groundwater (Wintgens et al. 2015). In recent times, there has been a surge of interest in newer pollutants, including pesticides, pharmaceuticals, and microplastics, in India's groundwater. However, there is limited information on the sources, occurrence, and potential impact of these pollutants on India's groundwater (Pal et al. 2009, Sultan et al. 2024). In recent years, increasing concern has also been raised regarding emerging contaminants such as pesticide residues, pharmaceutical compounds, personal care products, antibiotic residues, PFAS (per- and polyfluoroalkyl substances), endocrine-disrupting compounds, and microplastics in groundwater systems. These contaminants primarily originate from agricultural runoff, untreated sewage discharge, landfill leachates, industrial effluents, hospital wastewater, and improper disposal of domestic and pharmaceutical waste. Although systematic nationwide data for these contaminants in Indian groundwater remain limited, several localized investigations have reported detectable concentrations in urban, peri-urban, and agriculturally intensive regions. Long-term exposure to such contaminants may pose ecological risks, antimicrobial

resistance concerns, endocrine disruption, and chronic human health impacts. However, due to limited monitoring data, inconsistent analytical methods, and lack of standardized national datasets, the present review mainly focuses on conventional inorganic groundwater contaminants such as fluoride, arsenic, nitrate, salinity, and heavy metals, which remain the most widely reported and critical groundwater quality concerns across India.

3.6. Contamination Sources

The diagram identifies the various sources of contamination that affect the groundwater in India. The sources include runoff from agriculture, which contains fertilizers and pesticides; industrial waste, which contains heavy metals and chemical compounds; domestic sewage, which contains nutrients; mining, which results in acid and the release of metals; and geogenic sources, which contain fluoride and arsenic compounds that are released from the interaction between rocks and water. The pathways that are represented in the diagram include infiltration, leaching, and lateral movement from surface water bodies. It is important to note that the variations in the reported values of groundwater quality from various sources may be attributed to various factors. For example, some sources may have been based on single-time sampling, while others may have been based on seasonal sampling. This may have caused inconsistencies

in the reported values. It is significant to note that the groundwater quality data collected and compiled in this review have been obtained from various studies conducted over different time periods, hydrogeological settings, and methodological approaches. These factors can have a major impact on the data. For example, groundwater samples collected before the monsoon season have shown high concentrations of contaminants (Krishnappa et al. 2024, Shirke et al. 2020). whereas samples collected after the monsoon season have shown low concentrations (Sahoo et al. 2024). Similarly, alluvial aquifers have shown different geochemical characteristics compared to hard rock aquifers (Mawari et al. 2022). Therefore, direct comparison of values across studies should be interpreted cautiously. The variation in physicochemical parameters and contaminant concentrations is related to both natural and anthropogenic sources. This is significant in understanding the patterns of groundwater chemistry discussed above. Fig. 4 demonstrates that groundwater contamination pathways in India are highly interconnected. Agricultural runoff contributes primarily to nitrate and pesticide contamination, while industrial effluents are major sources of heavy metals and toxic chemicals. Geogenic contamination pathways differ significantly from anthropogenic pathways because they are strongly controlled by aquifer mineralogy and groundwater chemistry. The figure further highlights that contaminant migration




Fig. 4: Major geogenic and anthropogenic groundwater contamination sources and contaminant transport pathways in Indian aquifer systems (synthesized from the literature).

occurs through infiltration, leaching, and groundwater flow processes, emphasizing the importance of hydrogeological conditions in contaminant transport.

3.7. Natural Geogenic Sources

Water can also be contaminated by natural geological and hydrogeochemical processes. Fluoride concentration increases due to the weathering of fluoride-bearing minerals like fluorite, apatite, biotite, and hornblende, which are present in rock formations of granite, gneiss, and volcanic origins (Malik et al. 2023). Also, groundwater with an alkaline nature (pH above 7.5), low calcium content, and high bicarbonate and sodium levels is more likely to dissolve more fluoride and move it around (Barathkumar et al. 2025). As for the issue of arsenic contamination, it is mostly associated with alluvial aquifers where iron and manganese oxyhydroxide dissolution occurs reductively under anaerobic conditions (Cherukumilli et al. 2017). Microbial processes also enhance the degradation rate of organic matter and alter geochemical conditions, moving towards more reducing conditions, which in turn increases the release of arsenic in groundwater (Kalaiselvi & Pillai 2019). Under arid and semi-arid conditions, salinity problems are mainly driven by the rapid rate of evaporation, the dissolution of salt minerals such as halite and gypsum, and the lack of recharge to groundwater (Vijaylaxmi & Khan 2021). For example, in coastal aquifers, seawater intrusion occurs due to the pumping of groundwater, which changes the hydraulic gradients and causes seawater to flow inland (Saha et al. 2014). Heavy metals such as iron, manganese, uranium, and thorium are found in specific geological formations and are transported through groundwater depending on parameters such as pH and redox reactions (Karunanidhi et al. 2024).

3.8. Anthropogenic Sources

3.8.1. Agricultural Activities

Agricultural practices in most areas of India also contribute significantly to the contamination of groundwater. This is particularly so because of the extensive use of chemical fertilizers, which increases the nitrate and phosphate content in the water (Kumar et al. 2017). The use of pesticides, coupled with irrigation return flows, further contributes to the degradation of water quality. Moreover, in areas such as Punjab, Haryana, and Gujarat, the nitrate content in the water is above 100 mg L⁻¹, which is attributed to the extensive use of nitrogen fertilizers in these areas (Mandal & Sanyal 2019). Livestock and poultry farming also contributes to the contamination of the water table (Narsimha & Sudarshan 2017).

3.8.2. Industrial Effluents

Industrial activities such as textile finishing, chemical and pharmaceutical production, leather tanning, electroplating, and mining also pose a major threat to groundwater. These activities may release heavy metals such as chromium, lead, cadmium, nickel, copper, and zinc, along with organic compounds and other toxic chemicals (Muthusamy et al. 2023). When effluent treatment is inadequate and waste is disposed of in unlined areas, chemicals can readily leach into groundwater (Nagaraj & Masilamani 2023). Consequently, high concentrations of heavy metals are found in industrial areas of Gujarat, Maharashtra, Tamil Nadu, and West Bengal (Bhuiyan & Ray 2017).

3.8.3. Domestic Sewage and Solid Waste

In addition, poor sewage disposal and unreliable sanitation facilities in urban areas also cause groundwater contamination. When untreated sewage enters the ground, it carries nitrates, phosphates, chlorides, and various organic compounds with it, thereby contaminating groundwater (Yasaswini et al. 2024). In rural areas, open defecation also causes environmental pollution, leading to contamination of water bodies (Mukherjee et al. 2021). Moreover, when solid wastes are disposed of in open dumps, leachate may carry heavy metals, organic compounds, and dissolved solids, thereby contaminating groundwater (Thalod & Sharma 2024).

3.8.4. Mining Operations

Mining activities such as coal, iron, and bauxite mining cause acid mine drainage, suspended solids, and elevated levels of heavy metals (Panneerselvam et al. 2023). Mining-related groundwater pollution is significant in Jharkhand, Chhattisgarh, and Odisha (Reddy et al. 2022).

3.9. Contamination Pathways and Mechanisms

Groundwater contamination occurs in various forms. Groundwater may be contaminated by seepage from surface sources, downward percolation from the unsaturated zone, and horizontal movement from already contaminated surface sources such as rivers, lakes, and canals. Groundwater may also be contaminated vertically or horizontally along fractures and joints, especially in hard rocks (Cherukumilli et al. 2017). Contaminated groundwater does not move in an arbitrary manner; it is influenced by various elements. The behavior and movement of the contaminant depend on the characteristics of the aquifer, i.e., the ease with which both water and contaminants pass through it (permeability), the available void spaces (porosity), and the minerals that react with the contaminant. Other elements include the hydrogeological characteristics, i.e., the rate at which the water table replenishes, the rate at which the groundwater

moves, and the depth at which the water table lies. Finally, there are the characteristics of the contaminant, i.e., its solubility how easily it moves, and the period it lasts in the environment without degrading. All these elements contribute to the behavior and movement of the contaminant in the aquifer system (Kalaiselvi & Pillai 2019)

4. HEALTH IMPLICATIONS

Fig. 5 depicts the pathway from contamination sources through exposure routes (drinking water, food chain, dermal contact) to health outcomes. It shows acute effects (gastrointestinal illness, methemoglobinemia) and chronic effects (fluorosis, arsenicosis, cancer, neurological disorders, kidney disease) with vulnerable populations (children, pregnant women, elderly) highlighted. The contamination

sources and pathways discussed above directly affect human exposure to polluted groundwater. The presence of pollutants such as fluoride, arsenic, nitrate, and heavy metals in groundwater poses serious health risks, which are discussed in the section below. Fig. 5 illustrates the relationship between groundwater contamination exposure pathways and associated human health outcomes. The figure indicates that drinking water remains the dominant exposure pathway for most groundwater contaminants in India. Long-term exposure to fluoride, arsenic, nitrate, and heavy metals contributes to chronic health effects including fluorosis, arsenicosis, neurological disorders, kidney damage, and carcinogenic risks. The figure also highlights that children and elderly populations are more vulnerable because of higher exposure sensitivity and physiological susceptibility.




Fig. 5: Major groundwater contamination exposure pathways and associated human health impacts in India.

4.1. Health Effects of Major Contaminants

4.1.1. Fluoride

When the amount of fluoride exceeds 1.5 mg L^{-1} , there is a high probability of causing fluorosis in both teeth and bones. In the case of teeth, fluorosis causes mottling and discoloration of the teeth, while in bones, it causes changes in the structure of the bones, joint pain, and lack of mobility (Das et al. 2009). In India, there is a high prevalence of fluorosis, and Previous national and regional assessments have estimated that nearly 66 million people in India may be exposed to fluoride-affected groundwater (Ghosh et al. 2024). In addition, it causes neurological effects, impacts IQ in children, and causes reproductive problems (Gupta et al. 2019).

4.1.2. Arsenic

Prolonged consumption of drinking water contaminated with arsenic causes a condition referred to as arsenicosis, which manifests itself in the form of changes in the color of the skin, keratosis, and other skin lesions (Gupta et al. 2015). Prolonged consumption also increases the risk of cardiovascular problems, neurological problems, diabetes, and various types of cancer, which affect the skin, lungs, bladder, and kidneys. A considerable number of people in eastern India continue to be at risk of arsenic poisoning through the drinking water supplied to them through the taps (Cherukumilli et al. 2018). Among the various compounds of arsenic, arsenite (As^{3+}) is found to be more poisonous and mobile than arsenate (As^{5+}) (Inaniyan & Raychoudhury 2019).

4.1.3. Nitrate

Drinking water with nitrate concentrations above 45 mg L^{-1} are particularly hazardous for infants, as it can cause methemoglobinemia, or 'blue baby syndrome,' which impairs the blood's capacity to carry oxygen. Adults who are frequently exposed to nitrates have a higher probability of developing gastric, thyroid, and reproductive system diseases (Kumar et al. 2017). In areas with heavy agricultural activity, there is a higher incidence of diseases associated with nitrate poisoning (Mandal & Sanyal, 2019).

4.1.4. Heavy Metals

Heavy metals in groundwater pose health risks. Lead affects the nervous system, resulting in reduced IQ, behavioral problems, and kidney damage (Singh et al. 2024). Cadmium causes a buildup of the metal in the kidneys, resulting in renal failure and bone disease (Awasthi et al. 2021). Hexavalent chromium is a known carcinogen and causes lung and digestive cancers (Khandaker & Rahman 2022). Iron and manganese, although essential, in high concentrations

make water unattractive and have a potential impact on the neurological system (Koley 2021).

4.1.5. Salinity and Hardness

Excessive levels of total dissolved solids and hardness in water cause kidney stones, heart and blood vessel problems, and stomach upsets (Ahada & Suthar 2019). Increased levels of sodium also pose a threat, especially for people suffering from hypertension or heart conditions (Yadav et al. 2019).

5. HEALTH RISK ASSESSMENT

For assessing the health risks associated with groundwater contamination, the hazard quotient (HQ) and hazard index (HI) are often employed as measures of noncarcinogenic risk, while the cancer risk (CR) is applied to those substances that are known to induce carcinogenicity (Khandaker & Rahman 2022).

If the HQ or HI is greater than or equal to 1, it indicates that a particular compound poses a potential health risk. In different regions of India, it has been reported that the HI value is greater than unity for 71 to 78% of groundwater samples contaminated with fluoride and nitrate (Marghade et al. 2023). Estimated cancer risks in areas contaminated by arsenic often approach and even surpass the acceptable limit of 10^{-4} . Several regional health risk assessment studies have reported that approximately 60–80% of analyzed groundwater samples in arsenic-affected regions exceeded the acceptable carcinogenic risk threshold. (Gupta et al. 2015). This also emphasizes the seriousness of the long-term risks of arsenic exposure through drinking water and the need to address the problem. Children are also at a higher risk than adults because they weigh less and drink more water per kilogram of weight than adults (Singh et al. 2024). Other vulnerable groups, like pregnant women, elderly people, and people with weakened immune systems, are also more susceptible to health risks and, therefore, need special consideration in assessing health risks and planning for water management (Awasthi et al. 2021). Fig. 5 illustrates the major routes by which groundwater contamination translates to human health effects. Considering the health hazards associated with groundwater contamination, the need for remediation and treatment of the groundwater arises. The next section highlights the different technologies that can be used in mitigating groundwater contamination in India. For assessing the health risks associated with groundwater contamination, the hazard quotient (HQ) and hazard index (HI) are often employed as measures of noncarcinogenic risk, while the cancer risk (CR) is applied to those substances that are known to induce carcinogenicity. Comparative evaluation of reviewed studies indicates that fluoride and arsenic contamination generally contribute the highest non-carcinogenic and carcinogenic health risks in

Indian groundwater systems. Fluoride-related hazard index (HI) values greater than 1 have frequently been reported in semi-arid and hard rock regions of Rajasthan, Telangana, Karnataka, and Tamil Nadu, particularly among children due to lower body weight and higher water intake rates. Similarly, arsenic-contaminated groundwater in eastern India, especially West Bengal and Bihar, consistently shows elevated carcinogenic risk values exceeding the acceptable USEPA risk threshold of 10^{-4} . In contrast, nitrate-associated health risks are comparatively more localized and strongly linked with agriculturally intensive regions such as Punjab, Haryana, and Gujarat. The reviewed literature therefore suggests that health risks from groundwater contamination in India are highly region-specific and controlled by both hydrogeological conditions and anthropogenic activities.

6. WATER QUALITY STANDARDS AND INDICES

Groundwater quality assessment entails its comparison with drinking water standards to ascertain its suitability for domestic consumption and other uses. These standards are guidelines for establishing benchmark values for water quality, which are used for the evaluation, regulation, and management of water quality. Besides individual parameter analysis, water quality indices are also employed to integrate two or more parameters into a single value, which is a measure of water quality in general.

6.1. Regulatory Standards

Groundwater quality standards are determined by different national and international organizations, with the aim of protecting public health and providing safe drinking water. In India, the Bureau of Indian Standards (BIS) specifies the limits of safe and permissible concentrations of drinking water quality, and the World Health Organization (WHO)

specifies the global reference values for the same. The United States Environmental Protection Agency (USEPA) specifies the guideline values, which are often used as a reference by researchers in their studies. These guidelines specify the threshold concentrations of major ions, nutrients, heavy metals, etc., present in the groundwater. These limits are critical in order to avoid health risks associated with the long-term use of groundwater (Krishnappa et al. 2024). Table 4 summarizes the permissible limits recommended by BIS, WHO, and USEPA for key groundwater quality parameters.

6.2. Water Quality Index (WQI)

The Water Quality Index (WQI) is a method to represent the overall water quality of groundwater with a single number, obtained by calculating several physicochemical parameters (Hanse et al. 2019). By integrating multiple water quality factors, the Water Quality Index simplifies complex information for decision-makers and water resource managers. Among the various methods available, the weighted arithmetic index method is the most popular method used in Indian groundwater research due to its ease and efficiency. However, significant methodological variability exists among WQI studies conducted across India. Different researchers have used different combinations of physicochemical parameters, weighting schemes, normalization approaches, and classification criteria depending on regional hydrogeological conditions and study objectives. While some studies considered only major physicochemical parameters such as pH, TDS, hardness, fluoride, and nitrate, others additionally incorporated heavy metals, microbiological indicators, or irrigation-related parameters. Similarly, the relative weights assigned to parameters vary considerably among studies, which directly influences the final WQI values and classification

Table 4: Permissible limits for groundwater quality parameters (Krishnappa et al. 2024).

Parameter	Unit	BIS (IS10500:2012)	WHO (2017)	USEPA (2018)	Health Significance
pH	-	6.5–8.5	6.5–8.5	6.5–8.5	Corrosion, taste
TDS	mg L ⁻¹	500 (2000) *	1000	500	Taste, scaling
Total Hardness	mg L ⁻¹	200 (600) *	500	-	Scaling, cardiovascular
Fluoride	mg L ⁻¹	1.0 (1.5) *	1.5	4.0	Fluorosis
Nitrate	mg L ⁻¹	45	50	45	Methemoglobinemia
Arsenic	µg L ⁻¹	10	10	10	Cancer, Arsenicosis
Lead	µg L ⁻¹	10	10	15	Neurological damage
Cadmium	µg L ⁻¹	3	3	5	Kidney damage
Chromium	µg L ⁻¹	50	50	100	Cancer
Iron	mg L ⁻¹	0.3	-	0.3	Aesthetic, taste

Note: Adapted from the Bureau of Indian Standards (IS 10500:2012), World Health Organization (2017), and USEPA (2018).

*Values in parentheses indicate maximum permissible limits in the absence of alternate sources.

categories. Therefore, direct comparison of WQI values across different regions should be interpreted cautiously, as methodological differences may contribute to variations in reported groundwater quality classifications. (Mukherjee & Singh 2022). The construction of WQI entails four major

stages, which include the selection of appropriate water parameters, assigning weights to each parameter depending on their relative importance in ensuring safe water for human consumption, determining individual water quality ratings, and finally, integrating all ratings to form a single index

Table 5: Major groundwater contaminants and associated health effects.

Contaminant	Sources	Affected Regions	Concentration Range	Health Effects	Indicative Population Exposure	Authors
Fluoride	Geogenic (granite, gneiss weathering)	Rajasthan, Tamil Nadu, Andhra Pradesh, Karnataka, Madhya Pradesh	1.5–12 mg L ⁻¹	Dental fluorosis, skeletal fluorosis, neurological effects	66 million	Das et al. 2009, Ghosh et al. 2024, Gupta et al. 2019
Arsenic	Geogenic (Fe-oxyhydroxide reduction)	West Bengal, Bihar, Uttar Pradesh, Assam, Chhattisgarh	10–3200 µg L ⁻¹	Arsenicosis, skin lesions, cancer (skin, lung, bladder)	Estimated from regional studies and published assessments	Gupta et al. 2015, Cherukumilli et al. 2018, Inaniyan & Raychoudhury 2019
Nitrate	Agricultural fertilizers, sewage, animal waste	Punjab, Haryana, Gujarat, Tamil Nadu, Delhi NCR	45–300 mg L ⁻¹	Methemoglobinemia (infants), gastric cancer, thyroid disorders	Estimated from regional agricultural groundwater studies	Marghade et al. 2023, Kumar et al. 2017, Mandal & Sanyal 2019
Heavy Metals (Pb, Cd, Cr, Cu, Zn)	Industrial effluents, mining, E-waste	Gujarat, Maharashtra, West Bengal, Jharkhand, Tamil Nadu	Variable (exceeding limits)	Neurological damage, kidney dysfunction, cancer, developmental disorders	Localized population exposure reported	Singh et al.2024, Awasthi et al.2021, Khandaker & Rahman, 2022
Salinity (High TDS, Cl ⁻)	Seawater intrusion, evaporation, mineral dissolution	Coastal regions (Gujarat, Tamil Nadu, Kerala, Goa), arid zones	TDS: 1500–5000 mg L ⁻¹	Kidney stones, cardiovascular issues, gastrointestinal disorders	35–45 million	Ahada & Suthar 2019, Hussain et al. 2012

Note: Population estimates are indicative and derived from multiple regional studies; actual values may vary with spatial and temporal factors. Values in Table 5 are compiled from regional investigations and published assessments rather than standardized national-scale datasets.

Table 6: Water Quality Index (WQI) classification and interpretation.

WQI Range	Water Quality Category	Suitability for Drinking	Treatment Required	Typical Characteristics	% of Indian Samples	Authors
<50	Excellent	Suitable without treatment	None	All parameters well within limits	15–25%	Hanse et al. 2019, Mukherjee & Singh 2022, Sahoo et al. 2024
50–100	Good	Suitable with minimal treatment	Simple disinfection	Most parameters within limits	40–64%	Hanse et al. 2019, Mukherjee & Singh 2022, Sahoo et al. 2024
100–200	Poor	Requires treatment	Conventional treatment	Some parameters exceed limits	8–35%	Ravindra & Mor.2019, Adimalla et al.2018
200–300	Very Poor	Requires advanced treatment	Advanced treatment needed	Multiple parameters exceed limits	2–15%	Ravindra & Mor 2019, Adimalla et al. 2018, Ahada & Suthar 2019)
>300	Unsuitable	Not suitable for drinking	Extensive treatment/alternate source	Severe contamination	3–11%	Ravindra & Mor. 2019, Adimalla et al.2018 Ahada & Suthar.2019

Note: Percentage ranges represent compiled estimates from multiple regional studies and may vary with hydrogeological conditions, sampling methods, and spatial coverage. These values represent trends from reviewed regional studies, not national-scale statistical estimates.

Table 7: Comparative health risk trends reported in Indian groundwater studies.

Contaminant	Dominant Risk Type	Most Vulnerable Regions	Commonly Affected Population	Reported Risk Trend	References
Fluoride	Non-carcinogenic (HI > 1)	Rajasthan, Telangana, Karnataka, Tamil Nadu	Children	Very High	Das et al. 2009, Ghosh et al. 2024, Gupta et al. 2019, Ahada & Suthar 2019
Arsenic	Carcinogenic (CR > 10 ⁻⁴)	West Bengal, Bihar, Assam	General population	Very High	Gupta et al. 2015, Cherukumilli et al. 2018, Inaniyan & Raychoudhury 2019
Nitrate	Non-carcinogenic	Punjab, Haryana, Gujarat	Infants	Moderate–High	Kumar et al. 2017, Mandal & Sanyal 2019, Marghade et al. 2023
Heavy Metals	Carcinogenic and non-carcinogenic	Industrial and mining regions	Adults and children	Moderate	Singh et al. 2024, Awasthi et al. 2021, Khandaker & Rahman 2022
Salinity/TDS	Chronic exposure-related risk	Coastal and arid regions	General population	Moderate	Ahada & Suthar 2019, Hussain et al. 2012

Note: Comparative health-risk trends are synthesized from published regional groundwater assessment studies conducted across different hydrogeological settings in India.

value (Sahoo et al. 2024). The parameters with higher health effects, e.g., nitrates, fluoride, and TDS, have higher weights in the equation. After determining the WQI, groundwater is normally grouped into five classes of water quality, namely, excellent (<50), good (50–100), poor (100–200), very poor (200–300), and not suitable for consumption (>300) (Ravindra & Mor 2019). Several reviewed studies have reported that a considerable proportion of groundwater samples fall within the “good” water quality category; however, the reported percentages vary depending on the adopted WQI methodology, parameter selection, weighting scheme, and classification criteria used in different studies. Approximately 15–30% lie in the poor to very poor range. Higher values of WQI are recorded in industrial and arid zones, while better water quality is recorded in zones with higher rates of recharge (Adimalla et al. 2018). The areas where the water quality index (WQI) is higher are the areas where industries are located, farming is done, and where the climate is arid to semi-arid (Ahada & Suthar 2019). It is worth noting that the differences in the percentages and concentration range reported by different studies can be attributed to differences in spatial coverage, sampling frequency, and methods of analysis. Thus, the data presented in this review is best interpreted as trends rather than exact national data. Further studies need to be undertaken on standardized data collection methods and statistical aggregation. Most of the studies have used normalized groundwater quality parameters against standard guideline values like BIS (IS 10500) and WHO limits. In the present review, the results have been interpreted in relation to standard values instead of comparing the actual values from different studies, which facilitates evaluation (Krishnappa et al. 2024). The WQI trends summarized in this review are therefore intended to provide indicative comparative

assessment of groundwater quality conditions across regions rather than standardized national-scale WQI rankings.

7. INTEGRATED ASSESSMENT FRAMEWORK FOR GROUNDWATER QUALITY IN INDIA

Based on the synthesis of the literature, an integrated framework has been proposed for connecting groundwater contamination sources, hydrogeochemical processes, health risks, and remediation strategies in the Indian context (Mukherjee & Singh 2022, Malik et al. 2023). Both anthropogenic and geogenic sources of groundwater contamination have been identified, and the hydrogeochemical processes influencing groundwater quality have also been analyzed (Singh et al. 2022, Yadav et al. 2018). Human health risks from groundwater contamination have also been discussed in the framework. Remediation strategies have also been included in the framework for addressing groundwater contamination in the Indian scenario. This framework will help in understanding groundwater contamination in a better way for sustainable groundwater management in the country. This integrated framework represents a novel contribution of this review by linking contamination sources, hydrogeochemical processes, health risks, and remediation strategies in a unified manner.

8. REMEDIATION STRATEGIES

The diagram presents the Integrated Remediation Technology Framework for Indian Groundwater. It is a decision guide to select appropriate cleanup technologies based on the nature of the contaminant, concentration of the contaminant, scale of the problem, and social and economic factors. The framework includes source control technologies, in-situ technologies like PRBs and phytoremediation, ex-situ




Fig. 6: Framework for selection of groundwater remediation technologies based on contaminant type and site conditions.

technologies like RO, adsorption, and electrocoagulation, and monitoring technologies to address the issue of groundwater in a holistic manner. Fig. 6 demonstrates that selection of groundwater remediation technologies depends on contaminant type, contaminant concentration, treatment scale, economic feasibility, and operational complexity. The framework indicates that advanced membrane technologies are highly effective for multi-contaminant removal but are associated with high operational cost and energy demand. In contrast, adsorption-based and nature-based technologies are more suitable for decentralized rural applications because of lower operational requirements and improved sustainability.

9. CONVENTIONAL TREATMENT TECHNOLOGIES

9.1. Reverse Osmosis (RO)

Reverse osmosis has emerged as a preferred treatment

technology for groundwater in India because of its high removal capabilities for a wide array of impurities. In this method, water is pressurized to pass through a semi-permeable membrane, resulting in a reduction of fluoride levels by 85–95%, arsenic by 90–98%, nitrate by 85–92%, heavy metals by 90–99%, and total dissolved solids by 95–98%. RO systems have been found to be highly effective and are used across residential, institutional, and community water treatment systems. However, there are some limitations to this technology, such as high energy requirements, membrane fouling, maintenance, and the difficulty of handling the reject brine. The cost of the equipment varies widely, from ₹5,000 to ₹25,000 for household systems to ₹5 to ₹50 lakhs for community systems (Muthusamy et al. 2023).

9.2. Nanofiltration (NF)

Nanofiltration operates at lower pressures compared to reverse osmosis treatment. The process has a strong ability

to remove divalent ions, organic matter, and a certain amount of dissolved solids. The process has a removal efficiency of 70–85% (Nagaraj & Masilamani, 2023). The process is effective in reducing hardness, sulfates, and some fluoride in groundwater supplies. Due to the low energy consumption and less brine disposal compared to reverse osmosis, the process is regarded as a cost-effective solution for brackish water treatment (Bhuiyan & Ray 2017).

9.3. Activated Alumina Adsorption

Activated alumina is frequently utilized for the adsorption-based removal of fluoride and arsenic from water. It has been seen to remove fluoride to the extent of 80 to 90%, and arsenic to the extent of 75 to 85%. However, this depends on the water being treated (Yasaswini et al. 2024). The method is preferred for its simple design, easy operation, and low cost, estimated at ₹2,000–₹10,000 for domestic-scale treatment. However, its effectiveness is pH-dependent, and regeneration or replacement is required periodically (Mukherjee et al. 2021).

9.4. Ion Exchange

Ion exchange resins selectively extract impurities, like fluoride, nitrates, and heavy metals, from groundwater. The method usually eliminates 85–95% of impurities, depending upon the type of resin and water quality (Thalod & Sharma 2024). However, ion exchange resins also have a drawback, as their periodic regeneration is necessary using chemical solutions, leading to a disposal problem. The effectiveness of ion exchange resins is also restricted by the lack of proper disposal facilities for solid wastes (Mandal & Sanyal 2019).

10. ADVANCED AND EMERGING TREATMENT TECHNOLOGIES

10.1. Electrocoagulation

Electrocoagulation has been shown to be a viable electrochemical technique for treating groundwater contaminated by fluoride, arsenic, heavy metals, and suspended solids. In this technique, coagulant species are produced in situ through sacrificial dissolution of electrodes, with reported removal efficiencies of 75–90% for fluoride, 80–95% for arsenic, and 85–98% for heavy metals. The primary advantages of this technique include the elimination of the need for chemical dosing, a compact design, and low levels of sludge formation. However, the technique has some limitations in real-world application, which include energy consumption (0.5–3 kWh m⁻³), electrode passivation, and the need for electrode replacement (Kalaiselvi & Pillai 2019, Vijaylaxmi & Khan 2021).

10.2. Adsorption-Based Technologies

Recently, breakthroughs in adsorption technology have led to the development of a new range of high-performing adsorbents for groundwater treatment. Biochar, clay, metal oxide, and complex nanostructured adsorbents have shown promising adsorption capacities for the removal of fluoride, arsenic, and various heavy metals, with removal efficiencies ranging from 80–99% (Saha et al. 2014). Of specific interest are adsorbents made from agricultural waste materials, which are rice husk, coconut shells, and neem leaves, and are gaining popularity as a cost-effective, green technology solution for groundwater treatment in rural communities (Karunanidhi et al. 2024). However, there are still challenges in the adsorption process, which include the regeneration of the adsorbents, the adsorbent's lifespan, and the disposal of the exhausted adsorbents, which need further optimization (Kumar et al. 2017).

10.3. Phytoremediation

Phytoremediation utilizes the natural uptake of pollutants by plants, thereby purifying water and soil. Water-loving plants, like the water hyacinth, duckweed, and vetiver, and those on land, like the sunflower, mustard, and ferns, have been observed to take up heavy metals, nutrients, and organic chemicals (Muthusamy et al. 2023). It offers an environmentally friendly and cost-effective approach for the remediation of shallow aquifers, groundwater with links to surface water, and constructed wetlands (Nagaraj & Masilamani 2023). However, its application is restricted by the low rate of cleanup, seasonal fluctuations, and the management and disposal of the polluted vegetation (Bhuiyan & Ray 2017).

10.4. Membrane Bioreactors (MBRs)

Membrane bioreactors utilize the combination of a biological process and membrane filtration for the effective removal of organic matter, nutrients, and pathogens (Yasaswini et al. 2024). The process has been found to be effective in the treatment of polluted groundwater and wastewater intended for aquifer recharge and reuse. The process has the advantage of high treatment efficiency in a space-saving design; however, the problem of membrane fouling and the operating costs need to be taken into consideration (Mukherjee et al. 2021).

10.5. Permeable Reactive Barriers

The permeable reactive barrier (PRB) technology is an innovative groundwater remediation technology. In the PRB technology, the contaminated groundwater is treated with the help of a reactive medium such as zero-valent iron,

activated carbon, or limestone (Thalod & Sharma 2024). The groundwater flows through the medium and is treated. The technology has been found to treat heavy metals, chlorinated organic compounds, and nutrient compounds (Cherukumilli et al. 2017).

10.6. Technology Selection and Implementation

The choice of technology is based on the type of contaminant present and its concentration, the quality characteristics of the water to be treated, the scale of the treatment (residential, community, or regional), cost viability, the complexity of the technology, and local technical capacity (Kalaiselvi & Pillai 2019). Table 8 summarizes the various remediation technologies. In most cases, the best results are achieved by combining different technologies into a hybrid technology.

For these technologies to be effective, the community needs to be engaged with capacity building and sustainable funding models. In rural and disadvantaged areas, policy support, subsidies, and awareness are key to the use of the technologies. Comparative evaluation of groundwater remediation technologies indicates that no single treatment method is universally suitable for all groundwater contamination conditions in India. Reverse osmosis demonstrates the highest overall removal efficiency for fluoride, arsenic, nitrate, salinity, and heavy metals; however, its high energy requirement, membrane fouling, and reject brine disposal significantly limit its applicability in economically weaker rural regions. Nanofiltration provides comparatively lower operational pressure and energy consumption but is less effective for complete dissolved contaminant removal.

Table 8: Comparison of groundwater remediation technologies.

Technology	Target Contaminants	Removal Efficiency	Capital Cost	Operating Cost	Advantages	Limitations	Rural Applicability	Authors
Reverse Osmosis	Fluoride, arsenic, nitrate, heavy metals, TDS	85–98%	High (₹5,000–50 Lakh)	High	High efficiency, multiple contaminants	Energy-intensive, brine disposal	Moderate	Muthusamy et al. 2023, Nagaraj & Masilamani 2023, Bhuiyan & Ray 2017
Activated Alumina	Fluoride, arsenic	75–90%	Low (₹2,000–10,000)	Low	Simple, low cost	pH-dependent, regeneration	High	Yasaswini et al. 2024, Mukherjee et al. 2021
Electrocoagulation	Fluoride, arsenic, heavy metals	75–98%	Moderate (₹10,000–5 Lakh)	Moderate	No chemicals, compact	Electrode replacement, energy	Moderate	Kalaiselvi & Pillai 2019, Vijaylaxmi & Khan 2021
Adsorption (Biochar, modified materials)	Fluoride, arsenic, heavy metals	80–99%	Low (₹1,000–8,000)	Very low	Low cost, sustainable	Regeneration, disposal	Very High	Saha et al. 2014, Karunanidhi et al. 2024, Kumar et al. 2017
Phytoremediation	Heavy metals, nutrients, organics	60–85%	Very low (₹500–5,000)	Very low	Eco-friendly, sustainable	Slow, seasonal, biomass disposal	High	Muthusamy et al. 2023, Nagaraj & Masilamani 2023, Bhuiyan & Ray 2017
Ion Exchange	Fluoride, nitrate, heavy metals	85–95%	Moderate (₹5,000–20,000)	Moderate	Selective removal	Brine disposal, regeneration	Moderate	Mandal & Sanyal 2019, Thalod & Sharma 2024
Nanofiltration	Hardness, sulfate, moderate TDS	70–85%	Moderate-High (₹8,000–30 Lakh)	Moderate	Lower pressure than RO	Partial removal	Low–Moderate	Nagaraj & Masilamani 2023, Bhuiyan & Ray 2017

Note: Synthesized from groundwater treatment and remediation studies. Cost ranges are indicative and may vary with system capacity, location, and operational conditions. Rural applicability in Table 8 is based on operational simplicity, energy requirements, maintenance demand, waste disposal complexity, and economic feasibility reported in the reviewed studies.

Adsorption-based technologies using activated alumina, biochar, and modified low-cost materials are considered more suitable for rural and decentralized applications because of their lower cost, operational simplicity, and minimal energy requirements, although periodic adsorbent regeneration and disposal remain important operational concerns. Electrocoagulation has shown promising removal efficiency for fluoride, arsenic, and heavy metals, but field-scale implementation is constrained by electrode passivation, electricity demand, and sludge management requirements. Similarly, ion exchange systems provide effective selective contaminant removal but generate secondary brine waste requiring appropriate disposal. Therefore, technology selection in the Indian context should consider contaminant type, hydrogeochemical conditions, treatment scale, operational complexity, waste disposal requirements, energy availability, and socio-economic feasibility, particularly in rural and resource-limited regions.

11. FUTURE PERSPECTIVES AND RESEARCH DIRECTIONS

11.1. Emerging Research Priorities

Future areas of research should address the following gaps in knowledge about groundwater quality in India. Some of the areas that need priority in the future are the following: emerging contaminants in groundwater, which include pharmaceuticals, pesticides, microplastics, and PFAS (Pal et al. 2009, Sultan et al. 2024). There is a pressing need for treatment technologies that are cost-effective, energy-efficient, and adaptable to local conditions, especially in rural areas (Karunanidhi et al. 2024). Advances in monitoring and forecasting can be achieved through integrating remote sensing, GIS, and machine learning, which will facilitate real-time continuous evaluation of groundwater quality (Thalod & Sharma 2024). Moreover, the impacts of climate change on groundwater recharge, the mobilization of contaminants, and the long-term trends in water quality require special research attention (Kaur et al. 2019). The research focus of the health sector needs to move beyond the examination of individual factors to the evaluation of the collective impacts of the risks that arise from the joint action of various contaminants (Khandaker & Rahman 2022). Moreover, the role of managed aquifer recharge (MAR) as a two-fold approach to enhancing groundwater quality and increasing its storability needs to attract more research attention on a regional scale (Dhakshinamoorthy & Selvarajan 2025). Based on the trends in groundwater quality being reported from different regions of India, it has been assessed that regions with high intensity in agricultural practices and groundwater depletion will have high nitrate and salinity problems in the near future (Kaur et

al. 2019, Kumar et al. 2017). Similarly, changes in recharge patterns due to climate variability will enhance the mobilization of geogenic contaminants like fluoride and arsenic in groundwater (Dhakshinamoorthy & Selvarajan 2025).

12. POLICY AND MANAGEMENT RECOMMENDATIONS

For effective groundwater quality management in India, it is important to adopt a multi-tier strategy. Strengthening the regulatory system and enforcement of regulations are significant steps towards curbing pollution caused by agricultural, industrial, and urban activities (Shukla & Saxena 2021). Preventive measures should focus on source control, including the adoption of sustainable agriculture, better management of industrial effluent disposal, and scientific disposal of solid and liquid waste (Kumar et al. 2017). A robust, comprehensive groundwater quality monitoring system is vital, which is supported by real-time data collection and information sharing among various regulatory agencies, scientists, and local governments (Gupta et al. 2022). Community engagement must be encouraged through decentralized water quality management, monitoring, and capacity-building programs to empower people to protect and manage groundwater resources (Hossain et al. 2021). Financial incentives, subsidies, and technical support are also necessary to promote the use of water treatment systems at the individual and community levels, especially in rural and disadvantaged areas (Karunanidhi et al. 2024). Another significant factor is to enhance public awareness of groundwater contamination issues and the associated health risks with the help of effective educational campaigns (Das et al. 2009). Ultimately, the key to sustainability is to enhance interdisciplinary research and cooperation between policymakers, scientists, and stakeholders to develop effective groundwater management strategies that are tailored to specific contexts (Kaliyan et al. 2022).

12.1. Technological Innovations

Recent advances in groundwater management reveal some promising technological approaches. Nanotechnology adsorbents and membrane filtration have demonstrated better efficacy in removing a range of impurities, from fluoride and arsenic to heavy metals (Saha et al. 2014). In energy-scarce environments, solar-powered water treatment could be a viable solution for reliable operation in a rural setting (Muthusamy et al. 2023). Real-time groundwater quality monitoring is being strengthened with the help of smart sensor technologies. Internet of Things (IoT) technology, for instance, enables rapid detection of changes in water quality through IoT-based monitoring systems. Simultaneously, AI and machine learning techniques are being used for the

prediction and assessment of groundwater quality (Thalod & Sharma 2024). Nature-based and hybrid technologies, like constructed wetlands and bioremediation, are gaining prominence as they operate with low operating costs and in a highly environmentally friendly manner (Nagaraj & Masilamani 2023). Decentralized treatment technologies, which are designed for households and communities, also hold promise as a solution for groundwater pollution, especially in rural areas (Karunanidhi et al. 2024).

12.2. Sustainable Water Resource Management

Ensuring the sustainability of groundwater resources over the long term requires an integrated water resources approach that connects demands, supplies, and ecosystems. In terms of cutting back on water use, the focus should be on saving water, improving irrigation efficiency, and enhancing the efficiency of water use (Prasad & Rao 2018). In terms of supplying more water, the focus should be on harvesting rainwater, recharging groundwater, and safely reusing treated water to enhance groundwater supplies while lessening the pressure on natural groundwater supplies (Dhakshinamoorthy & Selvarajan 2025). The key to the prevention of further deterioration in the quality of groundwater lies in the effective control of pollution, which includes the reduction of pollution, the application of best management practices, and the enforcement of environmental laws (Kumar et al. 2017). Another important factor is the protection and maintenance of ecosystems that rely on groundwater, which is necessary to maintain the quality of the water itself (Nagaraj & Masilamani 2023). Sustainable water management should be equitable, ensuring everyone has access to water that is safe and reliable, particularly for vulnerable and marginalized groups (Rajendran et al. 2024). As the climate changes, with shifts in rainfall, droughts, and increasingly extreme events, water management strategies should be responsive to these changes to create long-term sustainability (Kaur et al. 2019). Reaching the Sustainable Development Goal for water and sanitation, SDG 6, requires collaboration and coordination among government agencies, scientists, businesses, and social organizations (Shukla & Saxena 2021).

13. LIMITATIONS OF EXISTING STUDIES

Although extensive research has been conducted on groundwater quality in India, several limitations have been identified. One major limitation is that many studies are localized and based on limited sample sizes, which limits their representativeness (Hanse et al. 2019, Mukherjee & Singh 2022). Another limitation is the lack of long-term monitoring data, which limits understanding of temporal

trends (Mawari et al. 2022). Sampling and analytical techniques have also been inconsistent (Ahada & Suthar 2018). These limitations indicate that current groundwater research in India lacks sufficient organization and standardization, making it difficult to develop integrated, policy-relevant groundwater management strategies. Large, standardized monitoring programs, consistent analytical methods, and long-term datasets are therefore needed for reliable comparison and prediction of groundwater quality.

13.1. Data Variability and Comparability Considerations

Integrating the groundwater quality data from various studies may pose difficulties due to the inconsistencies in spatial distribution, time scales, and methodologies. The hydrogeological conditions, such as the presence of alluvial or hard rock formations, affect the movement of water and the presence of impurities. Although the review has provided an integrated overview of the groundwater quality trends in India, the results must be considered indicative rather than exact quantitative data (Mawari et al. 2022, Ahada & Suthar 2018). Population exposure estimates and contamination percentages reported in the reviewed literature are derived from regional investigations conducted under different hydrogeological and methodological conditions and should therefore be interpreted as indicative estimates rather than precise national-scale statistics.

13.2. Integrated Perspective on Groundwater Contamination

Groundwater contamination in India is a multifaceted phenomenon involving the interrelationship between sources of contamination, hydrogeochemical processes, and human health impacts (Mukherjee & Singh 2022, Malik et al. 2023). Human activities such as agriculture, industrial discharge, and urbanization are sources of groundwater contamination in India, while hydrogeochemical processes play an important role in groundwater chemistry, increasing the concentration of contaminants such as fluoride, arsenic, nitrate, and heavy metals in groundwater. Human health impacts of groundwater contamination in India are significant, considering the high concentration of contaminants in groundwater. Therefore, groundwater management in India involves an interrelationship between sources of contamination, groundwater chemistry, human health impacts, and appropriate measures for groundwater remediation.

14. CONCLUSIONS

India's groundwater is a complex patchwork, and its quality is determined by a combination of rock, climate, and human factors. When viewed by region, there are specific areas of

contamination. Fluoride is observed in many states, arsenic is a major threat in eastern states, nitrates are present in areas of high agricultural activity, and heavy metals are found in areas of industrial activity. When viewed as a whole, a major share of the population is affected by groundwater contamination, which is a threat to long-term health and sustainability. When physicochemical parameters are examined, a large number of groundwater samples indicate a level of contamination in terms of total dissolved solids, hardness, and nitrates, which indicates a combination of natural and anthropogenic factors. The level of contamination is a result of a combination of geogenic factors, which include rock–water interaction and redox reactions, as well as anthropogenic factors, which include excessive use of fertilizers, industrial effluents, poor sewage disposal, and mining activity. Health risk assessments indicate a high level of non-carcinogenic as well as carcinogenic risks in areas of contamination. Fluorosis, arsenic toxicity, methemoglobinemia, and heavy metal toxicity are major public health concerns, especially in children. The Water Quality Index results indicate that a part of the groundwater is potable, but a major share of it needs to be treated or an alternative source of water needs to be found. There are several options for remediation and treatment, including membrane systems, adsorption, electrochemical treatments, and nature-based solutions, among others, even though none of these options are universally applicable. It should be noted, however, that effective action will involve matching the contaminant profile, water chemistry, scale, cost, and local technical capacity. The present review demonstrates that groundwater contamination in India exhibits strong regional variability controlled by hydrogeological conditions, climatic factors, land-use practices, and anthropogenic pressures. Arsenic contamination remains the most critical concern in the eastern alluvial aquifer systems because of its severe carcinogenic risks and widespread population exposure. In contrast, fluoride contamination dominates the hard-rock terrains of western and southern India, where prolonged groundwater residence time and rock–water interaction enhance fluoride mobilization. Nitrate contamination is primarily associated with agriculturally intensive regions of northern and western India, indicating the growing influence of unsustainable fertilizer usage and poor sanitation practices. Heavy metal contamination is more localized and concentrated near industrial and mining regions, particularly in central and western India. These observations indicate that groundwater management strategies in India should be region-specific rather than adopting a single uniform national approach. From a management perspective, priority should be given to strengthening regional groundwater quality monitoring networks, promoting source-control strategies,

improving rural drinking-water treatment infrastructure, and implementing sustainable agricultural and industrial practices. Cost-effective decentralized treatment systems and community-based groundwater management approaches are particularly important for rural and economically weaker regions. The review also identifies several important research gaps, including insufficient long-term monitoring datasets, limited studies on emerging contaminants, lack of standardized groundwater quality assessment methodologies, and inadequate integration of hydrogeochemical modeling with health-risk assessment frameworks. Future research should focus on multidisciplinary approaches integrating hydrogeochemistry, GIS, remote sensing, machine learning, and climate-change assessment to support sustainable groundwater management and policy development in India. In the future, it will be important to focus on the understanding of emerging contaminants, cost-effective treatment options, advanced monitoring and prediction tools, and the assessment of the impact of climate change on groundwater quality. Integrating these priorities will support decision-making by researchers, policymakers, water managers, and public health professionals. Addressing groundwater quality degradation in India remains an important future priority.

15. ACKNOWLEDGEMENT

We would like to express our sincere appreciation to SVKM NMIMS Global University, Dhule, and the individuals whose contributions and support have greatly enhanced the quality of this research.

16. REFERENCES

- Adimalla, N., 2020. Controlling factors and mechanism of groundwater quality variation in semiarid region of South India: an approach of water quality index (WQI) and health risk assessment (HRA). *Environmental Geochemistry and Health*, 42(6), pp.1725-1752. [DOI]
- Adimalla, N., Li, P. and Qian, H., 2019. Evaluation of groundwater contamination for fluoride and nitrate in semi-arid region of Nirmal province, South India: a special emphasis on human health risk assessment (HHRA). *Human and Ecological Risk Assessment: An International Journal*, 25(5), pp.1107-1124. [DOI]
- Adimalla, N., Qian, H. and Li, P., 2020. Entropy water quality index and probabilistic health risk assessment from geochemistry of groundwaters in hard rock terrain of Nanganur County, South India. *Geochemistry*, 80(4 Supplement), 125544. [DOI]
- Ahada, C.P.S. and Suthar, S., 2018. Groundwater nitrate contamination and associated human health risk assessment in southern districts of Punjab, India. *Environmental Science and Pollution Research*, 25(25), pp.25336-25347. [DOI]
- Ahada, C.P.S. and Suthar, S., 2019. Assessment of human health risk associated with high groundwater fluoride intake in southern districts of Punjab, India. *Exposure and Health*, 11(2), pp.267-275. [DOI]
- Alam, M.A., Mailapalli, D.R. and Maheshwari, B., 2024. Assessing groundwater quality, health risks, and policy implications: a case study

- of West Medinipur district, West Bengal, India. *World Water Policy*, 10(1), pp.341-362. [DOI]
- Ambade, B., Sethi, S.S., Patidar, K., Gautam, S. and AlShehri, M.A., 2024. Assessing variability and hydrochemical characteristics of groundwater fluoride contamination and its associated health risks in East Singhbhum district of Jharkhand, India. *Journal of Hazardous Materials*, 478, 135498. [DOI]
- Awasthi, S., Chauhan, R., Dwivedi, S., Srivastava, S. and Tripathi, R.D., 2021. Arsenic contamination of groundwater and its mitigation strategies. In: A. Singh, M. Agrawal and S.B. Agrawal (eds.) *Water Pollution and Management Practices*. Springer, Singapore, pp.107-119. [DOI]
- Barathkumar, S., Sellamuthu, K., Sathiyabama, K., Malathi, P., Kumaraperumal, R., Devagi, P. and Pazhanivelan, S., 2025. Groundwater quality assessment in selected polluted hotspots of Tamil Nadu, India using geospatial and statistical approaches. *Desalination and Water Treatment*, 322, 101047. [DOI]
- Basha, U.I., Rajasekhar, M., Ghosh, S., Das, P. and Suresh, U., 2022. Spatial assessment of groundwater quality using CCME-WQI and hydrochemical indices: a case study from Talupula Mandal, Ananthapuramu district, South India. *Applied Water Science*, 12, 168. [DOI]
- Bhuiyan, C. and Ray, P.K.C., 2017. Groundwater quality zoning in the perspective of health hazards. *Water Resources Management*, 31(1), pp.251-267. [DOI]
- Biswas, T., Paul, S., Ruidas, D., Saha, A., Shit, M., Islam, A.R.M.T., Islam, A. and Costache, R., 2023. Evaluation of groundwater contamination and associated human health risk in a water-scarce hard rock-dominated region of India: Issues, management measures and policy recommendation. *Groundwater for Sustainable Development*, 23, 101039. [DOI]
- Central Ground Water Board (CGWB), 2017. *Ground Water Year Book of Kerala (2016-2017)*. Ministry of Water Resources, River Development and Ganga Rejuvenation, Government of India. Available at: CGWB.
- Chakraborty, B., Roy, S., Bera, A., Adhikary, P.P., Bera, B., Sengupta, D., Bhunia, G.S. and Shit, P.K., 2021. Geospatial assessment of groundwater quality for drinking through water quality index and human health risk index in an upland area of Chota Nagpur Plateau of West Bengal, India. In: P.K. Shit, P.P. Adhikary and D. Sengupta (eds.) *Spatial Modeling and Assessment of Environmental Contaminants: Risk Assessment and Remediation*. Springer, Cham, pp.327-358. [DOI]
- Chaudhry, A.K. and Sachdeva, P., 2021. Groundwater fluoride contamination and probabilistic health-risk assessment in fluoride endemic areas of the Lower Satluj basin, Punjab, India. *Research Square* [Preprint]. [DOI]
- Cherukumilli, K., Delaire, C., Amrose, S. and Gadgil, A.J., 2017. Factors governing the performance of bauxite for fluoride remediation of groundwater. *Environmental Science & Technology*, 51(4), pp.2321-2328. [DOI]
- Cherukumilli, K., Maurer, T., Hohman, J.N., Mehta, Y. and Gadgil, A.J., 2018. Effective remediation of groundwater fluoride with inexpensively processed Indian bauxite. *Environmental Science & Technology*, 52(8), pp.4711-4718. [DOI]
- Das, B., Rahman, M.M., Nayak, B., Pal, A., Chowdhury, U.K., Mukherjee, S.C., Saha, K.C., Pati, S., Quamruzzaman, Q. and Chakraborti, D., 2009. Groundwater arsenic contamination, its health effects and approach for mitigation in West Bengal, India and Bangladesh. *Water Quality, Exposure and Health*, 1(1), pp.5-21. [DOI]
- Dhakate, R., More, S., Duvva, L.K. and Enjamuri, S., 2023. Groundwater chemistry and health hazard risk valuation of fluoride and nitrate enhanced groundwater from a semi-urban region of South India. *Environmental Science and Pollution Research*, 30(15), pp.43554-43572. [DOI]
- Dhakshinamoorthy, R. and Selvarajan, G., 2025. Comprehensive evaluation of groundwater quality in Mayiladuthurai District, Tamilnadu: physico-chemical and heavy metal analysis. *Ecology, Environment & Conservation*, 31(2), pp.865-872. [DOI]
- Dhanya Raj and Shaji, E., 2017. Fluoride contamination in groundwater resources of Alleppey, Southern India. *Geoscience Frontiers*, 8(1), pp.117-124. [DOI]
- Gayathri, J.A., Raj, V.T., Sreelash, K., Maya, K. and Padmalal, D., 2025. Comprehensive evaluation of groundwater quality and human health risk assessment in an ecologically sensitive mountainous catchment in Southern Western Ghats. *Human and Ecological Risk Assessment: An International Journal*, 31(7-8), pp.921-953. [DOI]
- Ghosh, A., Stening, J. and Chakraborty, R., 2024. Recovery of ecosystems pollution by contaminants of potential concern using phytoremediation techniques. *Integrated Environmental Assessment and Management*, 20(6), pp.1987-2002. [DOI]
- Goyal, H., Lanjewar, R., Chitransh, S. and Mondal, P., 2025. A comprehensive exploration of groundwater quality of Ambagarh Chowki region, Chhattisgarh, India: water quality index, health risk, and ANN predictive modeling. *Water Environment Research*, 97(6), e70125. [DOI]
- Gupta, D.P., Chaudhary, S., Singh, A., Shukla, R. and Mishra, V.K., 2023. Hydrochemical assessment of groundwater quality in Narmada River basin (Central India). *Water Science & Technology: Water Supply*, 23(2), pp.459-481. [DOI]
- Gupta, P., Niveditha, P.S. and Sharada, S., 2019. Treatment of contaminated ground water using phytoremediation technique at Anantapur, India. *World Environment*, 9(2), pp.46-55. [DOI]
- Gupta, P., Roy, S. and Mahindrakar, A.B., 2015. Treatment of groundwater using phytoremediation technique at Kolar Gold Fields, India. *International Journal of Environmental Engineering*, 7(1), pp.11-34. [DOI]
- Hanse, A., Chabukdhara, M., Baruah, S.G., Boruah, H. and Gupta, S.K., 2019. Fluoride contamination in groundwater and associated health risks in Karbi Anglong district, Assam, Northeast India. *Environmental Monitoring and Assessment*, 191(12), 782. [DOI]
- Hossain, M., Patra, P.K., Ghosh, B., Khatun, A. and Nayek, S., 2021. Sensitive assessment of groundwater-associated, multi-exposure health hazards in a fluoride-enriched region of West Bengal, India. *Environmental Geochemistry and Health*, 43(11), pp.4515-4532. [DOI]
- Hussain, I., Arif, M. and Hussain, J., 2012. Fluoride contamination in drinking water in rural habitations of central Rajasthan, India. *Environmental Monitoring and Assessment*, 184(8), pp.5151-5158. [DOI]
- Inaniyan, M. and Raychoudhury, T., 2019. Application of activated carbon-metal composite for fluoride removal from contaminated groundwater in India. *International Journal of Environmental Science and Technology*, 16(11), pp.7545-7554. [DOI]
- Kadam, A., Wagh, V., Jacobs, J., Patil, S., Pawar, N., Umrikar, B., Sankhua, R. and Kumar, S., 2022. Integrated approach for the evaluation of groundwater quality through hydro geochemistry and human health risk from Shivganga river basin, Pune, Maharashtra, India. *Environmental Science and Pollution Research*, 29(3), pp.4311-4333. [DOI]
- Kalaiselvi, N. and Pillai, V.K., 2020. Dendritic growth and spread of electrochemical research in India. *Proceedings of the Indian National Science Academy*, 86(2), pp.1001-1013. [DOI]
- Kalaivanan, K., Karuppaiah, S. and Duraisamy, S., 2022. Assessment of groundwater quality and human health risk impact on fluoride concentration: a case study of Namakkal district, South India. *Research Square* [Preprint]. [DOI]
- Karanveer, Bala, R. and Das, D., 2022. Geochemical and health risk assessment of potentially toxic trace elements and nitrate via groundwater in agro-ecosystem of alluvial plain Punjab, India. *Human and Ecological Risk Assessment: An International Journal*, 28(9), pp.983-1011. [DOI]

- Karunanidhi, D., Aravinthasamy, P., Deepali, M., Subramani, T. and Shankar, K., 2021. Groundwater pollution and human health risks in an industrialized region of Southern India: impacts of the COVID-19 lockdown and the monsoon seasonal cycles. *Archives of Environmental Contamination and Toxicology*, 80(1), pp.259-276. [DOI]
- Karunanidhi, D., Aravinthasamy, P., Roy, P.D., Praveenkumar, R.M., Prasanth, K., Selvapraveen, S., Thowbeekrahman, A., Subramani, T. and Srinivasamoorthy, K., 2020. Evaluation of non-carcinogenic risks due to fluoride and nitrate contaminations in a groundwater of an urban part (Coimbatore region) of South India. *Environmental Monitoring and Assessment*, 192(2), 102. [DOI]
- Karunanidhi, D., Aravinthasamy, P., Roy, P.D., Subramani, T. and Jayasena, H.C., 2024. Nitrate contamination in groundwater and its evaluation of non-carcinogenic health hazards from Arjunanadi river basin, South India. *Groundwater for Sustainable Development*, 25, 101153. [DOI]
- Karunanidhi, D., Aravinthasamy, P., Subramani, T. and Muthusankar, G., 2021. Revealing drinking water quality issues and possible health risks based on water quality index (WQI) method in the Shanmuganadhi river basin of South India. *Environmental Geochemistry and Health*, 43(2), pp.931-948. [DOI]
- Kaur, G., Sharma, S. and Garg, U.K., 2019. Assessment of ground water contamination by inorganic impurities in Ferozepur district of Punjab state, India. *Asian Journal of Chemistry*, 31(3), pp.515-521. [DOI]
- Kaur, L. and Godara, P., 2025. An investigation of water quality index and health risks of fluoride and nitrate in the arid groundwater (India). *Macedonian Journal of Ecology and Environment*, 27(2), pp.99-113. [DOI]
- Khan, A.F., Srinivasamoorthy, K., Prakash, R., Gopinath, S., Saravanan, K., Vinnarasi, F., Babu, C. and Rabina, C., 2021. Human health risk assessment for fluoride and nitrate contamination in the groundwater: a case study from the east coast of Tamil Nadu and Puducherry, India. *Environmental Earth Sciences*, 80(21), 724. [DOI]
- Khan, N., Malik, A. and Nehra, K., 2021. Groundwater hydro-geochemistry, quality, microbiology and human health risk assessment in semi-arid area of Rajasthan, India: a chemometric approach. *Environmental Monitoring and Assessment*, 193(4), 234. [DOI]
- Khandaker, N.R. and Rahman, M.M., 2022. Understanding sustainable arsenic mitigation technology application in the Indian Subcontinent. In: S. Yadav, A.M. Negm and R.N. Yadava (eds.) *Wastewater Assessment, Treatment, Reuse and Development in India*. Springer International Publishing, Cham, pp.87-105. [DOI]
- Koley, S., 2021. Contemporary practices in groundwater arsenic remediation and wastewater management in West Bengal, India: a systematic review. *International Journal of Advanced Technology and Engineering Exploration*, 8(80), pp.797-821. [DOI]
- Kouser, B., Bala, A., Verma, O., Prashanth, M., Khosla, A. and Pir, R.A., 2022. Hydrochemistry for the assessment of groundwater quality in the Kathua region, Jammu and Kashmir, India. *Applied Water Science*, 12, 143. [DOI]
- Krishnappa, S.M., Channabasavaiah, S., Hanché, D., Ananda, K.D. and Muthanarasimha, A.P., 2024. Evaluating health risks associated with fluoride and nitrate contaminants in drinking water to residents living in the Chikkaballapur Taluk of Karnataka, India. *Environmental Quality Management*, 34(2), e22315. [DOI]
- Kumar, G., Kothandaraman, S., Karuppanan, K. and Hussain, S., 2024. Groundwater quality and suitability assessment in Tirupur region, Tamil Nadu, India. *Journal of Chemistry*, 2024, 6083772. [DOI]
- Kumar, M., Patel, A.K., Das, A., Kumar, P., Goswami, R., Deka, P. and Das, N., 2017. Hydrogeochemical controls on mobilization of arsenic and associated health risk in Nagaon district of the central Brahmaputra Plain, India. *Environmental Geochemistry and Health*, 39(1), pp.161-178. [DOI]
- Madhavan, S., Kolanuvada, S.R., Sampath, V., Roy, P.D., Moorthy, P., Natarajan, L. and Chokkalingam, L., 2023. Assessment of groundwater vulnerability using water quality index and solute transport model in Poiney sub-basin of South India. *Environmental Monitoring and Assessment*, 195(2), 272. [DOI]
- Malik, A., Katyal, D., Narwal, N., Kataria, N., Ayyamperumal, R. and Khoo, K.S., 2023. Sources, distribution, associated health risks and remedial technologies for inorganic contamination in groundwater: a review in specific context of the state of Haryana, India. *Environmental Research*, 236, 116696. [DOI]
- Malik, N., Malik, A. and Bishnoi, S., 2021. Assessment of groundwater hydro-geochemistry, quality, and human health risk in arid area of India using chemometric approach. *Arabian Journal of Geosciences*, 14, 1546. [DOI]
- Mandal, J. and Sanyal, S., 2019. Geospatial analysis of fluoride concentration in groundwater in Puruliya district, West Bengal. *Space and Culture, India*, 6(5), pp.71-86. [DOI]
- Marghade, D., Malpe, D.B., Duraisamy, K., Patil, P.D. and Li, P., 2021. Hydrogeochemical evaluation, suitability, and health risk assessment of groundwater in the watershed of Godavari basin, Maharashtra, Central India. *Environmental Science and Pollution Research*, 28(15), pp.18471-18494. [DOI]
- Marghade, D., Mehta, G., Shelare, S.D., Jadhav, G. and Nikam, K.C., 2023. Arsenic contamination in Indian groundwater: from origin to mitigation approaches for a sustainable future. *Water*, 15(23), 4125. [DOI]
- Maurya, J.K., Pradhan, S.N., Seema and Ghosh, A.K., 2021. Evaluation of ground water quality and health risk assessment due to nitrate and fluoride in the Middle Indo-Gangetic plains of India. *Human and Ecological Risk Assessment: An International Journal*, 27(5), pp.1349-1365. [DOI]
- Maurya, S., More, A.B., Gupta, G., Saxena, A. and Singh, A., 2025. Hydrochemical evaluation of groundwater quality and human health risk assessment of fluoride and nitrate: a case study of Pratapgarh district, UP, India. *Water, Air, & Soil Pollution*, 236(3), 204. [DOI]
- Mawari, G., Kumar, N., Sarkar, S., Frank, A.L., Daga, M.K., Singh, M.M., Joshi, T.K. and Singh, I., 2022. Human health risk assessment due to heavy metals in ground and surface water and association of diseases with drinking water sources: a study from Maharashtra, India. *Environmental Health Insights*, 16, 1178630221146020. [DOI]
- Mhaske, G.D., Bhalla, R. and Nalawade, P.M., 2022. Study of groundwater hydrochemistry and drinking appropriateness using water quality index (WQI) modelling in the Jam River basin, India. *Ecology, Environment and Conservation*, 28, pp.S469-S477. [DOI]
- Mukherjee, A., Saha, D., Harvey, C.F., Taylor, R.G., Ahmed, K.M. and Bhanja, S.N., 2015. Groundwater systems of the Indian sub-continent. *Journal of Hydrology: Regional Studies*, 4, pp.1-14. [DOI]
- Mukherjee, A., Sarkar, S., Chakraborty, M., Duttagupta, S., Bhattacharya, A., Saha, D., Bhattacharya, P., Mitra, A. and Gupta, S., 2021. Occurrence, predictors and hazards of elevated groundwater arsenic across India through field observations and regional-scale AI-based modeling. *Science of the Total Environment*, 759, 143511. [DOI]
- Mukherjee, I. and Singh, U.K., 2022. Environmental fate and health exposures of the geogenic and anthropogenic contaminants in potable groundwater of Lower Ganga basin, India. *Geoscience Frontiers*, 13(3), 101365. [DOI]
- Mukherjee, I., Singh, U. and Patra, P.K., 2019. Exploring a multi-exposure-pathway approach to assess human health risk associated with groundwater fluoride exposure in the semi-arid region of east India. *Chemosphere*, 233, pp.164-173. [DOI]
- Muthusamy, P., Paneerselvam, B., Kaliyappan, S.P., Almohamad, H., Aldosari, A.A. and Abdo, H.G., 2023. Pollution source identification and suitability assessment of groundwater quality for drinking purposes in semi-arid regions of the southern part of India. *Water*, 15(22), 3995. [DOI]
- Nagaraj, S. and Masilamani, U.S., 2023. Hydrogeochemical and multivariate statistical approaches to investigate the characteristics of groundwater

- quality in fluoride-enriched hard rock region in Tirupathur district of Tamil Nadu, India. *Environmental Science and Pollution Research*, 30(44), pp.99809-99829. [DOI]
- Narsimha, A. and Sudarshan, V., 2017. Contamination of fluoride in groundwater and its effect on human health: a case study in hard rock aquifers of Siddipet, Telangana state, India. *Applied Water Science*, 7, pp.2501-2512. [DOI]
- Naskar, S., Mandal, D., Biswas, B. and Mukherjee, C., 2025. Hydrochemical assessment of groundwater quality using pollution index, quality indexes, fuzzy-GIS, graphical and multivariate techniques from a mining region of Eastern India. *Discover Water*, 5, 48. [DOI]
- Pal, P., Sen, M., Manna, A., Pal, J., Pal, P., Roy, S. and Roy, P., 2009. Contamination of groundwater by arsenic: a review of occurrence, causes, impacts, remedies and membrane-based purification. *Journal of Integrative Environmental Sciences*, 6(4), pp.295-316. [DOI]
- Pandey, S., Mohapatra, G. and Arora, R., 2023. Groundwater quality, human health risks and major driving factors in arid and semi-arid regions of Rajasthan, India. *Journal of Cleaner Production*, 427, 139149. [DOI]
- Panneerselvam, B., Muniraj, K., Pande, C.B., Ravichandran, N., Thomas, M. and Karuppannan, S., 2022. Geochemical evaluation and human health risk assessment of nitrate-contaminated groundwater in an industrial area of South India. *Environmental Science and Pollution Research*, 29(57), pp.86202-86219. [DOI]
- Panneerselvam, B., Ravichandran, N., Kaliyappan, S.P., Karuppannan, S. and Bidorn, B., 2023. Quality and health risk assessment of groundwater for drinking and irrigation purpose in semi-arid region of India using entropy water quality and statistical techniques. *Water*, 15(3), 601. [DOI]
- Pattanaik, P. and Bhowmick, P.K., 2020. Determination of water quality index of drinking water sources and health risk assessment of arsenic-contaminated rural areas in Basirhat-I Block of West Bengal. *International Journal of Innovative Technology and Exploring Engineering*, 9(3), pp.677-684. [DOI]
- Raja, V. and Neelakantan, M., 2021. Evaluation of groundwater quality with health risk assessment of fluoride and nitrate in Virudhunagar district, Tamil Nadu, India. *Arabian Journal of Geosciences*, 14(1), 52. [DOI]
- Rajendran, R., Lakshmanan, E., Muddu, S., Rajmohan, N., Senapathi, V. and Brindha, K., 2025. Multi-indices approach for comprehensive appraisal of groundwater quality and the implication on human health in the Amaravathi basin, Southern India. *Environmental Monitoring and Assessment*, 197(2), 153. [DOI]
- Ramalingam, S., Panneerselvam, B. and Kaliappan, S.P., 2022. Effect of high nitrate contamination of groundwater on human health and water quality index in semi-arid region, South India. *Arabian Journal of Geosciences*, 15, 242. [DOI]
- Rathinam, R.K., Ramajayam, J.G., Mahalingam, V. and Sarangapani, M., 2024. Assessment of hydrogeochemical characteristics and potable quality of groundwater around Retteri lake in Thiruvallur district, Tamil Nadu, India. *Zastita Materijala*, 65(3), pp.516-523. [DOI]
- Ravindra, K. and Mor, S., 2019. Distribution and health risk assessment of arsenic and selected heavy metals in groundwater of Chandigarh, India. *Environmental Pollution*, 250, pp.820-830. [DOI]
- Reddy, C.K.V.C., Golla, V., Badapalli, P.K. and Reddy, N.B.Y., 2022. Evaluation of groundwater contamination for fluoride and nitrate in Nellore urban province, Southern India: a special emphasis on human health risk assessment (HHRA). *Applied Water Science*, 12, 32. [DOI]
- Sabinaya, S., Mahanty, B., Lhamo, P., Sahoo, S.K., Jha, V.N. and Sahoo, N.K., 2023. Application of a multivariate water quality index to evaluate the groundwater geochemistry of Jajpur district, Odisha, India and assessment of potential health risks. *Research Square* [Preprint]. [DOI]
- Saha, A. and Pal, S.C., 2025. Assessment and characterization of groundwater quality and related human health risk in a fluoride-enriched semi-arid region of Eastern India. *Water, Air, & Soil Pollution*, 236(11), 691. [DOI]
- Saha, D., Dwivedi, S.N. and Singh, R.K., 2014. Aquifer system response to intensive pumping in urban areas of the Gangetic plains, India: the case study of Patna. *Environmental Earth Sciences*, 71(4), pp.1721-1735. [DOI]
- Saha, R., Wankhede, T., Majumdar, R.C. and Das, I., 2024. Pan India fluoride hazard assessment in groundwater. *Journal of Hazardous Materials*, 478, 135543. [DOI]
- Sahu, B.L., Banjare, G.R., Ramteke, S., Patel, K.S. and Matini, L., 2017. Fluoride contamination of groundwater and toxicities in Dongargaon Block, Chhattisgarh, India. *Exposure and Health*, 9(2), pp.143-156. [DOI]
- Sane, P.B., Gaikwad, S.P., Mahale, R.R., Muley, A., Supekar, A., Dongare, A.N. and Bhagat, A.A., 2025. Unusually high levels of fluoride and nitrate in groundwater in Southern India: water quality indices and associated health hazard implications. *Journal of Geosciences Research*, 10(2), pp.114-122. [DOI]
- Saraswat, A., Nath, T., Omeka, M.E., Unigwe, C.O., Anyanwu, I.E., Ugar, S.I., Latore, A., Raza, M.B., Behera, B., Adhikary, P.P., Scopa, A. and AbdelRahman, M.A.E., 2023. Irrigation suitability and health risk assessment of groundwater resources in the Firozabad industrial area of north-Central India: an integrated indexical, statistical, and geospatial approach. *Frontiers in Environmental Science*, 11, 1116220. [DOI]
- Sarma, R.V. and Singh, S.K., 2023. Assessment of groundwater quality and human health risks of nitrate and fluoride contamination in a rapidly urbanizing region of India. *Environmental Science and Pollution Research*, 30(19), pp.55437-55454. [DOI]
- Shaikh, H., Gaikwad, H., Kadam, A. and Umrikar, B., 2020. Hydrogeochemical characterization of groundwater from semiarid region of Western India for drinking and agricultural purposes with special reference to water quality index and potential health risks assessment. *Applied Water Science*, 10, 204. [DOI]
- Sharma, T., Litoria, P.K., Bajwa, B.S. and Kaur, I., 2021. Appraisal of groundwater quality and associated risks in Mansa district (Punjab, India). *Environmental Monitoring and Assessment*, 193(4), 159. [DOI]
- Shirke, K.D., Kadam, A. and Pawar, N.J., 2020. Temporal variations in hydro-geochemistry and potential health risk assessment of groundwater from lithological diversity of semi-arid region, western Gujarat, India. *Applied Water Science*, 10, 156. [DOI]
- Shukla, S. and Saxena, A., 2021. Appraisal of groundwater quality with human health risk assessment in parts of Indo-Gangetic alluvial plain, North India. *Archives of Environmental Contamination and Toxicology*, 80(1), pp.55-73. [DOI]
- Sidhu, N., Kaur, L., Rishi, M.S., Din, S.N.U., Tewari, K. and Singh, P., 2024. Integrating multivariate hydrogeochemical analysis with human health risk assessment: an inverse geochemical and statistical modeling approach. *Journal of Geochemical Exploration*, 258, 107389. [DOI]
- Singh, A., Maurya, S., Saxena, A. and Gupta, G., 2024. A review of phytoremediation techniques used for removal of arsenic from groundwater. In: A. Kuanar, A.P. Das, D. Kar and M.P. Shah (eds.) *Sustainable Management of Environmental Pollutants through Phytoremediation*. CRC Press, pp.135-156. [DOI]
- Singh, G., Rishi, M.S., Herojeet, R., Kaur, L. and Sharma, K., 2020. Evaluation of groundwater quality and human health risks from fluoride and nitrate in semi-arid region of Northern India. *Environmental Geochemistry and Health*, 42(7), pp.1833-1862. [DOI]
- Singh, K., Singh, R.D. and Pandey, G., 2023. Hydrogeochemistry, solute source identification, and health risk assessment of groundwater of cancer prone region in India. *Water Science & Technology: Water Supply*, 23(1), pp.317-342. [DOI]
- Singh, P., Rishi, M.S. and Kaur, L., 2024. Multi-parametric analysis of groundwater quality to assess human health risk and hydrogeochemical processes in an agriculturally intensive alluvial aquifer of Northwest India. *International Journal of Environmental Analytical Chemistry*, 104(11), pp.2552-2570. [DOI]

- Singh, R., Singh, S., Das, A., Sahu, H., Durgam, M., Khose, S., Pohshna, C. and Mailapalli, D., 2024. Evaluating groundwater quality for drinking and irrigation purposes and health risk assessment in West Midnapore district of West Bengal, India. *Journal of Agricultural Engineering (India)*, 61(4). [DOI]
- Singh, R., Upreti, P., Allemailem, K.S., Almatroudi, A., Rahmani, A.H. and Albalawi, G.M., 2022. Geospatial assessment of groundwater quality and associated health problems in the western region of India. *Water*, 14(3), 296. [DOI]
- Srinivas, Y., Oliver, D.H., Raj, A.S. and Chandrasekar, N., 2013. Evaluation of groundwater quality in and around Nagercoil town, Tamilnadu, India: an integrated geochemical and GIS approach. *Applied Water Science*, 3(3), pp.631-651. [DOI]
- Subramaniyan, A. and Ganesan, S., 2024. Human health risk and water quality assessment due to fluoride and nitrate around Cauvery river basin, Southern India. *Environmental Monitoring and Assessment*, 196(10), 880. [DOI]
- Sultan, M.W., Qureshi, F., Ahmed, S., Kamyab, H., Rajendran, S., Ibrahim, H. and Yusuf, M., 2025. A comprehensive review on arsenic contamination in groundwater: sources, detection, mitigation strategies and cost analysis. *Environmental Research*, 265, 120457. [DOI]
- Taloor, A.K., Sharma, S., Pir, R.A. and Kumar, K., 2024. Appraisal of health risks associated with exposure of fluoride and nitrate contaminated springs in the Doda Kishtwar Ramban (DKR) region of Jammu and Kashmir, India. *Journal of Geochemical Exploration*, 257, 107380. [DOI]
- Tanwer, N., Deswal, M., Khyalia, P., Laura, J.S. and Khosla, B., 2023. Fluoride and nitrate in groundwater: a comprehensive analysis of health risk and potability of groundwater of Jhunjhunu district of Rajasthan, India. *Environmental Monitoring and Assessment*, 195(2), 267. [DOI]
- Thalod, S. and Sharma, G., 2024. Arsenic contamination in groundwater of Haryana's National Capital Region and its comparative assessment (Mahendragarh, Rewari, and Jhajjar districts). *Cuestiones de Fisioterapia*, 53(3), pp.5027-5039. [DOI]
- Ullengula, M., Dhakate, R., Rao, N.S., Suryabhagavan, K.V., More, S., Gugulothu, S., Badapalli, P.K. and Ratnalu, G.V., 2026. Temporal dynamics of groundwater quality and associated health risks in a semi-arid region of South India. *Discover Sustainability*, 7, 298. [DOI]
- Vijayakumar, C.R., Balasubramani, D.P. and Azamathulla, H.M., 2022. Assessment of groundwater quality and human health risk associated with chromium exposure in the industrial area of Ranipet, Tamil Nadu, India. *Journal of Water, Sanitation and Hygiene for Development*, 12(1), pp.58-67. [DOI]
- Vijaylaxmi and Khan, S., 2021. Fluoride remediation using membrane processes. In: R. Prasad (ed.) *Environmental Pollution and Remediation*. Springer, Singapore, pp.175-194. [DOI]
- Wagh, V., Mukate, S., Panaskar, D.B., Muley, A. and Sahu, U.L., 2019. Study of groundwater hydrochemistry and drinking suitability through water quality index (WQI) modelling in Kadava river basin, India. *SN Applied Sciences*, 1(10), 1251. [DOI]
- Wintgens, T., Nätörp, A., Elango, L. and Asolekar, S.R. (eds.), 2016. *Natural Water Treatment Systems for Safe and Sustainable Water Supply in the Indian Context: Saph Pani*. IWA Publishing, London. pp.1-335. [DOI]
- Yadav, K.K., Gupta, N., Kumar, V., Choudhary, P. and Khan, S.A., 2018. GIS-based evaluation of groundwater geochemistry and statistical determination of the fate of contaminants in shallow aquifers from different functional areas of Agra city, India: levels and spatial distributions. *RSC Advances*, 8(29), pp.15876-15889. [DOI]
- Yadav, K.K., Kumar, V., Gupta, N., Kumar, S., Rezaia, S. and Singh, N., 2019. Human health risk assessment: study of a population exposed to fluoride through groundwater of Agra city, India. *Regulatory Toxicology and Pharmacology*, 106, pp.68-80. [DOI]
- Yasaswini, G., Kushala, S., Santhosh, G.S.V., Naik, M.T.K., Mondal, M., Dey, U., Das, K., Sarkar, S. and Kumar, P., 2024. Occurrence and distribution of fluoride in groundwater and drinking water vulnerability of a tropical dry region of Andhra Pradesh, India. *Water*, 16(4), 577. [DOI]
- Yenugu, S.R., Vangala, S. and Badri, S., 2020. Monitoring of groundwater quality for drinking purposes using the WQI method and its health implications around inactive mines in Vemula-Vempalli region, Kadapa District, South India. *Applied Water Science*, 10(8), 202. [DOI]