

Spatial Analysis of Environmental Vulnerability Among Tribal Households in Garbada Taluka, Dahod District, Gujarat, India

Surabhi Pareek^{ID} and Suneeta Chandorkar^{†ID}

Department of Foods and Nutrition, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat, India

[†]Corresponding author: Suneeta Chandorkar; suneetachandorkar@gmail.com

Abbreviation: Nat. Env. & Poll. Technol.
Website: www.neptjournal.com

Received: 14-10-2025

Revised: 23-12-2025

Accepted: 06-01-2026

Key Words:

Environmental vulnerability index
 Shannon index
 Crop diversity
 Sanitation
 Tribal households
 Garbada Taluka
 GIS

Citation for the Paper:

Pareek, S. and Chandorkar, S., 2026. Spatial analysis of environmental vulnerability among tribal households in Garbada Taluka, Dahod District, Gujarat, India. *Nature Environment and Pollution Technology*, 25(3), B4407. <https://doi.org/10.46488/NEPT.2026.v25i03.B4407>

Note: From 2025, the journal has adopted the use of Article IDs in citations instead of traditional consecutive page numbers. Each article is now given individual page ranges starting from page 1.



Copyright: © 2026 by the authors

Licensee: Technoscience Publications

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

Tribal households in western India are highly exposed to environmental and infrastructural challenges that influence their livelihood and food security. Assessing vulnerability at multiple scales is essential for targeted adaptation planning. This study aimed to develop a composite Environmental Vulnerability Index (EVI) for tribal households in Garbada Taluka, Dahod District, Gujarat, and examine spatial disparities at both the village and household levels. Primary data from 645 households were analyzed. Crop diversity was calculated using the Shannon Index, and a composite EVI was constructed from four indicators: landholding size, crop diversity, water sufficiency, and sanitation access. Households were classified into low-, medium-, and high-vulnerability categories. Statistical analyses (descriptive statistics, correlations, and regression) were combined with GIS-based spatial mapping to examine patterns across villages and within communities. The descriptive results showed that two-thirds of the households fell into the medium vulnerability category, with smaller proportions classified as low or high. Regression analysis confirmed that landholding size is a significant predictor of household income. Spatial mapping identified Matwa, Zari Bujarg, and Garbada as high-vulnerability clusters, whereas villages such as Jesawada, Ambli, and Devda demonstrated relatively lower vulnerability. Household-level mapping revealed intra-village disparities, with highly vulnerable households present in resilient villages. The dual-scale EVI analysis highlights village-level hotspots and household-level variations in vulnerability. These findings are relevant for climate adaptation, agricultural diversification, and WASH interventions in tribal development schemes. Future research should integrate direct nutrition indicators with environmental vulnerability frameworks to strengthen the linkages with food and nutrition security.

1. INTRODUCTION

Tribal populations in India continue to experience high levels of vulnerability due to intersecting social, economic, and environmental challenges. Limited landholdings, reliance on rainfed agriculture, and inadequate access to safe water and sanitation contribute to fragile livelihoods. These conditions create cycles of poverty and environmental stress, leaving households with fewer opportunities to adapt to changing circumstances.

The Dahod district in eastern Gujarat is one of the predominantly tribal regions of the state. The livelihood of households in this area is largely dependent on agriculture, with maize being the primary crop. However, agricultural practices are characterized by small, fragmented landholdings and limited crop diversity. At the same time, household amenities such as water supply and sanitation remain inadequate in many villages in the region. Together, these factors shape the patterns of environmental vulnerability and influence household well-being. Previous studies have highlighted persistent poverty and livelihood challenges among tribal households in Gujarat (Shah & Kumar 2008), and newer analyses continue to emphasize the structural disadvantages faced by tribal populations across India's agrarian regions (Mathew 2024, Sharma & Mohan 2024). According to the

District Census Handbook: Dahod (Census of India 2011), Garbada Taluka and other parts of the district remain heavily tribal in composition, with low literacy levels and poor household amenities. The Aspirational Districts Dashboard (NITI Aayog 2023) further indicates that Dahod continues to lag behind state averages in key indicators related to health, nutrition, and education, reflecting persistent human development challenges in tribal regions.

Recent assessments have emphasized the importance of integrated approaches to vulnerability analysis. Empirical studies from tribal regions across India demonstrate that climate variability, limited livelihood diversification, and constrained adaptive capacity jointly shape household-level vulnerabilities (Kumar et al. 2023). The Climate Vulnerability Assessment for Adaptation Planning (DST 2021) underscores the role of GIS-based methods in identifying hotspots of risk across India, while the Gujarat State Action Plan on Climate Change and Health (2022–27) (Government of Gujarat 2024) highlights Dahod as a district facing multiple stressors related to environmental and health conditions. Institutional studies also highlight the gendered and intergenerational dimensions of vulnerability. The Swaminathan Research Foundation (MSSRF 2024) reports the disproportionate impact of climate change on women and children across India's agro-ecological zones, while the Council on Energy, Environment and Water (Mohanty & Wadhawan 2021) emphasizes the role of institutional capacity and climate-resilient agriculture in enhancing adaptive responses among smallholder and tribal communities. Complementary reviews (Mohanty 2024) argue for multidisciplinary approaches that link sustainability, rural livelihoods, and tribal well-being to policy frameworks for inclusive development. Studies have shown that maize–legume intercropping systems can enhance resource-use efficiency, soil fertility, and overall sustainability in smallholder farms (Dwivedi et al. 2016).

Against this backdrop, the present study was undertaken in Garbada Taluka of Dahod district, Gujarat, to develop an Environmental Vulnerability Index (EVI) at the household level. While national- and district-level environmental vulnerability assessments exist, they largely rely on aggregated indicators that mask household-level and intra-village disparities, particularly in tribal regions. Empirical studies using primary household data combined with GIS to assess environmental vulnerability at the micro-scale remain limited. This study addresses this gap by developing a household-level Environmental Vulnerability Index (EVI) and applying a dual-scale spatial analysis to identify both village-level hotspots and intra-village heterogeneity in Garbada Taluka. Using primary survey data, the EVI was constructed from land size, crop diversity, and WASH indicators, and spatial analysis was applied to examine

household- and village-level patterns of vulnerability in the study area. By highlighting these patterns, this study aims to provide an evidence base for local planning and strengthen resilience among tribal households.

2. MATERIALS AND METHODS

2.1. Study Area

The research was conducted in Garbada Taluka of Dahod District, Gujarat, India, one of the nine talukas of Dahod. The district has a predominantly tribal population and has been recognized as an Aspirational District by the Government of India due to persistent developmental challenges in education, health, agriculture, and infrastructure (NITI Aayog 2023). Garbada was selected as the study site because of its diverse topography, comprising hilly, plain, and medium regions, which provides a natural setting to examine geographical variations in agricultural practices and household vulnerability. Agriculture, particularly maize cultivation, is the primary livelihood in this region. The study was conducted between February 2023 and July 2023.

2.2. Study Design and Samples

A cross-sectional household survey was conducted in 645 households in Garbada Taluka. Households were selected using random GPS points generated in ArcGIS 10.5, enforcing a minimum of 50 m spacing to reduce clustering and spatial bias. This spatial randomization approach followed the GIS-based household selection method described by Mathenge et al. (2023). Random points were overlaid on high-resolution satellite imagery and village boundary layers and matched to actual household locations during field visits using GPS-enabled Android devices. This procedure ensured uniform geographic coverage and minimized the enumerator bias.

2.3. Data Collection

Primary data were collected through structured interviews and household-level observations using pretested questionnaires. The tool includes the following modules:

1. **Household characteristics:** family size, number of children, annual income, and landholding (acres).
2. **Agriculture:** production data for maize, rice, wheat, and soybeans.
3. **Water and sanitation:** household access to a sufficient water supply and toilet facilities (MSSRF 2024).
4. **Geolocation:** latitude and longitude of each household recorded using handheld GPS devices.

Ethical clearance was obtained from the Department of

Foods and Nutrition, Faculty of Family and Community Sciences, The Maharaja Sayajirao University of Baroda.

2.4. Variables and Indicators

2.4.1. Crop Diversity

Crop diversification was measured using the Shannon (H') and Simpson diversity indices (D) (Magurran 2021). Both indices were computed from the production data of maize, rice, wheat, and soybean. While both indices were examined descriptively, the Shannon index was chosen for regression and EVI construction because it captures both crop richness and evenness in diversified farming systems (Sibhatu et al. 2015).

2.4.2. WASH Indicators

Water sufficiency (yes/no) and toilet access (yes/no) were included as essential indicators of environmental health (DST 2021).

2.4.3. Household Income

Annual self-reported earnings were recorded and validated through cross-verification of agricultural output and market prices (Shukla et al. 2025).

2.4.4. Landholding Size

The total cultivated land (in acres) was measured and used to classify households into small, medium, and large categories, following the 33rd percentile cut-offs (Government of Gujarat 2024).

2.5. Environmental Vulnerability Index (EVI)

An Environmental Vulnerability Index (EVI) was constructed at the household level using four indicators: landholding size, crop diversity, water sufficiency, and toilet facilities. Each variable was coded as binary (0 = not vulnerable, 1 = vulnerable)

$$EVI_i = \sum_{j=1}^4 X_{ij}$$

Where:

- $X_{i1} = 1$ if landholding $\leq$ 33rd percentile (small landholding), else 0
- $X_{i2} = 1$ if Shannon diversity $\leq$ 33rd percentile (low crop diversity), else 0
- $X_{i3} = 1$ if the household lacked sufficient water, else 0
- $X_{i4} = 1$ if the household lacked a toilet facility, else 0

Thus, EVI_i ranged from 0 (no vulnerability) to 4 (highest vulnerability).

Households were categorized as:

1. Low vulnerability: scores 0–1
2. Medium vulnerability: scores 2–3
3. High vulnerability: score 4

This composite index approach aligns with contemporary frameworks that integrate multiple environmental and socioeconomic stressors into a single standardized measure of vulnerability. Similar additive or weighted-index methods have recently been applied in social and ecological contexts for micro-level vulnerability assessments, adapted here to the household scale (Smits & Huisman 2024). Equal weighting was adopted to maintain transparency and reduce subjectivity in the absence of empirically derived weights, a common approach in exploratory household-level vulnerability assessments. While binary coding improves interpretability and comparability across indicators, it may result in some loss of information from the underlying continuous variables, which is acknowledged as a limitation of the present study.

For village-level thematic maps, continuous indicator values were classified using percentile-based thresholds, where values $\leq$ 33rd percentile represented low, 34–66th percentile represented medium, and $\geq$ 67th percentile represented high levels of vulnerability or deprivation.

2.6. Data Analysis

2.6.1. Statistical Analysis

All quantitative analyses were performed using R software (version 4.3.3). Descriptive statistics (means, medians, and percentages) were computed to summarize household and village-level indicators. Spearman's rank correlation was used to examine the associations among key variables, including landholding size, crop diversity, water sufficiency, toilet access, and household income. To identify the determinants of household income, a linear regression model was estimated with log-transformed income as the dependent variable. Household- and village-level Environmental Vulnerability Index (EVI) scores were summarized to highlight vulnerability patterns across the study area. The combined use of R for statistical analysis and ArcGIS 10.5 for spatial visualization ensured robust analytical rigor and reproducibility of the results.

2.6.2. Spatial Analysis

Household coordinates and village boundary shapefiles were imported into ArcGIS 10.5 for spatial visualization. Five thematic maps were generated to illustrate spatial variability in key indicators.

1. Household-level EVI classification
2. Village-level mean EVI
3. Village-level crop diversity (Shannon index)
4. Village-level percentage of households with sufficient water

5. Village-level percentage of households with toilet facilities

3. RESULTS AND DISCUSSION

The findings of the study are presented at both the household and village levels to capture the individual and spatial patterns of vulnerability. The results include descriptive statistics of households, correlation and regression analyses to assess key determinants of income and vulnerability, and the construction of an Environmental Vulnerability Index (EVI). Spatial analysis using ArcGIS further illustrated inter-village variations in crop diversity, water sufficiency, sanitation, and EVI distribution. Together, these findings provide a comprehensive picture of the environmental vulnerability of tribal households in Garbada Taluka.

The results are presented below, with an interpretation provided alongside to place the findings in the context of the existing literature.

3.1. Household Characteristics

The surveyed households had an average landholding size of 3.25 acres, with a median landholding size of 4 acres (Table 1). The average household size was 6.2 members, with a median of 2 children. The average annual household income was ₹159,855, although the range was wide (₹10,000–700,000), reflecting the heterogeneity in livelihoods. Approximately 68% of households reported sufficient water availability, whereas 62% had toilet facilities. Crop diversity was modest, with a mean Shannon index of 1.12 and a Simpson index of 0.62, confirming the dominance of maize. Such dependence on maize as a staple crop under rainfall variability has been previously documented in Dahod (Chawra et al. 2024). Shah and Kumar (2008) also emphasized poverty and livelihood vulnerability tied to land access among Gujarat's tribal communities.

While crop diversity reflects agricultural resilience, its relationship with household resources and infrastructure was further explored through correlation and regression analysis.

Table 1: Descriptive statistics of surveyed households (n=645).

Variable	Mean/%	Median	Min–Max
Landholding size [acres]	3.25	4.0	0-15
Household size	6.2	7.0	2-15
Children in the household	2.1	2.0	0-6
Annual income [₹]	159,855	140,000	10,000-700,000
% with sufficient water	68%	-	-
% with toilet facility	62%	-	-
Shannon diversity (mean)	1.12	-	0-1.39
Simpson diversity (mean)	0.62	-	0-0.70

Spearman's correlation analysis (Table 2) showed a weak but significant positive relationship between landholding and income ($\rho = 0.079$, $p = 0.046$), suggesting that larger farms provide an income advantage. A significant correlation was also observed between water sufficiency and toilet access ($\rho = 0.086$, $p = 0.030$), reflecting the link between household infrastructure and sanitation. However, crop diversity was not significantly associated with income or water sufficiency, highlighting the persistence of monocropping of maize. These findings are consistent with national assessments that highlight landholding as a stronger determinant of household resilience compared to diversification (DST 2021, Smits & Huisman 2024).

3.2 Determinants of Household Income

Regression analysis confirmed that landholding size was the only significant predictor of household income ($\beta = 0.032$, $p = 0.028$) after adjusting for crop diversity, water sufficiency, and sanitation (Table 3). The adjusted R^2 was only 0.009, indicating that agricultural and infrastructure variables explained less than 1% of the income variation. This highlights the complexity of tribal livelihoods, where income depends on multiple non-farm and environmental factors. A similar weak explanatory power of agricultural determinants has been observed in Central India's vulnerability studies (Smits & Huisman 2024) and in Gujarat's State Action Plan on Climate Change and Health (Government of Gujarat 2024). The very low adjusted R^2 (0.009) suggests that agricultural and WASH-related variables alone are insufficient to explain household income variation in tribal settings. In districts such as Dahod, household incomes are widely reported to depend on diversified livelihood strategies, including non-farm employment and seasonal migration, which are beyond the scope of the present study. Therefore, the regression results should be interpreted as indicative associations rather than strong predictors of income, and future research may benefit from incorporating migration, wage labor, and social protection variables. Although crop diversity is widely linked to resilience and nutrition outcomes in the literature, it did not show a statistically significant association with household income in this analysis.

Table 2: Spearman correlations between key indicators.

Variables	rho [ρ]	p-value
Landholding vs Income	0.079	0.046*
Crop diversity vs Income	0.069	0.079
Crop diversity vs Water	- 0.024	0.548
Landholding vs Water	- 0.019	0.639
Toilet vs Water	0.086	0.030*

* $p < 0.05$, ns = not significant

Table 3: Regression results: Determinants of log household income.

Predictor	Estimate	Std. Error	t-value	p-value
Intercept	11.616	0.080	145.6	<0.001
Landholding (acres)	0.032	0.014	2.20	0.028 *
Crop diversity (Shannon)	0.052	0.071	0.73	0.465
Water sufficiency	0.061	0.046	1.34	0.181
Toilet facility	0.043	0.046	0.94	0.346

*Model fit: Adjusted $R^2 = 0.009$, $F(4, 640) = 2.43$, $p = 0.046$ ($p < 0.05$)

3.3 Spatial Analysis of Vulnerability

To complement the statistical findings, a spatial analysis was conducted to capture the geographic variation in agricultural practices, WASH infrastructure, and environmental vulnerability across Garbada Taluka. Mapping key indicators at both the village and household levels helps visualize disparities that may not be fully apparent in tabular summaries. This approach highlights clusters of high vulnerability and provides a basis for identifying priority intervention areas. Figs. 1–5 present the spatial distributions of crop diversity, sanitation, water sufficiency, and the composite Environmental Vulnerability Index (EVI).

3.3.1 Crop Diversity and Environmental Indicator

Crop diversity, measured using the Shannon Index, was lowest in Matwa, Zari Bujarg, and Garbada (red), reflecting maize-dominated monocropping, while Ambli, Jesawada, and Devda showed relatively higher diversity (green) (Fig. 1). Low Shannon scores indicate limited crop diversification, which is associated with higher vulnerability to climatic shocks and lower dietary diversity, which is essential for child health. Similar rainfall-driven monocropping patterns have been observed in Dahod (Chawra et al. 2024). At the nutrition level, studies show that agricultural diversification, as captured by the Shannon Index, is positively associated with household dietary adequacy (Smits & Huisman 2024). The map illustrates the spatial distribution of crop diversity across the villages. Villages shaded red represent low crop diversity (dominated by maize monocropping), yellow indicates medium diversity, and green denotes high diversity with a more balanced cultivation of cereals, pulses, and oilseeds. High-diversity villages (e.g., Ambli, Jesawada, and Devda) reflect more resilient farming systems, whereas low-diversity clusters (e.g., Matwa, Zari Bujarg, and Garbada) highlight areas vulnerable to climatic shocks and dietary monotony.

Sanitation access was poorest in Matwa, Vajelav, and Zari Bujarg, where fewer than 40% of households reported a toilet, whereas Jesawada and Bharsada performed better (Fig. 2). Poor sanitation is a well-documented determinant of

undernutrition in children. NITI Aayog (2023) and Pickering et al. (2019) reported that toilet access reduces the odds of stunting by 16–39%. This map presents the spatial distribution of sanitation coverage across the villages. Villages shaded red represent very low coverage, orange indicates low coverage, yellow indicates medium coverage, and green indicates high coverage. Very low coverage is concentrated in Matwa, Vajelav, and Zari Bujarg, whereas comparatively better access is observed in Jesawada, Bharsada, and Ambli. The clustering of red and orange zones across central and southern Garbada indicates critical sanitation deficits, which are strongly linked to child undernutrition through higher exposure to diarrheal and parasitic infections.

Water sufficiency was particularly low in Garbada, Zari Bujarg, and Matwa (<40%), whereas Patiya, Bhe, and Devda demonstrated better access (Fig. 3). These results align with evidence from the Maternal & Child Nutrition showing that children in households with sufficient water were significantly more likely to meet the minimum dietary diversity standards (Gao et al. 2022). This map shows the distribution of household water sufficiency across the villages. Villages shaded red, orange, yellow, and green represent very low, low, medium, and high access, respectively. Very low sufficiency is concentrated in Matwa, Zari Bujarg, Garbada, and Boriyala, whereas Patiya, Bhe, Jesawada, and Devda exhibit higher sufficiency. The clustering of water-deficit villages in the central and southern parts of Garbada reflects the underlying infrastructure and resource constraints. Research from rural India, including Uttarakhand, also highlights that improved water and sanitation access reduces stunting by lowering exposure to diarrheal and parasitic infections (Nair et al. 2017).

To integrate these multiple dimensions—land, diversity, water, and sanitation—into a single measure, a composite Environmental Vulnerability Index (EVI) was constructed.

3.3.2 Environmental Vulnerability Index (EVI)

The household-level EVI revealed that 53.8% of households were classified as medium-vulnerability, 41.4% as low-vulnerability, and 4.8% as high-vulnerability (Table 4). The dominance of medium vulnerability suggests that while extreme conditions are rare, most households face sustained risks owing to small landholdings, low crop diversity, and limited WASH infrastructure. These results align with the national Climate Vulnerability Assessment (DST 2021), which found medium vulnerability to be prevalent across large parts of India, and with hotspot analyses of regional disparities (Ravichandran et al. 2023).

While the EVI summary provides a quantitative distribution of vulnerability, spatial mapping allows these patterns to be visualized across villages.




Fig. 1: Village-wise mean crop diversity (Shannon Index) in Garbada Taluka.




Fig. 2: Village-wise percentage of households with toilet facilities in Garbada Taluka.




Fig. 3: Village-wise percentage of households with sufficient water in Garbada Taluka.

3.3.3. Village-Level Spatial Patterns

Figures 1–4 provide an overview of the village-level environmental indicators across Garbada Taluka. These maps highlight variations in crop diversity, sanitation, water sufficiency, and the aggregated Environmental Vulnerability Index (EVI). Together, they illustrate the uneven distribution of resources and resilience across villages. For instance, villages such as Matwa, Zari Bujarg, Garbada, and Simaliya Bujarg consistently emerged as high-vulnerability clusters (Fig. 4), marked by low crop diversity, poor sanitation, and inadequate water sufficiency. In contrast, villages such as Jesawada, Ambli, and Devda displayed relatively higher resilience, with more diversified production systems and better infrastructure. These spatial patterns are consistent with broader state-level assessments, such as the Government of Gujarat (2024), which identified the Dahod district as a climate- and environment-sensitive region.

Table 4: Distribution of Environmental Vulnerability Index (EVI) classes.

EVI Class	Criteria (score)	No. of households
Low	0–1	267
Medium	2–3	347
High	4	31

Although village-level patterns highlight clusters of vulnerability, households within the same village often differ substantially from each other. Household-level mapping helps to capture these intra-village disparities.

3.3.4. Household-Level Spatial Patterns

At the micro scale, Fig. 5 shows household-level EVI. Even within medium-vulnerability villages, such as Sahada and Pandadi, households were heterogeneous—some were resilient, whereas others were highly vulnerable. This intra-village disparity is often masked in aggregated data and underscores the value of household-scale mapping (Raj & Sofi 2023).

Comparing village- and household-level results helps assess whether vulnerability patterns are consistent across scales or if finer variations are masked in aggregated data. The household-level EVI provides a quantitative basis for capturing intra-village variability, as individual households are assigned numeric vulnerability scores ranging from 0 to 4. The observed co-existence of highly vulnerable and relatively resilient households within the same village reflects the uneven access to land, water, sanitation, and agricultural resources at the household scale. Such intra-village heterogeneity highlights the limitations of

relying solely on village-level averages and reinforces the importance of household-scale analysis for targeted planning and intervention.

3.3.5. Consistency Across Scales

A comparison of village-level (Fig 4) and household-level (Fig. 5) The Environmental Vulnerability Index (EVI) demonstrates both consistency and complementarity in the findings. At the aggregate scale, villages such as Matwa, Zari Bujarg, Garbada, and Vajelav were consistently classified as being highly vulnerable. This pattern was reinforced at the micro-scale, where a large share of households in these villages was mapped under the high vulnerability category. Simultaneously, household-level mapping revealed intra-village disparities that were not visible in the aggregate analysis, with even resilient villages such as Jesawada and Ambli containing a few high-vulnerability households.

These dual-scale findings align with earlier research on environmental conditions and health outcomes. For instance, household environmental conditions have been shown to directly affect morbidity in India (Brahmanandam & Nagarajan 2021), while behavioral interventions such as

waste segregation in Delhi demonstrate how socioeconomic factors shape vulnerability at the micro level (Kaur & Kaur 2024). These insights reinforce the importance of combining village- and household-level analyses, as undertaken in this study. This map shows the average environmental vulnerability of households, aggregated at the village level. Villages shaded green represent low vulnerability, yellow indicates medium vulnerability, and red indicates high vulnerability. High EVI values are concentrated in Zari Bujarg, Garbada, Simaliya Bujarg, and Nandhelav, while Jesawada, Ambli, Devda, and Gulbar show relatively lower vulnerability. The map highlights the regional clustering of vulnerability, with southern and central villages facing greater environmental stress than western and northern areas.

At the district and state levels, composite indices have been widely applied for climate vulnerability assessments. The Council on Energy, Environment, and Water (Shukla et al. 2025) produced a district-level vulnerability atlas of India, and the Department of Science and Technology's Climate Vulnerability Assessment for Adaptation Planning in India (DST 2021) combined environmental, economic, and social indicators to guide adaptation strategies. The clustering of




Fig. 4: Village-wise mean Environmental Vulnerability Index (EVI) in Garbada Taluka.

high-vulnerability villages in Garbada corresponds to similar spatial disparities documented in western and central India.

At the regional scale, studies using the Climate Vulnerability Index have highlighted significant variations in exposure and adaptive capacity across Indian states (Singh et al. 2021). Evidence from tribal areas in Central India shows that livelihood vulnerability is strongly mediated by crop diversity, landholding, and water access (Smits & Huisman 2024), whereas household-level vulnerability in the Western Ghats varies sharply with ecological and demographic factors (Raj & Sofi 2023). These parallels reinforce the present study's observation that intra-village heterogeneity in Garbada is as critical as inter-village differences (Fig. 5).

This map illustrates the distribution of environmental vulnerability at the household level. Households shaded

green, yellow, and red represent low, moderate, and high vulnerability, respectively. High-vulnerability households are scattered across villages such as Matwa, Garbada, and Zari Bujarg, whereas most households in Jesawada, Ambli, and Devda show low vulnerability. The household-level analysis reveals intra-village disparities, where even villages with medium overall vulnerability include both resilient and highly vulnerable households.

The integration of spatial mapping further strengthens the explanatory power of household survey data. Remote sensing-based eco-environmental assessments have demonstrated the value of combining climatic and anthropogenic indicators for dynamic vulnerability mapping (Lan et al. 2023). Although this study primarily relied on household survey data integrated with GIS, the consistency of its results with both composite indices and remote sensing




Fig. 5: Household-level Environmental Vulnerability Index (EVI) in Garbada Taluka.

approaches underscores their methodological robustness. The findings were subsequently interpreted in relation to the broader concerns of food and nutrition security, given the close link between environmental vulnerability, crop diversity, and household well-being. Similar GIS-based spatial studies have revealed how flood and drought risks influence food security in vulnerable regions (Rosalia et al. 2021). Given the cross-sectional design of this study, the observed relationships should be interpreted as associations rather than causal effects. The spatial analysis presented in this study is primarily descriptive, and formal spatial autocorrelation statistics (e.g., Moran's I or Getis-Ord G_i^*) were not applied; therefore, spatial clustering patterns should be interpreted with caution.

3.4. Implications for Food and Nutrition Security

Although direct nutrition indicators were not collected, the environmental variables used in the EVI are well-established determinants of food and nutrition. Crop diversity has been linked to dietary adequacy (Sibhatu et al. 2015), and access to water and sanitation has been associated with reduced child stunting and underweight (Nair et al. 2017, Pickering et al. 2019). The clustering of vulnerable villages in Garbada, therefore, suggests a potential overlap with nutritional insecurity, consistent with national studies linking climate vulnerability and child malnutrition (Gao et al. 2022, NITI Aayog 2023). However, a key limitation of this study is the absence of direct nutritional indicators, such as dietary diversity or anthropometric measures; therefore, links to food and nutrition security were interpreted contextually based on the existing literature rather than measured outcomes.

4. CONCLUSIONS

This study assessed environmental vulnerability among 645 tribal households in Garbada Taluka of Dahod District, Gujarat, using a composite Environmental Vulnerability Index (EVI) that incorporates landholding, crop diversity, water sufficiency, and sanitation access. The results demonstrated that most households fell into the medium vulnerability category, with villages such as Matwa, Zari Bujarg, and Garbada consistently emerging as hotspots of high vulnerability. Household-level mapping further revealed significant intra-village disparities, underscoring the importance of considering both aggregate and micro-level scales in vulnerability assessments.

Although nutritional outcomes were not directly measured, the environmental and infrastructural indicators used in the EVI are well-established determinants in the literature on food and nutrition security. The clustering of low crop diversity, poor sanitation, and inadequate water

sufficiency suggests potential heightened risks to livelihood and nutrition security in the study area. Evidence from India and other smallholder contexts shows that higher crop diversity is strongly associated with improved household dietary adequacy (Sibhatu et al. 2015).

From a policy perspective, these findings have implications for targeted and location-specific interventions. Villages with high EVI may warrant priority attention in climate adaptation planning, agricultural diversification programs, and WASH (water, sanitation, and hygiene) infrastructure development. Simultaneously, household-level disparities suggest the relevance of micro-level interventions, ensuring that vulnerable families within relatively resilient villages are not overlooked. Access to safe water and toilet facilities has been consistently linked with reduced risks of stunting and underweight among children by lowering exposure to diarrheal and parasitic infections, while food-based interventions that improve diet diversity have also proven effective in addressing multiple micronutrient deficiencies (Nair et al. 2017).

The dual-scale approach demonstrated here can support tribal development schemes, the State Action Plan on Climate Change, and national programs such as ICDS and POSHAN 2.0 by identifying critical hotspots and informing more targeted interventions. Future research should integrate direct nutrition indicators, such as dietary diversity and child growth outcomes, with environmental vulnerability frameworks to provide a more holistic understanding of food and nutrition security.

5. ACKNOWLEDGMENTS

The authors thank all the participants in this study.

6. REFERENCES

- Brahmanandam, N. and Nagarajan, R., 2021. Impact of change in household environment condition on morbidity in India: Evidence from longitudinal data. *PLOS ONE*, 16(3), p.e0247465. DOI
- Census of India, 2011. District Census Handbook: Dahod. Office of the Registrar General & Census Commissioner, Government of India. Retrieved October 31, 2025, from <https://www.census2011.co.in/census/district/199-dohad.html>
- Chawra, U., Kacha, H., Rani, R.R. and Bhabhor, G., 2024. Rainfall variability and crop planning of Dahod: A case study from a tribal district in Gujarat, India. *International Journal of Environment and Climate Change*, 14(4), pp.259-266. DOI
- Department of Science and Technology (DST), 2021. Climate Vulnerability Assessment for Adaptation Planning in India Using a Common Framework. Government of India. Retrieved October 31, 2025, from <https://dst.gov.in/document/reports/national-climate-vulnerability-assessment-report>
- Dwivedi, A., Singh, A., Tomar, S.S., Bankoti, P., Singh, O. and Kumar, K., 2016. Interaction effect of maize and mash-bean intercropping on sustainable production systems in the subtropical zone of India. *Nature*

- Environment and Pollution Technology*, 15(2), pp.661-665. Retrieved October 31, 2025, from [https://neptjournal.com/upload-images/NL-56-46-\(46\)B-3252.pdf](https://neptjournal.com/upload-images/NL-56-46-(46)B-3252.pdf)
- Gao, Y., Sheng, J., Mi, X., Zhou, M., Zou, S. and Zhou, H., 2022. Household water access, dietary diversity and nutritional status among preschoolers in poor rural areas of Central and Western China. *Nutrients*, 14(3), p.458. DOI
- Government of Gujarat, 2024. State Action Plan on Climate Change and Human Health – Gujarat (2022–27). Gandhinagar: Government of Gujarat & NCDC. Retrieved October 31, 2025, from https://ncdc.mohfw.gov.in/wp-content/uploads/2025/01/09_SAPCCHH_Gujarat_21-10-24.pdf
- Kaur, H. and Kaur, P., 2024. Factors determining household waste segregation behavior: An Indian case study. *International Journal of Experimental Research and Review*, 41(Special Volume), pp.83-95. DOI
- Kumar, P., Ashwani, Mishra, S., Thakur, S., Kumar, D. and Raman, V.A.V., 2023. Vulnerability of tribal communities to climate variability in Lahaul and Spiti, Himachal Pradesh, India. *Cografya Dergisi*, 47, pp.29-43. DOI
- Lan, G., Jiang, X., Xu, D., Guo, X., Wu, Y., Liu, Y. and Yang, Y., 2023. Ecological vulnerability assessment based on remote sensing ecological index (RSEI): A case of Zhongxian County, Chongqing. *Frontiers in Environmental Science*, 10, p.1074376. DOI
- Magurran, A.E., 2021. Measuring biological diversity. *Current Biology*, 31(19), pp.R1174-R1177. DOI
- Mathenge, M., Sonneveld, B.G.J.S. and Broerse, J.E.W., 2023. Mapping the spatial dimension of food insecurity using GIS-based indicators: A case of Western Kenya. *Food Security*, 15(2), pp.243-260. DOI
- Mathew, V.J., 2024. The evolution and status of Tribal Sub Plan through local self-government institutions: Reflections from Kerala's experience. ISEC Working Paper No. 583. Institute for Social and Economic Change. Retrieved October 31, 2025, from <https://www.isec.ac.in/wp-content/uploads/2025/04/WP-583-Vidya-J-Mathew-Final.pdf>
- M.S. Swaminathan Research Foundation (MSSRF), 2024. How Does Climate Change Impact Women and Children Across Agro-Ecological Zones in India: A Scoping Study. Chennai: MSSRF. Retrieved October 31, 2025, from https://www.mssrf.org/sites/default/files/2025-05/Report-CCW-Study-v2-a_0.pdf
- Mohanty, A. and Wadhawan, S., 2021. Mapping India's Climate Vulnerability: A District-Level Assessment. New Delhi: CEEW. Retrieved October 31, 2025, from <https://www.ceew.in/sites/default/files/ceew-study-on-climate-change-vulnerability-index-and-district-level-risk-assessment.pdf>
- Mohanty, B., 2024. A multidisciplinary review of emerging research on tribal communities, sustainable agriculture and rural livelihoods. *SSRN*, 9, p.76142. [DOI]
- Nair, M.K., Augustine, L.F. and Konapur, A., 2017. Food-based interventions to modify diet quality and diversity to address multiple micronutrient deficiency. *Frontiers in Public Health*, 3, p.277. DOI
- NITI Aayog, 2023. Deep-Dive: Insights from Champions of Change – The Aspirational Districts Dashboard. Government of India. Retrieved October 31, 2025, from <https://www.niti.gov.in/sites/default/files/2023-03/DEEP-DIVE-Insights-from-Champions-of-Change-The-Aspirational-Districts-Dashboard.pdf>
- Pickering, A.J., Null, C., Winch, P.J., Mangwadu, G., Arnold, B.F., Prendergast, A.J., Njenga, S.M., Rahman, M., Ntozini, R., Benjamin-Chung, J. et al., 2019. The WASH Benefits and SHINE trials: Interpretation of WASH intervention effects on linear growth and diarrhoea. *The Lancet Global Health*, 7(8), pp.e1139-e1146. DOI
- Raj, R. and Sofi, A.A., 2023. Does climate change lead to severe household-level vulnerability? Evidence from the Western Ghats of Kerala, India. *Land Use Policy*, 130, p.106655. DOI
- Ravichandran, V., Kantamaneni, K., Singh, A., Nair, A., Janakiraman, A. and Prem Kumar, S., 2023. Mapping hotspot clustering: An approach to study the spatial pattern of quality of living households in India. *Remote Sensing Applications: Society and Environment*, 31, p.101007. DOI
- Rosalia, A.C.T., Suryanto and Hakim, L., 2021. Spatial analysis of the impact of flood and drought on food security index. *Nature Environment and Pollution Technology*, 20(2), pp.721-727. DOI
- Shah, A. and Kumar, A., 2008. The Grassroots of Poverty: Tribal Livelihoods and Development in Gujarat. GIDR Working Paper No. 187. Gujarat Institute of Development Research. Retrieved October 31, 2025, from <https://gidr.ac.in/pdf/WP-187.pdf>
- Sharma, P. and Mohan, K., 2024. Location, Livelihoods and Tribal People: Exploring the Linkages in India. Independent Research Publication. Retrieved October 31, 2025, from https://www.researchgate.net/publication/383238589_Location_Livelihoods_and_Tribal_People_Exploring_the_Linkages_in_India
- Shukla, S., Abraham, S. and Parekh, S., 2025. Advancing Climate-Resilient Agriculture in India by Strengthening Institutional Capacity: Lessons from Bihar and Odisha. New Delhi: CEEW. Retrieved October 31, 2025, from <https://www.ceew.in/sites/default/files/how-can-india-scale-climate-resilient-agriculture-practices-and-climate-smart-farming.pdf>
- Sibhatu, K.T., Krishna, V.V. and Qaim, M., 2015. Production diversity and dietary diversity in smallholder farm households. *Proceedings of the National Academy of Sciences (PNAS)*, 112(34), pp.10657-10662. DOI
- Singh, C., Madhavan, M. and Arvind, J., 2021. Climate change adaptation in Indian cities: A review of existing actions and spaces for triple wins. *Urban Climate*, 36, p.100783. DOI
- Smits, J. and Huisman, J., 2024. The GDL Vulnerability Index (GVI). *Social Indicators Research*, 174, pp.721-741. DOI