

Vol. 24

Open Access Journal

D1725

Original Research Paper

https://doi.org/10.46488/NEPT.2025.v24i02.D1725

Thermodynamic Modeling Studies on Biosorption of Reactive Amoxicillin Antibiotic by Pithophora Macroalgae in Aqueous Solution

Murad M. Khamayseh¹[†]^(D) and Rana Kidak²^(D)

¹Department of Environmental Sciences, Cyprus International University, Nicosia 99010, Turkey ²Department of Environmental Engineering, Faculty of Engineering, Cyprus International University, Nicosia 99010, Turkey [†]Corresponding author: Murad M. Khamayseh; murad.kha32@gmail.com

Abbreviation: Nat. Env. & Poll. Technol. Website: www.neptjournal.com

Received: 05-09-2024 Revised: 16-11-2024 Accepted: 20-11-2024

Key Words:

Amoxicillin **Biosorption** Biosorbent Pithophora Macroalgae

Citation for the Paper:

Khamayseh, M. M. and Kidak, R., 2025. Thermodynamic modeling studies on biosorption of reactive amoxicillin antibiotic by Pithophora macroalgae in aqueous solution. Nature Environment and Pollution Technology, 24(2), p. D1725. https://doi. org/10.46488/NEPT.2025.v24i02.D1725

Note: From year 2025, the journal uses Article ID instead of page numbers in citation of the published articles.



Copyright: © 2025 by the authors Licensee: Technoscience Publications This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/4.0/).

ABSTRACT

Antibiotic removal poses a serious risk to the environment due to its intricate structure. Consequently, scientists are developing new and efficient techniques to remove antibiotic compounds from wastewater. The goal of this study is to employ green Pithophora macroalgae to remove the antibiotic amoxicillin (AMX) from a water-based solution. With a focus on understanding the process, this study assesses the application of reacting AMX biosorption on the biomass of Pithophora algae in aqueous solutions using thermodynamic modeling. The determined thermodynamic characteristics show that an endothermic process is used in the biosorption of the antibiotic AMX, considering that AMX has a positive electrical charge of ΔH° at 49.796 KJ.moL⁻¹. As ΔG° has a positive charge (2.982 kJ.moL⁻¹, 3.718 kJ.moL⁻¹, and 4.793 kJ.moL⁻¹) for AMX at (298 K, 303 K, and 308 K, respectively. This positive result indicates that the reaction is not feasible or spontaneous. The decrease in chaos at the liquid/solid interface caused by AMX biosorption on Pithophora macro algae is reflected in the negative charge of ΔS° , which was -176.735 kJ.moL⁻¹. The effect of temperature on the biosorption of AMX was investigated for different initial AMX concentrations. At a lower temperature of 298 K, the AMX molecules were more likely to diffuse into the internal pores of the Pithophora algae. This suggests that the diffusion rate of the adsorbate (AMX) across the bulk and pore boundaries of the biosorbent particles may be increased at lower temperatures. The findings of this study indicate that the biomass of the macroalgae Pithophora is a valuable biosorbent for the biosorption of AMX antibiotics, and it may have potential applications in the treatment of wastewater.

INTRODUCTION

Biosorption is a sustainable method for removing contaminants and heavy metals from aqueous solutions (Balarak et al. 2017). It is a process where contaminants are extracted from water or wastewater using a biosorbent. Biosorbents are materials that can be found naturally or produced artificially, and they can selectively bind contaminants and other chemicals from aqueous solutions (Arihilam & Arihilam, 2019). They are frequently employed in the cleanup of polluted soils and water. Because of its straightforward operation, low cost of raw materials, ability to reduce spent water, and ability to keep contaminants out of natural resources, biosorption has become a viable method for removing toxins from water and soil (Abd & Mohammed-Ridha 2021).

Amoxicillin (AMX) is a beta-lactam antibiotic that is used for the treatment of bacterial infections. Because amoxicillin is so inexpensive, it is prescribed extensively throughout the world. With a half-life of roughly one to two hours, it is known that this antibiotic is very vulnerable to hydrolytic and oxidative degradation in aqueous solutions (Aljeboree et al. 2020). It is well known that this antibiotic is extremely reactive, acting much like hydroxyl radicals. Due to its reactivity, using it in water systems may cause the creation of hazardous intermediates, which may have an impact on aquatic life (Matsubara et al. 2020).

One potential method for removing AMX from water systems is biosorption. The majority of AMX is allegedly released into the sewage system unaltered, as stated by Jung et al. (2012). Water samples contain about 1.3 μ g.mL⁻¹ of it (Välitalo et al. 2017). Before AMX is released into the environment, the sewage system does not adequately remove it (Lai et al. 2017). On the other hand, amoxicillin can be removed more successfully via genetic engineering, and the biological process of removing antibiotics from macroalgae does not require chemical intervention (Chakhtouna et al. 2021). Many methods, including adsorption, coagulation, electrolysis, filtration, photocatalysis, coagulation, accelerated oxidation, and biological degradation, have been used to eradicate antibiotics (Yan et al. 2022).

The high energy consumption, high material costs, and additional pollution caused by the addition of additional chemicals are some of the disadvantages of these technologies (Boukhelkhal et al. 2016). Adsorption is the most widely used and successful technique for removing contaminants from wastewater (Asim et al. 2021). Its uncomplicated appearance and ease of usage make it simple to install. Because biosorption is inexpensive, efficient, and safe for the environment, it has attracted a lot of attention as an alternative to antibiotics. This method binds and concentrates pollutants in water by using biological material from live and dead biomass that has been heat-dried and chemically treated (Dimitrakopoulou et al. 2012). Because of its straightforward operation, low cost of raw materials, ability to minimize spent water, and ability to keep contaminants out of natural resources, biosorption has become a promising method for removing pollutants from soil and water (Limousy et al. 2017). With a focus on comprehending its mechanism, this paper reviews some recent developments in equilibrium and kinetics research on the biosorption of reactive Amoxicillin Antibiotic (AMX) on Pithophora Macroalgae in aqueous solutions. Pithophora is a type of filamentous green algae. It can grow in dense mats either on the surface or at the bottom. This algae is often described as having a rough, coarse texture, and its growth pattern resembles a tangled mass of cotton or wool. One of the most economical adsorbents is green algae, which contains cellulosic polysaccharides that react with a large number of proteins to form glycoproteins. Numerous functionalized groups, such as amino, carboxyl, sulfate, and hydroxyl, are present in these materials and are essential to the biosorption process (Romera et al. 2007). The thermodynamic values obtained from the study suggest that the sorption of levofloxacin (LVX) antibiotics is governed by an endothermic process. The findings indicate that the biomass of Pithophora macroalgae

serves as an effective biosorbent for the removal of LVX antibiotics, presenting a promising alternative method for the elimination of antibiotics from aqueous environments. Furthermore, Pithophora demonstrates potential applicability in real-world wastewater treatment scenarios, having proven its efficacy as a biosorbent for antibiotic compounds (Khamayseh & Kidak 2023). According to the study conducted by Kalyani et al. (2021), the thermodynamic analysis revealed that the biosorption processes are both exothermic and spontaneous. The biosorbents derived from Pithophora cleveana Wittrock and Mimusops elengi exhibited maximum removal efficiencies of 74.11% and 73.11%, respectively, with corresponding maximum uptake capacities of 13.58 mg.g⁻¹ and 12.96 mg.g⁻¹. Furthermore, the thermodynamic parameters, including ΔG° , ΔH° , and ΔS° , indicated that the biosorption process is spontaneous and exothermic. These findings suggest that the biosorbents derived from Pithophora cleveana Wittrock and Mimusops elengi possess significant potential for the removal of lead metal ions from aqueous solutions, attributable to their effective absorption capabilities (Gaddam et al. 2020). Additionally, Pithophora has demonstrated its applicability in real-world wastewater treatment scenarios and has proven to be an effective biosorbent for the biosorption of antibiotics.

Algal biomass treated with an alkaline solution will bio-absorb pharmaceutical chemicals. The primary goal of this research is to use the readily available green Pithophora macroalgae to extract AMX from aqueous solutions. The starting pH, temperature, agitation speed, and adsorbent dosage all have an impact on how well the green macroalgae biosorb antibiotic is. For equilibrium studies, the isotherm biosorption and kinetics models and thermodynamic modeling are also assessed. Therefore, our objective was to examine the effectiveness of Pithophora macroalgae in the removal of AMX from aqueous solutions. Pithophora is a filamentous green alga that creates free-floating mats, which refers to Cladophorales members (O'Neal et al. 1985).

MATERIALS AND METHODS

Materials and Instruments

The following materials and instruments were used in this experiment: 10 L distilled water; 10 L tab water; 1 L Methanol; 1 g Amoxicillin antibiotic; 8 Erlenmeyer flasks; 8 Lab shaker bottles; one measuring cylinder; 2 Pipettes; 1 Flask; 1 beaker; volumetric cylinder; 0.1 NaOH; 0.1 HCL solution; and weighing Scale (Saturius CP, model CPA 2250). In addition, 1 mortar and pestle, 1 magnetic stirrer or batch, a Bath water shaker, and a UV-visible spectrophotometer



were used. (National Center for Biotechnology Information 2022).

The Procedure for Preparing the Stock Solution

The amoxicillin (AMX) antibiotic component was obtained from a local pharmacy in Istanbul, Turkey, for analytical reference purposes without undergoing any further processing. The antibiotics' characteristics are listed in Table 1; the following process was used to use the AMX: The antibiotic was precisely weighed by weighing scale. The stock solution $(1g.L^{-1})$ was prepared by grounding the antibiotics, 1 g of AMX antibiotic using mortar and pestle to make them as powders in the lab. Dissolving accurately weighted amounts of the antibiotics in suitable solvents.

Table 2: The concentrations of AMX and its absorptions.

Conc. mg.L ⁻¹	absorbance
10	0.345
30	1.108
50	1.512
70	1.912
100	2.89
115	3.31
125	3.59
150	4.56

The AMX antibiotic was dissolved in methanol and water. Then, the antibiotic powders were added to the magnetic stirrer for around 2 h at 70°C to speed up the dissolving. After that, the stock solutions were put in the refrigerator at 4°C further to use it. 0.1 M NaOH and 0.1 M HCl were used to adjust the pH.

Many standards calibration charts were prepared for a wide range of concentrations that were close to the anticipated concentration of the analyte, measuring the absorbance of different Antibiotic (AMX) concentrations at different (λ) nanometers and unknown concentrations of antibiotics before and after adsorption were computed from the calibration chart. In this work, the calibration curve was plotted using different wavelengths as given for the AMX at 231 nm. UV spectrometer, which would be based on the wavelength obtained during spectrum research. As shown in Fig. 1, the calibration curve was obtained for the range of 10 to 150 mg.L⁻¹. N.B: the experiment was repeated three times, and the mean values and standard deviations from the mean values



CALIBRATION OF AMX

Fig. 1: Calibration curve of AMX at different concentrations.

were recorded. Table 2 presents the various concentrations of AMX alongside their corresponding absorbance values. A double beam/visible spectrophotometer (Shimadzu UV-2450) was employed to measure absorbance at the specific wavelength characteristic of the compound, facilitating the determination of an unknown antibiotic concentration.

The measurement of antibiotic concentration is widely regarded as one of the most commonly utilized methodologies. When appropriately calibrated, this technique can exclusively analyze the UV-visible spectral properties of the substance, thereby providing accurate estimations of AMX concentration. The UV spectrophotometer utilized in the laboratory during the experiments is depicted by the spectrophotometer device (Fig. 2). The standard deviation of the measurements is reported to be 1.41. The best-fitting isotherm has been found in recent years using a variety of error analysis techniques, including the coefficient of determination (\mathbb{R}^2), the sum of the errors squared, a hybrid error function, Marquardt's percent standard deviation, the average relative error, and the sum of absolute errors which noted that the \mathbb{R}^2 of linear regression analysis alone is not a suitable metric for assessing an isotherm model's fit quality (Allen et al. 2003).

Determination of pH and Zeta Potential Charge

The pH of the growth media was assessed at the commencement and conclusion of each trial, as well as on designated sample days and during the preparation of the medium, utilizing a pH meter (WTW Inolab pH 7110 masa Tipi pH meter). Additionally, the pH of the metal removal media was adjusted



Fig. 2: Shimadzu UV-2450 double-beam UV/visible spectrophotometer.



Fig. 3: Effect of initial pH and contact time on sorption of AMX (Khamayseh & KIdak 2024).

before the initiation of the experiment. To ensure accurate pH measurements, a two-point calibration with buffer solutions was routinely conducted. The determination of the zero-point charge (ZPC) of the Pithophora algae was carried out at a pH of 7.6, following the mass titration method as described by Cristiano et al. (2011). Specifically, fifty milliliters of 0.01 M NaNO₃ were placed in Erlenmeyer flasks, and appropriate biosorbent dosages ranging from 0.1 to 1.5 grams were introduced. The samples were agitated in a mechanical shaker for twenty-four hours, after which the residual pH value was measured.

The neutral charge of the bio-sorbent was assessed utilizing the pH drift method. Under optimal conditions,

the kinetic profiles illustrating the biosorption removal of AMX, employing non-linear forms across varying solution pH levels, are presented in Fig. 3. A gradual biosorption process was observed, characterized by an increase in contact time across all pH levels investigated. The AMX antibiotic was effectively removed at an initial pH level, reaching equilibrium after 120 min, beyond which no further sorption occurred. The influence of pH on the adsorption process can be attributed to the solubility of pollutants and the presence of active functional groups on the adsorbents, which facilitate electrostatic interactions between the antibiotic molecules and the surface of the algae. (Khamayseh & Kidak 2024).



Fig. 4: Fresh Pithophora macroalgae harvested from the CIU University (2022).



Fig. 5: Pithophora cells under electron microscope, CIU labs-North Cyprus.



Fig. 6: The relation between ln Kd and 1/T of AMX.

Table 3: Thermodynamics for biosorption of AMX at different temperatures.

Adsorbate	T[℃]	qe [mg.g ⁻¹]	Thermodynamic parameters				
			$\Delta G^{\circ} [kJ.moL^{-1}]$	$\Delta H^{o} [kJ.moL^{-1}k]$	$\Delta S^{o} [J moL^{-1} K)]$		
AMX	25	18.0	2.982	49.796	-176.735		
	30	16.0	3.718				
	35	12.1	4.793				

Collection and Identification of the Algae

This study made use of the (seaweed) Green *Pithophora* Macroalgae (GPM) as in Fig. 4. Between October and November 2021, during the winter months, they were gathered from the pond at Cyprus International University (CIU). To create the 40 g powdery form, around 8 kg of pond-harvested material was pulverized in a milling machine. After which they were dried at 40 degrees and grounded at 45 microns.

Under the electron microscope shown in Fig. 5, Pithophora was seen, which is made up of filaments irregularly branching and frequently include akinetes; many inflated reproductive cells resemble spores. From lime green to a dark greenish brown, it can be any color. When gas bubbles created by the plant are caught within the thick algal growth, they become buoyant, and the surface mats typically form in warmer weather. These mats may momentarily sink to the bottom if they are disturbed by significant wind or rain events.

RESULTS AND DISCUSSION

Effects of Temperature

The temperature is typically affected by processes involving heat or mass transport. A thermodynamic process alters the thermodynamic state of a system. r is defined as the degree of chaos or uncertainty in a system. Enthalpy is a key concept in thermodynamics. It is the overall amount of heat in the system. This indicates that as more energy is provided, the enthalpy rises. To determine whether antibiotic adsorption by Pithophora algae occurs spontaneously, entropy and energy factors should be considered. An experimental investigation was conducted to examine the influence of temperature on the adsorption of AMX by Pithophora algae. The findings indicated that the capacity for antibiotic adsorption diminishes as the temperature increases. (Khamayseh & Kidak 2024).

At temperature ranges of 298 K to 308 K, the removal effectiveness of AMX medication at various start concentrations was investigated. A 100 mL solution was applied to 0.5 g of biomass in Erlenmeyer flasks. After that, it was shaken in a temperature-controlled shaker at 150 rpm. Samples were taken anywhere from five to eighty-one minutes apart at various times. The equations were used to calculate Gibb's free energy (ΔG°). (4.1), (4.2), and (4.3) (Aravindhan et al. 2009).

$$\Delta G^{\circ} = \Delta H^{\circ} - T \Delta S^{\circ} \qquad \dots (4.1)$$

$$\Delta G^{\underline{o}} = -RT \ln Kd \qquad \dots (4.2)$$

$$K_d = \frac{q_e}{c_e} \qquad \dots (4.3)$$

Where T is the absolute temperature (K), R is the gas

constant (8.314 J moL⁻¹.K⁻¹), qe is the amount of antibiotic absorbed onto the biosorbent at equilibrium time (mg/g), and Ce is the equilibrium concentration of the sorbate present antibiotics (mg.L⁻¹).

Van't Hoff plots ln Kd vs. 1/T (Fig. 6) can be used to determine entropy (Δ S°) and enthalpy (Δ H°) based on their slope and intercept. Also, entropy (Δ S) and enthalpy (Δ H) can be determined using the following equations (4.4) and (4.5).

$\Delta H^{o} =$	Slope *	R gas constant	(4.4)

$$\Delta S^{\circ} = \text{Intercept } * \text{ R gas constant} \qquad \dots (4.5)$$

The Langmuir isotherm equilibrium constants at 298 K, 303 K, and 308 K were used to calculate ΔG° , ΔH° , and ΔS° for AMX adsorption by *Pithophora* (Table 3). At increasing temperatures, the biosorption % shows a trend toward decline. This decrease may be explained by an increase in the propensity of ions to break bonds and move from the solid to the bulk state, the destruction or deactivation of active sites on the surface of the biosorbent as a result of bond breaks, or the mobilization of ions in solution as a result of highly energized metal ions (Sulaymon et al. 2013). The coefficient of correlation value, R², of the van't Hoff plot in Fig. 6 was deemed satisfactory at 0.996 for AMX. For the necessary system, the thermodynamic parameters ΔG° , ΔS° , and ΔH° were measured. As can be shown in Table 3, the charge of ΔG° for AMX is positive (2.982 kJ.moL⁻¹, 3.718 kJ.moL⁻¹, and 4.793 kJ.moL⁻¹) at (298 K, 303 K, and 308 K, respectively. This positive value suggests that the reaction is neither possible nor spontaneous.

According to the ΔH° values, the biosorption process could be viewed as a physical sorption that is facilitated by a chemical reaction. Given that the electrical charge of ΔH° for AMX is positive at 49.796 KJ.moL⁻¹. This shows that the method of biosorption was endothermic, indicating that the antibiotics and biomass have a significant interaction. This finding indicates that the biosorption process is endothermic, suggesting a significant interaction between the biomass and the antibiotics. Consequently, algal biomass can be effectively utilized for the removal of AMX at ambient temperature. Entropy effects can be considered as the driving force of the biosorption process. Consequently, the biomass of algae demonstrates efficacy in the removal of AMX at ambient temperature. It can be posited that entropy effects are instrumental in driving the biosorption process. The reduction in disorder at the liquid/solid interface, resulting from the biosorption of AMX onto Pithophora macroalgae, is indicated by the negative value of ΔS° , which is measured at -176.735 kJ.moL⁻¹.

The measured qe values of the biomass of AMX decreased substantially from 18.0 to 12.1 mg/g when the



Fig. 7: Effect of temperature on the equilibrium AMX and uptake capacity.

Table 4: Biosorption isotherm constants obtained for various models obtained through linear regression analysis for AMX removal by Pithophora biosorbent at different temperature.

Antibiotic	T [K]	qe $_{\rm EXP}$ (mg/g)	Freundlich			Langmuir		
			K_{f} [mg.g ⁻¹]	Ν	\mathbb{R}^2	$Q_m [mg.g^{-1}]$	K[L.mg []]]	R^2
AMX	298	18.0	1.17	1.43	0.930	25.83	0.0345	0.944
	303	16.0	1.07	1.51	0.921	18.79	0.0367	0.954
	308	12.1	1.03	1.63	0.841	15.69	0.076	0.942

temperature of an antibiotic solution rose from 298 K to 308 K, as depicted in Fig. 7's plot of the influence of the biosorption of AMX antibiotic. This may indicate a decrease in the viscosity of the antibiotic-containing aqueous solution. As a result, the total volume of the biosorbent particles and the adsorbate's diffusion rate across pore borders may both rise (Mohammed et al. 2020).

Biosorption Isotherm at Different Temperatures

At all the temperatures investigated, Table 4 demonstrates that the adsorption data of AMX can be better described by the Langmuir isotherm than the Freundlich isotherm. According to the Langmuir isotherm's values for qm, the two antibiotics under study have a significant amount of



For AMX biosorption at 298 K, 303 K, and 308 K, pseudo-first order, pseudo-second order, and intraparticle diffusion kinetics are presented. a) Pseudo-first order of AMX b) Pseudo-second- order of AMX, c) intraparticle diffusion

Fig. 8: Biosorption Kinetics at different temperatures: (a-c).

Adsorbent	C_o [mg. ⁻¹ L]	$q_{e,EXP}$ [mg.g ⁻¹]	Pseudo-first order			Pseudo-se	Pseudo-second order			Intraparticle diffusion		
Pithophora Algae			$K_1([]$	$q_{e,cal}$ [mg.g ⁻¹]	R_2	$K_2(g/mg min^{-1})$	$q_{e,cal}$ [mg.g ⁻¹]	<i>R</i> ₂	K_{DIF} [g. g ⁻¹ min ^{-1/2})	C [g.g ⁻¹]	R ²	
	10	1.52	0.0091	1.31	0.985	0.0641	1.60	0.999	0z0687	0.7059	0.965	
	30	4.80	0.0047	2.91	0.942	0.0093	5.32	0.993	0.2932	1.2117	0.976	
	50	8.00	0.0057	6.18	0.996	0.0031	9.48	0.986	0.5830	0.7619	0.986	
	70	10.60	0.0052	7.89	0.988	0.0025	11.79	0.988	0.7532	1.1859	0.984	
	100	13.40	0.0094	10.93	0.989	0.0026	15.40	0.994	0.9256	2.3055	0.979	
	115	14.80	0.0092	11.70	0.991	0.0025	16.88	0.996	1.0127	2.7025	0.977	
	125	15.40	0.0104	11.38	0.993	0.0033	17.08	0.999	0.9851	3.8705	0.956	
	150	18.00	0.0135	14.66	0.983	0.0029	20.02	0.998	1.1795	4.4310	0.944	

Table 5: Results comparison between Pseudo-first order, pseudo-second order kinetics and intraparticle diffusion for AMX biosorption at 25°C (Khamayseh & Kidaz 2024).

adsorption capacity for algae biosorbent and adsorption capacity for AMX. The experimental qm values of the biomass of AMX dramatically dropped from 25.83 to 15.69 mg.g⁻¹. This may indicate a decrease in the viscosity of the antibiotic-containing aqueous solution. As a result, the adsorbate's diffusion rate may rise through the biosorbent particles' pores and bulk borders (Mohammed et al. 2020). The greater R² of the Langmuir isotherm model in Table 4 for all AMX values indicates that it provides the best fitting for the experimental data. Having a 25.83 mg/g maximal capacity for sorption, the experimental data for AMX biosorption most closely resembles the Langmuir isotherm model (Khamayseh & Kidak 2024).

Biosorption Kinetics at Different Temperatures

The Biosorption kinetics determine the sorption mechanism, which involves mass transfer, diffusion, and reaction on the biosorbent surface, as well as the reaction rate. This involves how the system sorption qualities vary over time and how much of the surface is covered, which in turn provides information about how quickly a process is happening. The design and regeneration of the biosorbent rely heavily on the rates of adsorption since sorption and desorption processes are both time-dependent. The experimental data are analyzed

using dynamic models to study the component of biosorption, the potential rate and governing steps, mass transfer, and the chemical reaction process (Ho & Mckay 1999). In other words, the sorption equilibrium also reveals details about the adsorption process dynamics. Based on the physicochemical characteristics of the antibiotic and the bio-sorbent, the kinetics of the biosorption process aid in assessing the effectiveness and identifying the type of sorption mechanism (Kerkez-Kuyumcu et al. 2016). The capacity for biosorption at equilibrium and the rate of solute binding on the surface of the biological material may both be calculated using biosorption kinetics. This material gives crucial details regarding a potential biosorption mechanism that includes chemical reactions and diffusion (Kratochvil & Volesky 1998). Three kinetic models-pseudo-first-order (PFO), pseudo-second-order (PSO), and intraparticle diffusion models-were employed to investigate the biosorption kinetics of AMX. The rate constants for these models were determined through linear regression analysis of the experimental data, as presented in Table 5. The experiment was conducted under optimal conditions, utilizing an initial concentration of 50 mg.L⁻¹, and the kinetics were examined at a temperature of 25°C.

The pseudo-second-order (PSO) model demonstrated a superior fit to the experimental data, exhibiting a

Table 6: Results comparison between Pseudo-first order, pseudo-second order kinetics and Intraparticle diffusion for AMX biosorption at different temperatures.

	Temp.	emp. q _e ,EXP)	p. q _e ,EXP) Pseudo-First Order			Pseudo-Second Order			Intraparticle Diffusion		
		[mg.g ⁻¹]	$K_1[[\min^{-1}]]$	$q_{e,cal}$ [mg.g ⁻¹]	R^2	g/K ₂ (mg/min)	$q_{e,cal} \mathrm{[mg.g^{-1}]}$	R ²	K _{dif} [mg/g min ^{1/2})	C [mg.g ⁻¹]	R^2
AMX	298 K	8.00	0.0057	6.18	0.996	0.0031	9.48	0.999	0.5830	0.7619	0.986
	303 K	7.60	0.0064	6.10	0.986	0.0029	9.18	0.975	0.5687	0.593	0.971
	308 K	7.03	0.0062	5.49	0.943	0.0034	8.40	0.981	0.529	0.546	0.961

regression coefficient (R²) of 0.999 and a rate constant of 0.0641 g.mg⁻¹·min⁻¹ at an initial amoxicillin (AMX) concentration of 10 mg.L⁻¹. Additionally, at the maximum AMX concentration of 150 mg.L⁻¹, the PSO rate constant (K₂) was determined to be 0.0029 g.mg^{-1} .min⁻¹, with an R² value of 0.998 at a temperature of 25°C. The PSO model was more obvious to fit with the experimental data (Table 5), demonstrating that the PSO model fitted better with the experimental data than the PFO model and the intraparticle diffusion model in terms of regression coefficients for all AMX concentrations. This is further shown by contrasting the experimental data's qe values with the one inferred from the plots (qe.CAL). The intraparticle diffusion method was then used to examine the data to gain additional insight into the adsorption mechanism. The potential of intra-particle diffusion was explored using experimental data and the Weber-Morris intra-particle diffusion model. The lower \mathbf{R}^2 values obtained corroborate the conclusion drawn from Table 5 data analysis: intraparticle diffusion is not able to explain the experimental results, the intraparticle diffusion cannot explain the experimental results, which is confirmed by the lower R² values attained in the AMX antibiotic. All things considered, the PSO model appears to have the greatest match to the study data, indicating that the biosorption of AMX onto Pithophora algae is caused by chemisorption interaction which is responsible for the biosorption of AMX. This correlates with previous research (Oba & Aydinlik 2022, Dan et al. 2021, Hamadeen et al. 2021, Pezoti et al. 2016). Pseudo-first-order and pseudo-second-order kinetic model equations were used to model the experimental data (Fig. 8 a, b). To determine whether film diffusion or intraparticle diffusion is the rate-controlling step, kinetic parameters were calculated using the Weber and Morris kinetic model to investigate intraparticle diffusion Fig. 8 (c). It can be said that the pseudo-second-order kinetic model and the experimental data closely match each other Additionally, qe, calc based on the pseudo-second-order isotherm has values that are comparable to qe, exp. The temperature has an impact on parameter values like k₂, qe, _{calc}, but not in a way that makes it necessary to treat it less than 298 K. For Kinetic analyses, Table 6 displays the values for the K_{diff} and C parameters from the modeling along with the rate constants, k, and adsorption capacity. The regression coefficient, or R², values are also displayed in the same table. According to Table 6, the experiment was carried out on the optimal condition at 50 mg.L⁻¹ initial concentration, and the kinetic was studied at different temperatures. It was seen the ge decreased by increasing the temperature in all types of the kinetics under 298 K, 303 K, and 308 K. So, the best-fit temperature is 298 K, which indicates high qe in pseudo-second order as the following order 9.48, 9.18, and 8.40 (mg.g⁻¹) for AMX

at 298 K, 303K, and 308K, respectively. We looked at the Freundlich and Langmuir isotherms to see which model fit the experimental data best. The experimental results' kinetic model adheres to the pseudo-second-order ($R^2 = 0.999$). In practical water and wastewater treatment applications, the biomass of the macroalgae Pithophora has been demonstrated in this study to be a potential biosorbent for the biosorption of AMX antibiotics (Khamayseh & Kidak 2024).

CONCLUSIONS

There is significant interest in identifying effective strategies for the removal of highly hazardous organic chemicals from water and wastewater. Among these strategies, adsorption has emerged as a recognized method for eliminating organic contaminants, noted for its efficiency, cost-effectiveness, and ability to yield high-quality treated effluent. Thermodynamic analyses have indicated that the biosorption process is endothermic and nonspontaneous. The findings of this study suggest that Pithophora algae represent a promising biosorbent for the extraction of the antibiotic amoxicillin from aqueous environments in practical wastewater applications. Pithophora was selected for this research due to its prevalence in urban areas, propensity for algal blooms, and natural dominance in nutrient-rich or eutrophic conditions. Furthermore, its filamentous structure contributes to reduced harvesting costs. When compared to other materials, algal biomass is the most frequently utilized biosorbent. Algae possess a high capacity for binding metal ions due to their untreated state and the characteristics of their cell walls, rendering them a cost-effective biosorbent option. Future investigations should focus on the specific functional characteristics of Pithophora's cell wall, particularly the presence of chlorophyll and its implications for the biosorption process. Additionally, further research is necessary to explore modeling, the regeneration of biosorbent materials, and the testing of adsorbed raw biomasses in relation to antibiotics. Advancements in techniques that enhance biosorbent processing will be essential for optimizing the application of biosorbent technology and regenerating biomass. • It is advised to use modified pithophora algae in a subsequent isotherm study to compare raw, unmodified algae with modified macroalgae treated with NaOH, formaldehyde, and CaCl₂ under different temperatures related to the thermodynamics methods. The potential for biological process improvement is limited because cells are not metabolizing, and there is no ability to biologically change the metal cationic and anionic state. The scientific understanding of the ion-binding mechanisms of these materials remains limited, highlighting the need for ongoing research and development in this area.

REFERENCES

- Abd, I.N. and Mohammed-Ridha, M.J., 2021. Simultaneous adsorption of tetracycline and amoxicillin by Cladophora and Spirulina algae biomass. *Iraqi Journal of Agricultural Sciences*, 52(5), pp.1290–1303.
- Aljeboree, A.M., Alshirifi, A.N. and Alkaim, A.F., 2020. Removal of amoxicillin drug onto nanocomposite surface using ultrasound assistance. *Plant Archives*, 20(1), pp.3135–3140.
- Allen, S.J., Gan, Q., Matthews, R. and Johnson, P.A., 2003. Comparison of optimized isotherm models for basic dye adsorption by kudzu. *Bioresource Technology*, 88(2), pp.143–152.
- Aravindhan, R., Rao, J.R. and Nair, B.U., 2009. Application of a chemically modified green macro alga as a biosorbent for phenol removal. *Journal* of Environmental Management, 90(5), pp.1877–1883.
- Arihilam, N. and Arihilam, E., 2019. Impact and control of anthropogenic pollution on the ecosystem – A review. *Journal of Bioscience and Biotechnology Discovery*, 4, pp.54–59. DOI.
- Asim, N., Amin, M.H., Alghoul, M.A., Sulaiman, S.N.A., Razali, H., Akhtaruzzaman, M. and Sopian, K., 2021. Developing chemically treated waste biomass adsorbent for dye removal. *Journal of Natural Fibers*, 18(7), pp.968–977.
- Balarak, D., Joghatayi, A., Mostafapour, F.K. and Azarpira, H., 2017. Biosorption of amoxicillin from contaminated water onto palm bark biomass. *International Journal of Life Science and Pharma Research*, 7(1), pp.L-9.
- Boukhelkhal, A., Benkortbi, O., Hamadache, M., Ghalem, N., Hanini, S. and Amrane, A., 2016. Adsorptive removal of amoxicillin from wastewater using wheat grains: Equilibrium, kinetic, thermodynamic studies, and mass transfer. *Desalination and Water Treatment*, 57(56), pp.27035–27047.
- Chakhtouna, H., Benzeid, H., Zari, N. and Bouhfid, R., 2021. Functional CoFe₂O₄-modified biochar derived from banana pseudostem as an efficient adsorbent for the removal of amoxicillin from water. *Separation and Purification Technology*, 266, 118592.
- Cristiano, E., Hu, Y.J., Siegfried, M., Kaplan, D. and Nitsche, H., 2011. A comparison of point of zero charge measurement methodology. *Clays* and Clay Minerals, 59(2), pp.107–115.
- Dan, A., Chen, C.X., Zou, M.Y., Deng, Y.Y., Zhang, X.M., Du, J.J. and Yang, Y., 2021. Removal efficiency, kinetic, and behavior of antibiotics from sewage treatment plant effluent in a hybrid constructed wetland and a layered biological filter. *Journal of Environmental Management*, 288, 112435.
- Dimitrakopoulou, D., Rethemiotaki, I., Frontistis, Z., Xekoukoulotakis, N.P., Venieri, D. and Mantzavinos, D., 2012. Degradation, mineralization, and antibiotic inactivation of amoxicillin by UV-A/TiO₂ photocatalysis. *Journal of Environmental Management*, 98, pp.168–174.
- Gaddam, K., SivananainthaPerumal, M. and Ravindiran, G., 2020. Removal of lead metal ion using biowaste of *Pithophora cleveana* Wittrock and *Mimusops elengi. Energy Sources, Part A: Recovery, Utilization, and Environmental Effects*, pp.1–19.
- Hamadeen, M., Elsayed, A., Elkhatib, A., Mohamed, E., Badawy, A., Samir, A. and Abdelgaleil, M., 2021. Novel low-cost nanoparticles for enhanced removal of chlorpyrifos from wastewater: Sorption kinetics, and mechanistic studies. *Arabian Journal of Chemistry*, 14, 102981. DOI
- Ho, Y.S. and McKay, G., 1999. Pseudo-second order model for sorption processes. *Process Biochemistry*, 34(5), pp.451–465. DOI
- Jung, Y., Kim, W., Yoon, Y., Kang, J., Hong, Y. and Kim, H., 2012. Removal of amoxicillin by UV and UV/H₂O₂ processes. *Science of The Total Environment*, 420, pp.160–167. DOI
- Kalyani, G., Gokulan, R. and Sujatha, S., 2021. Biosorption of zinc metal ion in aqueous solution using biowaste of *Pithophora cleveana*

Wittrock and *Minusops elengi*. *Desalination and Water Treatment*, 218, pp.363–371.

- Kerkez-Kuyumcu, Ö., Bayazit, Ş.S. and Salam, M.A., 2016. Antibiotic amoxicillin removal from aqueous solution using magnetically modified graphene nanoplatelets. *Journal of Industrial and Engineering Chemistry*, 36, pp.198–205.
- Khamayseh, M.M. and Kidak, R., 2023. Equilibrium, kinetics, and thermodynamics study on the biosorption of reactive levofloxacin antibiotic on *Pithophora* macroalgae in aqueous solution. *Environmental Monitoring and Assessment*, 195(2), 301.
- Khamayseh, M.M. and Kidak, R., 2024. Biosorption of reactive amoxicillin antibiotic on *Pithophora* macroalgae in aqueous solution. Equilibrium and kinetic studies. *Desalination and Water Treatment*, 100669.
- Kratochvil, D. and Volesky, B., 1998. Advances in the biosorption of heavy metals. *Trends in Biotechnology*, 16(7), pp.291–300.
- Lai, C., Liu, X., Qin, L., Zhang, C. and Zeng, G., 2017. Chitosan-wrapped gold nanoparticles for hydrogen-bonding recognition and colorimetric determination of the antibiotic kanamycin. *Microchimica Acta*, 184(7), pp.2097–2105. DOI
- Limousy, L., Ghouma, I., Ouederni, A. and Jeguirim, M., 2017. Amoxicillin removal from aqueous solution using activated carbon prepared by chemical activation of olive stone. *Environmental Science and Pollution Research*, 24(11), pp.9993–10004.
- Matsubara, M.E., Helwig, K., Hunter, C., Roberts, J., Subtil, E.L. and Coelho, L.H.G., 2020. Amoxicillin removal by pre-denitrification membrane bioreactor (A/O-MBR): performance evaluation, degradation by-products, and antibiotic resistant bacteria. *Ecotoxicology and Environmental Safety*, 192, 110258.
- Mohammed, A.A., Al-Musawi, T.J., Kareem, S.L., Zarrabi, M. and Al-Ma'abreh, A.M., 2020. Simultaneous adsorption of tetracycline, amoxicillin, and ciprofloxacin by pistachio shell powder coated with zinc oxide nanoparticles. *Arabian Journal of Chemistry*, 13(3), pp.4629–4643.
- National Center for Biotechnology Information., 2022. PubChem Compound Summary for CID 149096, Levofloxacin. Retrieved November 9, 2022 from https://pubchem.ncbi.nlm.nih.gov/compound/Levofloxacin
- Oba, O.A. and Pasaoglulari Aydinlik, N., 2022. Preparation of mesoporous activated carbon from novel African walnut shells (AWS) for deltamethrin removal: kinetics and equilibrium studies. *Applied Water Science*, 12(7), pp.1–20.
- O'Neal, S.W., Lembi, C.A. and Spencer, D.F., 1985. Productivity of the filamentous alga *Pithophora oedogonia* (Chlorophyta) in Surrey Lake, Indiana. *Journal of Phycology*, 21(4), pp.562–569.
- Pezoti, O., Cazetta, A.L., Bedin, K.C., Souza, L.S., Martins, A.C., Silva, T.L. and Almeida, V.C., 2016. NaOH-activated carbon of high surface area produced from guava seeds as a high-efficiency adsorbent for amoxicillin removal: kinetic, isotherm and thermodynamic studies. *Chemical Engineering Journal*, 288, pp.778–788.
- Romera, E., González, F., Ballester, A., Blázquez, M.L. and Muñoz, J.A., 2007. Comparative study of biosorption of heavy metals using different types of algae. *Bioresource Technology*, 98(17), pp.3344–3353.
- Sulaymon, A.H., Mohammed, A.A. and Al-Musawi, T.J., 2013. Removal of lead, cadmium, copper, and arsenic ions using biosorption: equilibrium and kinetic studies. *Desalination and Water Treatment*, 51(22–24), pp.4424–4434.
- Välitalo, P., Kruglova, A., Mikola, A. and Vahala, R., 2017. Toxicological impacts of antibiotics on aquatic micro-organisms: a mini-review. *International Journal of Hygiene and Environmental Health*, 220(3), pp.558–569.
- Yan, L., Yan, N., Gao, X.Y., Liu, Y. and Liu, Z.P., 2022. Degradation of amoxicillin by newly isolated *Bosea sp.* Ads-6. *Science of The Total Environment*, 828, 154411.