

## REVIEW ARTICLE

<https://doi.org/10.46488/NEPT.2026.v25i04.B4431>

## OPEN ACCESS

# Evaluating Green Infrastructure Interventions for Land Surface Temperature Reduction: A PRISMA-Based Systematic Review

**Avinash Kumar Singh† and Manoj Kumar**

Department of Architecture and Planning, National Institute of Technology, Patna-800005, Bihar, India

†Corresponding author: Avinash Kumar Singh; [avinash.singh1603arch@nitp.ac.in](mailto:avinash.singh1603arch@nitp.ac.in)

## ARTICLE HISTORY

Received: 06-02-2026

Revised: 08-04-2026

Accepted: 10-04-2026

## KEYWORDS

Land surface temperature  
Urban heat island  
Green infrastructure  
Urban morphology  
PRISMA

## CITE THIS ARTICLE

Singh, A.K. and Kumar, M., 2026. Evaluating green infrastructure interventions for land surface temperature reduction: A PRISMA-based systematic review. *Nature Environment and Pollution Technology*, 25(4), B4431. <https://doi.org/10.46488/NEPT.2026.v25i04.B4431>

## COPYRIGHT &amp; LICENCE

© 2026 The Authors  
Open Access under CC BY 4.0.

## PUBLISHED BY



## ABSTRACT

Rapid urbanization has intensified land surface temperatures (LST) in cities worldwide, exacerbating the urban heat island effect and increasing risks to thermal discomfort, public health, and energy demand. Green infrastructure (GI), including street trees, urban parks, green roofs, vertical greenery, and blue-green systems, has emerged as a key nature-based strategy for mitigating urban surface heat. This study presents a PRISMA-guided systematic review combined with bibliometric analysis to evaluate the effectiveness of GI interventions in reducing LST within built environments. A comprehensive search of the Web of Science Core Collection identified 211 records, of which 61 peer-reviewed studies published between 2012 and 2026 met the predefined inclusion criteria. Bibliometric analysis using VOSviewer reveals a rapidly expanding research field, characterized by exponential growth ( $R^2 = 0.9135$ ) and strong thematic convergence around keywords such as green infrastructure, land surface temperature, urban heat island, and built environment. Keyword co-occurrence mapping identifies multiple interconnected research clusters, highlighting integration across thermal regulation, urban morphology, climate adaptation, and methodological innovation, while also revealing the relative underrepresentation of social and equity-focused dimensions. Geographically, research output is concentrated in Asia, Europe, and Australia, indicating regional imbalances in evidence generation. The systematic synthesis shows that tree-based green infrastructure and urban parks consistently produce the largest LST reductions, commonly ranging from 2–4°C, with higher localized reductions, whereas green roofs and vertical greenery systems deliver more localized but critical cooling benefits in high-density urban areas. Cooling effectiveness is strongly context-dependent, shaped by climate zone, urban morphology, vegetation structure, and spatial configuration. Notably, several studies demonstrate that small-scale GI interventions can significantly reduce surface heat stress in informal settlements, suggesting potential equity implications, particularly in heat-vulnerable urban areas, although systematic evidence remains limited. By integrating bibliometric mapping with PRISMA-based content analysis, this review provides a context-sensitive, design-aware, and climate-responsive synthesis to support evidence-based urban heat mitigation and climate-resilient planning.

## 1. INTRODUCTION AND BACKGROUND

## 1.1. Urban Heat Challenges and Land Surface Temperature (LST)

Urbanization has substantially transformed natural landscapes, replacing vegetation and permeable surfaces with impervious materials such as asphalt, concrete, and metal (Mohan & Kandya 2015, Seto et al. 2012). This transformation has led to the amplification of Land Surface Temperature (LST) in cities, a phenomenon closely associated with the Urban Heat Island (UHI) effect (Evola et al. 2017, Nath 2025, Vyas et al. 2014). The UHI effect is characterized by higher daytime

and nighttime temperatures in urban areas relative to surrounding rural environments, driven primarily by solar heat absorption, limited evapotranspiration, anthropogenic heat emissions, and reduced ventilation (Gorai et al. 2024, Halder et al. 2025). Elevated LST is associated with increased surface heat exposure and can contribute to conditions that influence thermal discomfort, energy demand, and heat-related risks, although it does not directly represent human thermal experience (Abuwaer et al. 2023, Ullah et al. 2024, Ullah & Al-Ghamdi 2023).

Understanding the dynamics of LST is therefore critical for developing effective urban heat mitigation strategies. Temperature gradients between built and natural areas vary with urban density, building materials, surface albedo, and local climate conditions (Azmeer et al. 2024, Halder et al. 2025). High-density urban fabrics, particularly compact high-rise districts, are prone to intensified heat accumulation due to restricted airflow and increased radiation trapping. Conversely, urban open spaces and low-density areas benefit from higher convective cooling and larger potential for integrating vegetation-based interventions (Kumar et al. 2025).

These climatic and urban morphological complexities underscore the need for targeted, context-specific approaches to mitigate urban heat, particularly in rapidly urbanizing regions experiencing extreme temperatures (Rupapara et al. 2025). The evidence from past decades suggests that green

infrastructure (GI) interventions offer a promising solution for passive cooling by leveraging natural processes (Huang et al. 2022, Jin et al. 2024). GI is an effective strategy for urban heat mitigation, although its performance varies significantly depending on climatic conditions, urban morphology, and intervention type. In this study, the term GI is used as an umbrella concept encompassing urban greenery, including trees, parks, green roofs, vertical greenery systems, and blue-green infrastructure

## 1.2. Green Infrastructure as a Nature-Based Solution

GI encompasses a diverse array of interventions, including street trees, parks, urban forests, green roofs, and vertical greenery systems, which collectively aim to reduce urban heat and improve thermal comfort (Halder et al. 2026, Wong et al. 2021). The primary mechanisms through which GI reduces LST are shading, which limits direct solar radiation on urban surfaces, and evapotranspiration, whereby plants release water vapor to dissipate heat into the surrounding atmosphere (de Quadros & Mizgier 2023, Ying 2022). Together, these mechanisms contribute to significant surface and air temperature reductions, often producing perceptible cooling benefits for urban residents. Fig. 1 illustrates the dual mechanisms through which GI reduces urban temperatures: shading reduces heat absorption at the surface, while evapotranspiration dissipates heat into the atmosphere, collectively lowering LST.




Fig. 1: Mechanisms of Green Infrastructure LST Reduction.

Empirical studies have shown that tree canopies can reduce surface temperatures by up to 10°C, while urban parks create localized “cool islands” with air temperature reductions of 1–3°C. Green roofs and green walls, although generally providing smaller reductions in pedestrian-level temperatures, contribute to surface cooling, energy efficiency in buildings, and improvement of local microclimate conditions (Hurdic et al. 2024, Kanga et al. 2022, Luo & Wu 2021, Rahman et al. 2020, Weng et al. 2004). The effectiveness of these interventions is highly sensitive to factors such as vegetation type, canopy density, spatial arrangement, and species selection.

Also, multi-layered vegetation strategies, integrating trees, shrubs, and ground cover, enhance thermal performance by increasing the leaf area index and creating diverse microclimates. Proper planning of vegetation layout, including tree spacing and canopy continuity, has been shown to produce 20–30% higher cooling efficiency compared to scattered, single-layer plantings (Chen et al. 2022, Gherri 2023, Perini & Magliocco 2014). Selecting drought-tolerant and native species ensures both thermal performance and long-term sustainability, especially in arid and semi-arid urban regions (Lee & Jim 2018, Yang & Ma 2024).

### 1.3. Empirical Evidence on LST Reduction

Research on the thermal impact of GI has expanded considerably over the last decade, employing both field measurements and simulation-based approaches. Field studies capture local microclimatic variations and validate thermal performance at the pedestrian scale, while simulation models such as ENVI-met, CFD, and coupled urban climate-energy models enable scenario testing and optimization of vegetation placement (Battista et al. 2016, Chow et al. 2016, Liu et al. 2023, Pandey et al. 2021, Toparlar et al. 2017, Yang et al. 2023).

Cooling effectiveness has been found to vary according to climatic context, with hot-arid cities benefiting the most from evapotranspiration due to low atmospheric humidity and high solar radiation (Bowen et al. 2024, Cheung et al. 2021). In contrast, hot-humid climates rely more heavily on shading, as moisture saturation limits latent heat flux from evapotranspiration (Cheung & Jim 2018, Yang et al. 2023). Urban density also plays a significant role: medium-density areas often achieve optimal cooling by balancing vegetation coverage with airflow, whereas high-density districts may experience constrained benefits due to limited ventilation (Kumar & Shukla 2022). Spatial configuration of vegetation within urban canyons, including street orientation, tree height, and canopy coverage, further modulates thermal outcomes. Studies have demonstrated that continuous canopies produce

greater reductions in pedestrian-level LST than scattered plantings, while multi-layered arrangements combining trees, shrubs, and ground cover yield the largest cumulative cooling benefits (Ornelas et al. 2023, Pradeep Kumar et al. 2023, Zhou et al. 2011). These findings underscore the necessity as influenced by urban morphology considerations into green infrastructure planning for maximum effectiveness.

It is important to distinguish between LST and human thermal comfort. While LST provides a spatially explicit measure of surface heat patterns, it does not directly capture air temperature, mean radiant temperature, or human thermal indices such as PET or UTCI. Therefore, LST should be interpreted as a proxy indicator of surface heat exposure rather than a direct measure of human thermal stress.

### 1.4. Integration with Urban Morphology and Planning

The effectiveness of GI in reducing LST is inseparable from the geometrical and functional characteristics of the built environment. Urban morphology variables, including building height, spacing, and street canyon geometry, influence wind flow, solar exposure, and heat accumulation, which in turn affect vegetation performance (Chen et al. 2021, Huang et al. 2025, Jiang et al. 2021). Optimizing vegetation placement within these morphological constraints enhances both shading and convective cooling effects, yielding more effective LST mitigation (Kikon et al. 2023, Zhang et al. 2003).

Local Climate Zone (LCZ) classifications provide a framework for assessing and comparing green infrastructure performance across diverse urban forms (Kotharkar & Bagade 2018). Integration of GI with urban planning enables performance-based strategies, such as prescribing minimum canopy coverage, spatial distribution, and multi-layered planting to achieve targeted cooling outcomes (Fang et al. 2023). In addition, urban planners and policymakers can leverage these strategies to align GI interventions with broader goals, including energy efficiency, air quality improvement, and has potential implications for thermal comfort (Liu et al. 2021, O'Donnell et al. 2017, Shao et al. 2021). Sustainable urban design approaches also advocate combining GI with complementary measures such as reflective pavements, water features, and shading structures to maximize overall thermal comfort (de Oliveira et al. 2022, Derksen et al. 2017, Tzoulas et al. 2007). This holistic perspective ensures that vegetation interventions are not only effective in isolation but synergistically contribute to a cooler, more livable urban environment.

### 1.5. Knowledge Gaps and Study Rationale

Despite the growing body of research, several critical knowledge gaps remain. Many studies focus on single

intervention types, isolated urban contexts, or short-term temperature reductions, which limits the generalizability of findings (Hunter et al. 2015, Mittermüller et al. 2021). Additionally, methodological heterogeneity, including differences in measurement techniques (satellite-derived vs. ground-based LST), simulation assumptions, and temporal scales, complicates cross-study comparisons (Feng 2022, Zhang et al. 2024).

A systematic synthesis is therefore required to consolidate evidence on GI interventions and their effectiveness in LST reduction, considering the influence of climate, urban morphology, and vegetation type. This study adopts a PRISMA-guided systematic review to:

- Map the current state of research on GI and LST reduction.
- Evaluate methodologies and measurement approaches used in existing studies.
- Assess cooling effectiveness across different urban forms and climatic contexts.
- Identify gaps in knowledge and provide evidence-based recommendations for future research and urban planning.

By focusing on these objectives, the study aims to provide a comprehensive, climate-sensitive understanding of GI performance, supporting urban designers and policymakers in implementing effective LST mitigation strategies. This aligns with the study objective of identifying research gaps and supports the need for more context-sensitive and human-centered evaluation frameworks.

Unlike previous reviews, this study not only consolidates empirical evidence on LST reduction by GI but also categorizes interventions by urban morphology, vegetation type, and climate zone, offering a more context-specific understanding of cooling effectiveness. Furthermore, by applying the PRISMA framework systematically, this review enhances transparency, reproducibility, and methodological rigor, providing an updated reference for policymakers, urban planners, and researchers designing evidence-based urban cooling strategies.

## 2. MATERIALS AND METHODS

In this research, a mixed-methods approach was adopted by combining a PRISMA-based systematic review with bibliometric analysis using VOSviewer. This approach integrates qualitative synthesis of the literature with quantitative mapping of research trends to examine the relationship between GI and LST across built environments (Arruda 2022, Kirby 2023, Page et al. 2021, van Eck 2010).

PRISMA process led to the selection of the papers used for both qualitative synthesis and bibliometric analysis. The mixed-methods approach has been instrumental in achieving the objectives of this study (Li et al. 2022). Expanding upon previous discussions, this study integrates the structured qualitative depth of a systematic review with the quantitative insights of bibliometric analysis to examine the relationship between green infrastructure (GI) and land surface temperature (LST) across built environments. The research protocol is shown in Fig. 2. The study follows the PRISMA guidelines and checklist (Page et al. 2021).

A structured data extraction framework was developed to ensure consistency across the reviewed studies. Each study was coded based on predefined variables, including:

- (i) type of green infrastructure (e.g., trees, parks, green roofs, blue-green systems),
- (ii) urban morphology (e.g., compact, high-density, suburban),
- (iii) climatic or geographical context,
- (iv) LST measurement method (e.g., remote sensing, simulation, field measurement), and
- (v) reported cooling outcomes.

These categories were defined based on common classifications used in urban climate and green infrastructure literature and were applied consistently across all included studies.

### 2.1. The PRISMA Framework

The PRISMA framework was employed to systematically collect, screen, and analyse articles from the Web of Science (WOS) database. This review is based on the WOS database, which, while ensuring high-quality and peer-reviewed sources, may not capture all relevant studies indexed in other databases such as Scopus or Google Scholar. This represents a limitation in terms of literature coverage. This method enhances transparency and reproducibility, which are vital for academic rigor (Chaudhuri & Kumar 2022, Fu et al. 2022). A web search was conducted on January 04, 2026, as the initial step. Boolean operators (AND, OR) were applied to refine the results and ensure the selection of pertinent studies. The search string was designed to balance comprehensiveness and specificity by incorporating multiple synonymous terms related to land surface temperature, urban heat, and green infrastructure. The selected terms were chosen to ensure alignment with the core research objectives while minimizing retrieval of irrelevant studies. “LST” is first introduced in this paper and is a widely used metric in urban climate and remote sensing studies, particularly for assessing surface heat patterns and urban heat island effects.

Table 1: Criteria for eligible literature through WOS (excluded if not meeting these criteria).

| No. | Criteria                                                                                                                                                                                                                                                                     |
|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1   | Search Terms<br>(Topic/Title/<br>Abstract)<br>("Land Surface Temperature" OR "LST" OR "Surface Temperature" OR "Urban Heat" OR "UHI")<br>AND ("Green Infrastructure" OR "Urban Greenery" OR "Urban Green Space")<br>AND ("Urban Environment*" OR "Built Environment")        |
| 2   | Document Type<br>Article                                                                                                                                                                                                                                                     |
| 3   | Language<br>English                                                                                                                                                                                                                                                          |
| 4   | Database<br>Web of Science (WOS) database                                                                                                                                                                                                                                    |
| 5   | Irrelevant<br>Disciplines<br>Subject categories retrieved excluding:<br><i>Pharmacology, toxicology, and pharmaceuticals;</i><br>Physics and Astronomy;<br>Chemical Engineering;<br>Computer Sciences;<br>Biochemistry, Genetics and Molecular Biology;<br>Decision Sciences |

Furthermore, comprehensive relevant literature retrieval is based on LST in GI and Built Environment. Alternative terms such as 'urban greening' or 'ecosystem services' were considered but were not included to maintain focus on studies explicitly addressing thermal performance. Criteria for eligible literature through WOS are further classified in Table 1. The specific search query used in basic search mode; the search term was - ("Land Surface Temperature" OR "LST" OR "Surface Temperature" OR "Urban Heat" OR "UHI") AND ("Green Infrastructure" OR "Urban Greenery" OR "Urban Green Space") AND ("Urban Environment\*" OR "Built Environment") and then search method topic was selected.

The WOS core collection database provided 211 journal articles using this search. Fig. 2 illustrates the PRISMA 2020-compliant study selection process, detailing the number of records at each stage (identification, screening, eligibility, and inclusion), along with explicit reasons for exclusion. This ensures transparency and reproducibility of the review process. The process began with the identification stage, where 211 records were retrieved from the WOS database. This broad search aimed to capture a wide range of relevant studies within the research scope. During this phase, 26 articles were excluded because they were not in English, were review papers, or were categorized as early access articles, resulting in 185 records proceeding to the next phase.

The screening stage involved a structured evaluation of the remaining records based on predefined inclusion criteria. A total of 65 studies were excluded due to disciplinary misalignment. The remaining records were assessed for relevance, defined as explicit examination of the relationship between GI and LST within urban or built environments.

Screening was conducted sequentially at title, abstract, and full-text levels to ensure consistency and minimize subjectivity. Following this process, 120 studies were retained for further assessment, of which 59 were excluded due to lack of alignment with the research scope. The final selection comprised 61 studies that met all inclusion criteria.

In the eligibility stage, the full texts of the 61 remaining records were reviewed against predefined criteria, including relevance to the research themes of LST, GI, and the built environment. Finally, the inclusion stage resulted in 61 studies being selected for detailed analysis. These studies formed the foundation of the systematic review, providing critical insights into key themes, methodologies, and research gaps.

By employing the PRISMA framework, this review ensured a structured and reliable process, yielding a robust dataset that highlights both the methods adopted and research gaps in the field. This approach not only synthesizes existing knowledge but also facilitates future research by systematically identifying synergies and challenges in urban greenery and thermal comfort.

In the supplementary material, a comprehensive list of all 61 studies included in the systematic review is presented, along with citation details. Each study is systematically documented with information such as authors, year of publication, title, and number of citations, providing a transparent overview of the research corpus. The meticulous application of the PRISMA framework ensures that all selected studies are highly relevant to the study's themes and research objectives.

To make sure the papers included in systematic reviews are of high quality, Dybå & Dingsøyr in 2008 developed a checklist of questions (Dybå & Dingsøyr 2008). Three




Fig. 2: PRISMA framework of the study, Source: PRISMA 2020 (Page et al. 2021).

primary criteria serve as the foundation for these inquiries: relevance, rigor, and credibility. Data were extracted systematically from each study, including GI type, urban morphology, climate context, LST measurement method, and reported cooling effects. While no formal numerical quality scoring or sensitivity analysis was conducted, the review carefully considered the scope, methodology, and context of each study to identify patterns and trends. Potential limitations, such as geographic concentration of studies or methodological variability, were acknowledged and considered in the synthesis. This assessment informed the interpretation of results and helped identify potential sources of bias, including limitations related to data sources, spatial scale, and modeling assumptions. Given the heterogeneity of study designs and outcomes, a qualitative assessment approach was considered more appropriate than a standardized scoring system.

This structured approach ensured a robust and transparent review, providing a comprehensive overview of the current

evidence on the effects of green infrastructure on urban LST, while highlighting research gaps and opportunities for future studies. A complete list of all included studies, with citation details, is provided in the supplementary material.

## 2.2. Bibliometric Analysis

For the bibliometric analysis, this study utilized VOSviewer software (version 1.6.19) in conjunction with the WOS database. VOSviewer was employed to generate visual bibliometric maps (Arruda 2022, van Eck 2010), illustrating the co-occurrence of keywords and emerging research trends in literature related to LST and GI across built environments. This technique allowed for the identification of frequently used terms, research foci, and linkages between concepts, offering insights into the conceptual structure of the field. In addition to keyword analysis, the study examined the geographic distribution of research by analyzing country affiliation contributions. Publication trends were also assessed to determine the temporal evolution of scholarly

interest in the topic. These bibliometric insights complement the qualitative synthesis by mapping the scope and trajectory of academic engagement, thereby strengthening the review's capacity to identify research gaps and emerging areas of focus. For keyword co-occurrence analysis, author keywords were extracted from the selected studies. A minimum occurrence threshold was applied to include only frequently used terms, ensuring robustness of the network. Co-occurrence relationships were normalized using the association strength method, and clustering was performed using the VOSviewer clustering algorithm to group related themes.

Given the diversity of study designs, spatial scales, climatic contexts, and outcome metrics (as summarized in Table 2), a formal meta-analysis was not considered appropriate. Instead, a structured quantitative synthesis approach was adopted to systematically compare reported LST reductions across studies.

### 3. RESULTS

#### 3.1. Publication Trends

The publication trend illustrated in Fig. 3 reveals a clear and accelerating growth in scholarly interest in GI interventions for LST reduction over the past decade. From 2012 to 2019, research output in this field remained relatively sparse, with only one publication recorded in each of the selected early years. This limited output suggests that, during this period, GI–LST relationships were still an emerging research niche, often embedded within broader studies on urban climate, land use, or sustainability rather than being a primary focus. A

notable shift occurs from 2020 onwards, where the number of publications increases substantially. The annual output rises from three studies in 2020 to eight in 2021, followed by sustained growth through 2022 and 2023. This surge coincides with growing global concern over urban heat islands, climate change impacts, and heat-related health risks, as well as improved access to high-resolution remote sensing data and geospatial analytical tools. These developments have enabled more robust and spatially explicit assessments of how green roofs, urban forests, parks, and other GI strategies influence LST.

The upward trajectory becomes particularly pronounced after 2023, reaching 19 publications in 2025, the highest value in the review period. The fitted exponential trendline ( $R^2 = 0.9135$ ) indicates a strong and consistent growth pattern, suggesting that research in this domain is not only increasing but accelerating. This reflects the consolidation of GI-based thermal mitigation as a critical research theme within urban climate adaptation and sustainable city planning. Overall, the observed publication trend underscores the timeliness and relevance of this PRISMA-based systematic review. The rapid increase in recent studies highlights the need for a structured synthesis to evaluate the effectiveness of different green infrastructure interventions, identify methodological patterns and gaps, and inform evidence-based urban planning and climate resilience strategies.

#### 3.2. Analysis of Co-Occurring Keywords and Density Visualization

The keyword co-occurrence network (Fig. 4), generated using VOSviewer, illustrates the conceptual structure of



Fig. 3: Annual publication trend in GI–LST research with an exponential trendline.

the research field by mapping the frequency and strength of relationships between terms. The co-occurring keyword map produced using VOSviewer offers a nuanced representation of the conceptual structure and thematic breadth of research on GI and LST reduction. Based on 501 author keywords, the network is organized into 23 clusters, with a total link strength of 5209, indicating a highly connected and mature body of literature. The identified clusters represent groups of closely related keywords that frequently co-occur within the literature, indicating dominant thematic areas. Total link strength reflects the overall degree of connectivity among keywords, with higher values indicating stronger conceptual integration across the research field.

At the center of the network, keywords such as green infrastructure, temperature, urban environment, climate change, and built environment exhibit the highest connectivity, reflecting their foundational role in shaping the discourse. Their prominence suggests that GI–temperature interactions are most often examined within broader discussions of urban climate dynamics and the physical characteristics of cities. The dense interlinkages among these core terms indicate strong conceptual integration rather than fragmented thematic development.

Several clusters are clearly oriented toward thermal processes and measurement approaches, with recurring terms

including land surface temperature, surface temperature, air temperature, energy balance, and urban heat. These clusters underscore the dominant reliance on remote sensing, surface energy modeling, and climate simulations to quantify the cooling effects of green infrastructure. Closely connected keywords such as land cover, urbanization, density, and buildings highlight the recognition that GI performance is strongly conditioned by urban form, spatial configuration, and land-use patterns.

Methodological diversity is also evident across the network. Clusters containing terms such as simulation, evaluation tools, leaf area index, canopy shading, Google Street View, and citizen science point to increasing use of innovative data sources and mixed-method approaches. These methods enable more detailed characterization of vegetation structure, street-level thermal environments, and user perceptions, complementing traditional satellite-based analyses. Emerging conceptual framings are reflected in clusters associated with nature-based solutions, multifunctionality, biodiversity, and bioclimatic design.

Overall, the identification of 23 interrelated clusters with a high total link strength reveals a well-established yet evolving research field. The keyword network highlights both the consolidation of core themes, particularly the




Fig. 4: Co-occurring keyword map visualization using VOSviewer.

GI-LST relationship and the increasing integration of health, resilience, and methodological innovation.

Fig. 5 presents the keyword density visualization generated using VOSviewer, offering a complementary perspective to the network map by highlighting the relative concentration and prominence of research themes within the literature on GI and temperature mitigation. The density map highlights areas of high research intensity through darker and more saturated regions, enabling clearer identification of dominant and peripheral research themes.

The bibliometric findings highlight a strong concentration of research on biophysical aspects of temperature reduction, while social, behavioral, and equity-related dimensions remain underexplored. This gap is consistent with the findings of the systematic review and suggests the need for more integrated, interdisciplinary approaches in future research.

Results reveal a research field that is highly concentrated around urban thermal regulation, yet gradually diversifying towards potential implications for thermal comfort outcomes, multifunctionality, and innovative data sources. This imbalance underscores the need for future studies to better integrate social, ecological, and methodological dimensions into GI effectiveness assessments, an objective directly

addressed by the systematic synthesis undertaken in this review.

These clusters can be broadly interpreted into three dominant thematic domains:

- (i) thermal processes and measurement approaches (e.g., LST, surface temperature, energy balance),
- (ii) urban form and morphology (e.g., density, land use, urban structure), and
- (iii) nature-based and design-oriented strategies (e.g., green infrastructure, ecosystem services, resilience).

Importantly, the analysis reveals comparatively weaker representation of keywords related to human thermal comfort, social vulnerability, and equity. This indicates a research gap in integrating biophysical temperature reduction with human-centered outcomes, which is further explored in the systematic synthesis.

### 3.3. Analysis of Country Distribution

Fig. 6 illustrates the country-wise distribution of publications included in this systematic review, highlighting the geographical patterns of research activity on GI interventions for temperature reduction. The results reveal a clear concentration of scholarly output in a limited number of




Fig. 5: Keyword density visualization generated using VOSviewer, illustrating the thematic concentration and research intensity of co-occurring keywords in studies on green infrastructure and urban temperature mitigation.

countries, reflecting uneven global engagement with this research theme.

China emerges as the leading contributor, accounting for the highest number of publications ( $n = 15$ ). This dominance underscores the country's strong research focus on urban heat mitigation, rapid urbanization challenges, and the widespread application of remote sensing and spatial analysis techniques in GI-temperature studies. Australia follows with nine publications, reflecting sustained interest in climate-sensitive urban planning and heat resilience strategies in the context of extreme temperature exposure. Italy ranks third with seven studies, indicating a growing European emphasis on nature-based solutions and urban climate adaptation.


Germany and the United States contribute six publications each, demonstrating consistent research engagement in assessing the thermal and environmental performance of green spaces, urban forests, and other GI elements. England and India follow with five publications each, highlighting emerging but still moderate research activity, particularly in relation to urban sustainability and climate adaptation. Austria and Switzerland contribute three studies each, while Canada accounts for two publications, suggesting more limited but focused contributions. This geographical imbalance points to a need for broader cross-regional studies and greater inclusion of diverse climatic, socio-economic, and urban contexts. Importantly, the concentration of studies in specific regions may influence the reported effectiveness of GI, as cooling performance is highly dependent on local climate conditions, vegetation characteristics, and urban morphology.

As a result, findings derived predominantly from temperate and rapidly urbanizing regions may not be directly transferable to underrepresented regions such as Africa or parts of South America, where climatic conditions, urban forms, and resource constraints differ significantly. This highlights the need for caution when generalizing results and underscores the importance of context-specific evaluation of GI interventions.

The country distribution indicates that research on GI and land surface temperature reduction is predominantly concentrated in Asia, Europe, and Australia, with comparatively fewer contributions from North America and limited representation from the Global South beyond China and India. This geographical imbalance points to a need for broader cross-regional studies and greater inclusion of diverse climatic, socio-economic, and urban contexts. Addressing this gap is essential for improving the generalizability of evidence and informing context-sensitive green infrastructure planning at a global scale.

### 3.4. Content Analysis

This section presents the results of the content analysis of 61 selected studies that examined the comparative performance of land surface temperature (LST) and green infrastructure (GI) in built environments, as summarized in Table 2. The synthesis was conducted using a qualitative coding approach, in which recurring patterns and relationships across studies were identified and grouped according to GI type, urban morphology, and climatic context. This approach



Source: WOS

Fig. 6: Geographical distribution of publications.

Table 2: Content analysis of 61 selected studies that examined the comparative performance of LST and GI in Built environments.

| S. No. | Authors                       | Year | GI Type                             | Urban Morphology          | LST Measurement Method      | Key Inferences                                                                                        |
|--------|-------------------------------|------|-------------------------------------|---------------------------|-----------------------------|-------------------------------------------------------------------------------------------------------|
| 1.     | Oraiopoulos et al. 2026       | 2026 | Urban green infrastructure          | Informal settlements      | Remote sensing + field data | GI significantly reduces surface heat stress and improves habitability in dense informal settlements. |
| 2.     | Berger et al. 2026            | 2026 | Native vegetation                   | Residential urban areas   | Field experiments           | Native plant-based GI enhances thermal regulation under urban heat stress conditions.                 |
| 3.     | Helmreich et al. 2025         | 2025 | Infiltration swales (blue-green GI) | Urban neighborhoods       | ENVI-met simulation         | Blue-green GI provides measurable surface cooling while improving stormwater management.              |
| 4.     | Yang & La Roche 2025          | 2025 | Tree canopies, green roofs          | Dense urban blocks        | ENVI-met                    | Tree canopies outperform green roofs in reducing daytime LST in compact urban areas.                  |
| 5.     | Lazovic et al. 2025           | 2025 | Roadside greenery                   | Traffic corridors         | Numerical simulation        | Roadside GI mitigates LST and contributes to improved thermal conditions along high-traffic streets.  |
| 6.     | Hogeweiij et al. 2025         | 2025 | Urban parks                         | High-density city center  | Landsat LST                 | Urban parks function as thermal sinks with cooling effects extending into surrounding areas.          |
| 7.     | Liu et al. 2025               | 2025 | Green roofs                         | High-rise urban districts | Remote sensing              | Green roofs reduce roof-level LST but show limited neighborhood-scale cooling.                        |
| 8.     | Esperon-Rodriguez et al. 2025 | 2025 | Street trees                        | Street canyon morphology  | CFD + ENVI-met              | Tree height and spacing strongly influence cooling performance in street canyons.                     |
| 9.     | Roversi & Longo 2025          | 2025 | Blue-green corridors                | Riverfront urban areas    | Satellite imagery           | Combined vegetation-water systems generate stronger LST reductions than vegetation alone.             |
| 10.    | Bachofen et al. 2025          | 2025 | Pocket parks                        | Mixed-use urban zones     | Landsat                     | Small green spaces significantly reduce localized surface heat hotspots.                              |
| 11.    | Ramadan et al. 2025           | 2025 | Urban forests                       | Metropolitan scale        | MODIS                       | Large urban forests lower regional LST more effectively than fragmented green spaces.                 |
| 12.    | Cuce et al. 2025              | 2025 | Vertical greenery systems           | High-rise districts       | Field measurements          | Vertical GI reduces façade temperatures but has limited city-scale influence.                         |
| 13.    | Karimi et al. 2025            | 2025 | Green belts                         | Urban fringe              | Remote sensing              | Continuous green belts outperform fragmented GI in reducing surface temperatures.                     |
| 14.    | Chen & Hanlon 2025            | 2025 | Grassland GI                        | Low-density suburbs       | Landsat                     | Grass-based GI provides moderate cooling but is less effective than tree cover.                       |
| 15.    | Yang et al. 2025              | 2025 | Wetlands                            | Peri-urban areas          | Satellite imagery           | Urban wetlands exhibit strong daytime LST reduction due to evapotranspiration.                        |
| 16.    | Seidu et al. 2025             | 2025 | Urban trees                         | Residential neighborhoods | Field + satellite           | Increased tree canopy cover correlates with lower surface temperatures.                               |
| 17.    | Pattnaik et al. 2025          | 2025 | Green roofs                         | Compact urban form        | ENVI-met                    | Roof greening effectiveness depends on building height and roof coverage ratio.                       |
| 18.    | Al-Zghoul & Al-Homoud 2025    | 2025 | Linear greenways                    | Transport corridors       | Remote sensing              | Linear GI reduces thermal fragmentation across urban landscapes.                                      |
| 19.    | Bera & Nag 2025               | 2025 | Mixed GI systems                    | Dense urban fabric        | Landsat + modeling          | Integrated GI networks outperform isolated interventions in LST mitigation.                           |
| 20.    | Ni et al. 2025                | 2025 | Urban parks                         | Compact city              | Remote sensing              | Park size and shape strongly influence cooling intensity and reach.                                   |
| 21.    | Del Rosario et al. 2025       | 2025 | Street greenery                     | High-density streets      | Field measurements          | Shaded streets show significantly lower surface temperatures during peak heat.                        |
| 22.    | Yazdi et al. 2024             | 2024 | Blue-green infrastructure           | Flood-prone urban areas   | ENVI-met                    | Blue-green GI provides dual benefits of cooling and climate resilience.                               |
| 23.    | Chen et al. 2024              | 2024 | Green walls                         | High-rise buildings       | Thermal imaging             | Green walls lower wall surface temperatures but have localized effects.                               |
| 24.    | Chen et al. 2024              | 2024 | Urban vegetation                    | City-wide                 | Landsat time series         | Expansion of urban greenery correlates with long-term LST decline.                                    |
| 25.    | Frosini et al. 2024           | 2024 | Green roofs                         | Commercial districts      | Remote sensing              | Commercial roof greening reduces extreme surface heat accumulation.                                   |

Table Cont....

| S. No. | Authors                   | Year | GI Type                | Urban Morphology          | LST Measurement Method | Key Inferences                                                             |
|--------|---------------------------|------|------------------------|---------------------------|------------------------|----------------------------------------------------------------------------|
| 26.    | Chu et al. 2024           | 2024 | Urban trees            | Residential blocks        | Field sensors          | Tree shade significantly reduces ground and pavement temperatures.         |
| 27.    | Jimenez & de Adana 2024   | 2024 | Urban parks            | Mixed-density areas       | MODIS                  | Larger parks exhibit stronger and more stable cooling effects.             |
| 28.    | Guan & Zhang 2024         | 2024 | Vegetated courtyards   | Compact housing           | ENVI-met               | Internal green courtyards improve microclimate in dense housing complexes. |
| 29.    | Liu et al. 2024           | 2024 | Blue-green networks    | Metropolitan regions      | Remote sensing         | Network connectivity enhances regional LST mitigation.                     |
| 30.    | Azmeer et al. 2024        | 2024 | Urban agriculture      | Peri-urban zones          | Landsat                | Urban farming areas show measurable surface cooling benefits.              |
| 31.    | Ibric et al. 2024         | 2024 | Street trees           | Urban road networks       | Field + GIS            | Tree-lined streets reduce surface heat accumulation during summer.         |
| 32.    | Zakrzewska et al. 2023    | 2023 | Green roofs            | Residential buildings     | Thermal monitoring     | Roof vegetation moderates diurnal surface temperature extremes.            |
| 33.    | Fong et al. 2023          | 2023 | Urban greenery         | Informal settlements      | Remote sensing         | Even small-scale GI interventions reduce extreme LST in vulnerable areas.  |
| 34.    | Pragati et al. 2023       | 2023 | Urban parks            | City center               | Landsat                | Park cooling intensity decreases with distance from park edges.            |
| 35.    | La Rosa & Li 2023         | 2023 | Green corridors        | Compact cities            | ENVI-met               | Corridors facilitate spatial diffusion of cooling effects.                 |
| 36.    | Sobrinho et al. 2023      | 2023 | Blue infrastructure    | Waterfront districts      | Satellite imagery      | Water bodies enhance adjacent green cooling effects.                       |
| 37.    | Pritipadmaja et al. 2023  | 2023 | Urban trees            | Suburban areas            | Field measurements     | Mature trees provide stronger cooling than young vegetation.               |
| 38.    | Okumus & Terzi 2023       | 2023 | Urban greenery         | Rapidly urbanizing cities | Landsat                | Loss of GI intensifies surface urban heat island effects.                  |
| 39.    | Irfeey et al. 2023        | 2023 | Street greenery        | Street canyon             | CFD + field data       | Combined ventilation and vegetation regulate canyon LST.                   |
| 40.    | Schaffernicht et al. 2023 | 2023 | Urban parks            | Metropolitan scale        | MODIS                  | Parks contribute to city-scale thermal regulation.                         |
| 41.    | Urban et al. 2022         | 2022 | Green roofs            | Dense commercial areas    | Remote sensing         | Roof greening effectiveness varies with roof material and coverage.        |
| 42.    | Graef et al. 2022         | 2022 | Blue-green GI          | Coastal cities            | Satellite imagery      | Coastal blue-green systems moderate extreme surface heat.                  |
| 43.    | Reyhani et al. 2022       | 2022 | Nature-based solutions | Urban neighborhoods       | ENVI-met               | Integrated NBS designs enhance urban heat resilience.                      |
| 44.    | Anderson et al. 2022      | 2022 | Urban vegetation       | Mixed-use districts       | Landsat                | Vegetation density inversely correlates with LST intensity.                |
| 45.    | Shan et al. 2022          | 2022 | Urban parks            | High-density zones        | Remote sensing         | Park cooling is strongest during daytime heat peaks.                       |
| 46.    | Salvalai et al. 2022      | 2022 | Green roofs            | Residential districts     | Field measurements     | Green roofs reduce surface heat but require adequate maintenance.          |
| 47.    | O'Keeffe et al. 2022      | 2022 | Urban trees            | Compact cities            | Remote sensing         | Tree canopy coverage is a key determinant of LST reduction.                |
| 48.    | Chun et al. 2021          | 2021 | Blue-green corridors   | Urban regions             | GIS + Landsat          | Corridor continuity enhances spatial cooling efficiency.                   |
| 49.    | Vulova et al. 2021        | 2021 | Urban greenery         | City-wide                 | MODIS                  | City-scale GI distribution shapes surface thermal patterns.                |
| 50.    | Kabano et al. 2021        | 2021 | Street trees           | Commercial streets        | Field monitoring       | Shaded commercial streets experience lower surface temperatures.           |
| 51.    | Zhang et al. 2021         | 2021 | Green walls            | High-rise buildings       | Thermal sensors        | Vertical GI cools building envelopes under extreme heat.                   |
| 52.    | Fuentes et al. 2021       | 2021 | Urban parks            | Dense urban cores         | Landsat                | Larger parks produce more stable cooling effects.                          |

Table Cont....

| S. No. | Authors                | Year | GI Type                    | Urban Morphology        | LST Measurement Method  | Key Inferences                                                      |
|--------|------------------------|------|----------------------------|-------------------------|-------------------------|---------------------------------------------------------------------|
| 53.    | Back et al. 2021       | 2021 | Urban greenery             | Metropolitan areas      | Remote sensing          | Fragmentation of GI reduces its cooling effectiveness.              |
| 54.    | Bouzouidja et al. 2021 | 2021 | Blue infrastructure        | Urban watersheds        | Satellite imagery       | Water bodies significantly moderate surrounding LST.                |
| 55.    | Amani-Beni et al. 2021 | 2021 | Urban forests              | City outskirts          | MODIS                   | Peripheral forests contribute to regional cooling.                  |
| 56.    | Shin et al. 2020       | 2020 | Urban greenery             | Rapidly urbanizing city | Landsat time series     | Expansion of GI correlates with declining urban LST trends.         |
| 57.    | Lu et al. 2020         | 2020 | Street greenery            | Street canyon           | Field + CFD             | Vegetation and airflow jointly regulate urban surface temperatures. |
| 58.    | Sabatino et al. 2020   | 2020 | Nature-based solutions     | Urban neighborhoods     | ENVI-met                | GI-based designs improve thermal resilience under heat stress.      |
| 59.    | Makido et al. 2019     | 2019 | Green walls and roofs      | High-density urban      | Case study + monitoring | Building-scale GI reduces surface temperature and energy demand.    |
| 60.    | Revell & Anda 2014     | 2014 | Urban greenery             | City-scale              | Landsat                 | Spatial patterns of GI determine high-temperature zones.            |
| 61.    | Su et al. 2012         | 2012 | Mixed green infrastructure | Metropolitan regions    | Remote sensing          | GI density is inversely related to urban surface heat intensity.    |

enabled systematic comparison while accommodating methodological heterogeneity across the reviewed studies.

The synthesis was structured to directly address the research objectives outlined in the introduction, particularly by consolidating empirical evidence on LST reduction associated with GI interventions and categorizing findings based on urban morphology, vegetation type, and climate zone. This facilitates a more context-specific understanding of cooling effectiveness.

The aim of this synthesis is to extract recurring patterns, conceptual orientations, and practical insights across multiple dimensions of evaluation. Such a structured approach not only supports understanding of the current state of knowledge but also helps identify methodological inconsistencies and research gaps that may hinder evidence-based planning and policy development. By organizing the review in this manner, the analysis provides a more nuanced understanding of the practical advantages, trade-offs, and contextual applicability of LST in addressing challenges related to urban sustainability and infrastructure resilience.

The “key inferences” reported in Table 2 were derived through a structured synthesis of each study’s primary findings. Rather than subjective interpretation, these inferences summarize the main conclusions presented by the original authors, with emphasis on the relationship between GI characteristics and LST reduction. To minimize subjectivity, consistent coding criteria were applied across all studies during the data extraction and synthesis process.

This PRISMA-guided systematic review synthesized evidence from 61 peer-reviewed studies published between 2012 and 2026, focusing on the effectiveness of GI

interventions in reducing LST in urban environments. The temporal distribution of studies shows a marked increase after 2018, reflecting growing scientific and policy concern regarding urban heat under climate change (Su et al. 2012, Revell & Anda 2014, Makido et al. 2019, Lu et al. 2020, Di Sabatino et al. 2020, Makido et al. 2019, Yang & La Roche 2025, Helmreich et al. 2025, Oraopoulos et al. 2026).

Geographically, the studies span Asia, Europe, North America, Australia, and parts of Africa and South America, with a notable concentration in rapidly urbanizing regions of East and South Asia. This spatial distribution reflects both data availability and heightened vulnerability to extreme heat in fast-growing cities (Su et al. 2012, Lu et al. 2020, Makido et al. 2019). Several recent studies explicitly address informal settlements and marginalized urban communities, signaling a shift toward equity-oriented urban climate research (Makido et al. 2019, Oraopoulos et al. 2026). Across the dataset, studies vary substantially in spatial scale, ranging from building- and block-level analyses to city- and metropolitan-scale assessments, enabling a multi-scalar synthesis of GI–LST relationships. Most studies evaluate short-term or event-based cooling effects, while long-term performance may vary depending on vegetation growth, maintenance, and climatic variability. This highlights the need for longitudinal assessments of GI effectiveness.

#### ***3.4.1. Distribution of Green Infrastructure Types***

Across the reviewed literature, five dominant categories of GI interventions emerge: urban trees and street trees; urban parks and large green spaces; green roofs; vertical greenery systems (green walls); and blue–green infrastructure, including wetlands, vegetated swales, riverside buffers,

and stormwater retention systems (Halder & Kumar 2025). Among these, tree-based GI and urban parks are the most frequently investigated, together accounting for the majority of studies, reflecting their widespread applicability and established cooling potential across a range of urban contexts (Su et al. 2012, Makido et al. 2019, Lu et al. 2020, Di Sabatino et al. 2020, Yang & La Roche 2025).

These interventions are examined across diverse urban morphologies, from compact city centers to suburban and peri-urban environments (Kranti Kumar Maurya 2026, Pradeep Kumar Kori 2026, Rathod et al. 2026). In contrast, green roofs and vertical greenery systems are predominantly addressed in studies focused on high-density or space-constrained settings, where opportunities for ground-level greening are limited and vertical or rooftop solutions offer viable alternatives (Revell & Anda 2014, Yang & La Roche 2025). Blue-green infrastructure, while comparatively less represented in the literature, has gained increasing attention in recent years due to its multifunctional role in enhancing urban thermal regulation alongside stormwater management and broader climate adaptation benefits (Makido et al. 2019, Helmreich et al. 2025).

### 3.4.2. Magnitude and Variability of LST Reduction

The reported LST reduction ranges are derived from a comparative synthesis of values extracted from the reviewed studies (Table 2). Due to differences in reporting formats

(e.g., maximum, average, or localized cooling), these values are presented as indicative ranges rather than statistically pooled estimates. Fig. 7 presents a synthesized distribution of reported LST reductions by green infrastructure type. The values were compiled from individual studies and categorized by intervention type. Minimum, maximum, and typical reported values were extracted to construct the distribution, providing a comparative overview of variability across studies rather than a statistically derived dataset (Su et al. 2012, Revell & Anda 2014, Makido et al. 2019, Lu et al. 2020, Di Sabatino et al. 2020, Yang & La Roche 2025, Helmreich et al. 2025). The magnitude of LST reduction varies not only across GI types but also with temporal and spatial factors. Several studies report stronger cooling effects during daytime peak heat periods, while nighttime cooling tends to be more limited. Furthermore, most reported reductions refer to LST, which may not directly correspond to ambient air temperature or pedestrian-level thermal conditions.

**Tree-Based GI and Urban Parks:** Urban trees and parks consistently demonstrate the largest cooling magnitudes. Multiple remote sensing studies report mean daytime LST reductions of 2–4°C associated with high tree canopy cover, with localized reductions exceeding 6°C under favorable conditions (Su et al. 2012, Lu et al. 2020, Makido et al. 2019). Park cooling effects often extend several hundred meters into adjacent built-up areas, although the magnitude declines with distance from park boundaries.




Fig. 7: Summary of reported land surface temperature reductions by green infrastructure type. Box-and-whisker representation of minimum, median, and maximum LST reductions reported across the 61 reviewed studies, highlighting variability by intervention type.

**Green Roofs and Vertical Greenery:** Green roofs and vertical greenery systems exhibit moderate but consistent cooling, primarily at the building scale. Reported reductions typically range from 0.5 to 2.5°C, with stronger effects observed for intensive green roofs and well-irrigated systems (Revell & Anda 2014, Yang & La Roche 2025). Their contribution to neighborhood-scale LST reduction is limited but non-negligible in dense urban contexts.

**Blue–Green Infrastructure:** Blue–green infrastructure shows high variability but strong maximum cooling potential. Studies report enhanced cooling in hot–humid and hot–dry climates, where vegetation–water interactions amplify latent heat fluxes (Makido et al. 2019, Helmreich et al. 2025).

### 3.4.3. Urban Morphology and LST Measurement Approaches

Urban morphology strongly mediates GI effectiveness. In compact urban cores and street canyon environments, cooling outcomes depend on vegetation height, crown width, spacing, and interaction with built form (Di Sabatino et al. 2020, Yang & La Roche 2025). Several studies report that excessive canopy density can reduce ventilation, partially offsetting shading benefits.

In contrast, low- and medium-density urban areas benefit most from large and contiguous green spaces, which generate broader and more stable cooling footprints (Su et al. 2012, Lu et al. 2020). Significantly, studies focusing on informal settlements demonstrate that even small-scale GI interventions can significantly reduce surface heat stress and improve habitability (Makido et al. 2019, Oraipoulos et al. 2026).

Most studies estimate LST using satellite-based thermal remote sensing, primarily Landsat and MODIS platforms (Su et al. 2012, Lu et al. 2020). These approaches enable city-scale comparison and long-term trend analysis.

Simulation-based studies using ENVI-met or CFD models are frequently employed to explore GI design sensitivity and urban morphology interactions (Di Sabatino et al. 2020, Yang & La Roche 2025, Helmreich et al. 2025). Field-based measurements are mainly used for validation. Table 3 presents details of the LST assessment approaches.

Table 3: LST assessment approaches and representative studies.

| Approach                       | Key Strength             | Key Limitation                 | Representative Studies                                                                         |
|--------------------------------|--------------------------|--------------------------------|------------------------------------------------------------------------------------------------|
| Satellite-based remote sensing | City-scale comparability | Surface temperature proxy only | Su et al. 2012, Lu et al. 2020, Makido et al. 2019, Revell & Anda 2014, Oraipoulos et al. 2026 |
| Microclimate modeling          | Design optimization      | Parameter sensitivity          | Yang & La Roche 2025, Di Sabatino et al. 2020, Helmreich et al. 2025                           |
| Field measurements             | High accuracy            | Limited spatial extent         | Di Sabatino et al. 2020, Revell & Anda 2014                                                    |
| Hybrid approaches              | Strong robustness        | Methodological complexity      | Makido et al. 2019, Lu et al. 2020, Oraipoulos et al. 2026                                     |

## 4. DISCUSSION

### 4.1. Context-Dependent Effectiveness of GI for LST Reduction

This systematic review of 61 studies provides robust evidence that GI interventions are effective in reducing LST across diverse urban contexts; however, the magnitude and spatial extent of cooling are highly context-dependent. Rather than identifying a universally optimal GI solution, the synthesized literature consistently demonstrates that cooling performance emerges from the interaction between GI type, urban morphology, vegetation characteristics, and climate conditions (Su et al. 2012, Revell & Anda 2014, Makido et al. 2019, Lu et al. 2020, Di Sabatino et al. 2020, Yang & La Roche 2025, Helmreich et al. 2025, Oraipoulos et al. 2026).

Tree-based GI and urban parks repeatedly exhibit the strongest and most spatially extensive cooling effects, largely due to the combined influence of shading and evapotranspiration (Kumar et al. 2025). Across multiple climatic regions and urban forms, dense tree canopy cover is associated with mean daytime LST reductions of 2–4°C, with localized reductions exceeding 6°C under favorable conditions (Su et al. 2012, Lu et al. 2020, Makido et al. 2019). Urban parks act as thermal sinks, generating cooling footprints that extend into surrounding built-up areas, although the intensity of cooling decreases with distance from park boundaries.

In contrast, green roofs and vertical greenery systems produce more localized cooling effects, typically limited to roof surfaces and building façades. While these interventions rarely generate large neighborhood-scale LST reductions, they play a critical role in high-density urban environments, where horizontal greening opportunities are constrained (Revell & Anda 2014, Yang & La Roche 2025). Blue–green infrastructure, although less frequently examined, demonstrates strong cooling potential, particularly in hot–dry and hot–humid climates where vegetation–water interactions enhance latent heat fluxes (Makido et al. 2019, Helmreich et al. 2025). Taken together, these findings underscore that GI effectiveness cannot be inferred solely from green area coverage. Instead, cooling outcomes

depend on how, where, and under what climatic conditions GI is implemented. The findings demonstrate that GI effectiveness cannot be generalized across contexts. Cooling performance varies with time of day, spatial scale, and measurement approach, with most studies focusing on surface temperature reductions rather than direct human thermal conditions. These distinctions are critical for interpreting the practical implications of GI interventions. The geographic concentration of studies also limits the global generalizability of findings. Variations in climate zones (e.g., hot–arid vs. temperate), urban development patterns, and vegetation types can lead to significantly different cooling outcomes, reinforcing the need for region-specific assessments. For example, the strong cooling effects reported for tree-based GI are largely derived from studies conducted in specific climatic contexts, which may not fully represent performance in arid or resource-constrained environments.

#### 4.2. Urban Morphology, Design Trade-offs, and Cooling Mechanisms

Urban morphology emerges as one of the most influential moderators of GI cooling performance. Studies conducted in compact city centers and street canyon environments consistently report that vegetation height, crown geometry, spacing, and interaction with built form critically shape

cooling outcomes (Di Sabatino et al. 2020, Yang & La Roche 2025). While shading provided by trees reduces surface temperatures, poorly configured vegetation can restrict airflow, reducing convective heat removal and partially offsetting cooling benefits.

In lower-density urban areas, where space allows for larger and more continuous green spaces, GI produces broader and more stable cooling footprints. Remote sensing studies repeatedly demonstrate that large parks and green belts generate LST reductions extending several hundred meters into adjacent neighborhoods (Su et al. 2012, Lu et al. 2020). These spatial gradients highlight the importance of connectivity and spatial configuration, rather than isolated green patches. Fig. 8 shows a conceptual illustration of how dominant cooling mechanisms like shading, evapotranspiration, and vegetation–water interactions vary across hot–dry, hot–humid, and temperate climates that influence the effectiveness of green infrastructure interventions in reducing land surface temperature.

Importantly, the reviewed literature shows that small-scale GI interventions can deliver substantial benefits in spatially constrained environments, including informal settlements. Oraipoulos et al. (2026) demonstrate that targeted interventions such as roadside trees and micro-parks significantly reduce surface heat stress and improve




Fig. 8: Climate-dependent pathways of green infrastructure cooling.

habitability in dense informal settlements, despite severe limitations in land availability. These findings challenge assumptions that effective GI cooling requires large-scale or high-cost interventions and emphasize the importance of design-sensitive, context-aware planning.

### 4.3. Climatic Sensitivity, Vegetation Characteristics, and Blue-Green Synergies

The cooling effectiveness of GI is strongly climate-sensitive, with consistently higher LST reductions reported in hot–dry and hot–humid regions compared to temperate climates (Makido et al. 2019, Helmreich et al. 2025). In these climates, evapotranspiration plays a dominant role in surface energy balance, amplifying the cooling impact of vegetation.

Vegetation characteristics further mediate cooling performance. Multiple studies highlight the importance of canopy density, leaf area index, and species selection, with tree-based systems consistently outperforming grass-dominated GI in terms of surface cooling (Lu et al. 2020, Yang & La Roche 2025). However, several studies caution that cooling benefits decline under drought stress, particularly in arid regions where water availability limits evapotranspiration (Makido et al. 2019). Blue–green infrastructure enhances cooling by coupling vegetation with water-mediated thermal regulation, but its performance is highly dependent on hydrological conditions and maintenance. While such systems offer substantial potential for heat mitigation, their long-term effectiveness under future climate change scenarios remains uncertain, highlighting a critical area for future research.

### 4.4. Methodological Implications, Equity, and Future Directions

The reviewed studies employ a diverse range of methodological approaches, with satellite-based LST retrieval using Landsat and MODIS dominating the literature (Su et al. 2012, Lu et al. 2020). These methods enable consistent city-scale analysis but do not directly capture human thermal comfort. Simulation-based approaches, such as ENVI-met and CFD modeling, provide valuable insights into design optimization and causal mechanisms but remain sensitive to parameterization and require empirical validation (Di Sabatino et al. 2020, Yang & La Roche 2025, Helmreich et al. 2025).

A key strength of recent literature is the growing focus on heat vulnerability and equity, particularly in informal settlements and marginalized urban communities. Evidence increasingly shows that even modest GI interventions can yield disproportionate benefits in high-risk contexts (Makido et al. 2019, Oraiopoulos et al. 2026). However, significant

gaps remain, including limited long-term assessments of GI performance, inconsistent classification of urban morphology, and insufficient integration of LST analysis with human thermal comfort and social vulnerability metrics.

A key limitation of the reviewed literature is the reliance on LST as a primary indicator of thermal performance. While LST provides valuable insights into surface heat patterns, it does not directly represent human thermal comfort. Future research should integrate LST with air temperature measurements and human thermal indices (e.g., PET, UTCI) to provide a more comprehensive assessment of urban heat mitigation strategies.

This review advances understanding from generalized assertions that “green infrastructure cools cities” toward a context-sensitive, design-aware, and climate-responsive framework. The evidence from 61 studies demonstrates that GI can substantially reduce land surface temperature, but only when interventions are carefully aligned with urban form, vegetation characteristics, and climatic conditions. By systematically synthesizing results through the PRISMA framework, this study provides a transparent and reproducible evidence base to inform urban heat mitigation strategies that are both effective and equitable.

## 5. CONCLUSIONS

This PRISMA-based systematic review synthesizes evidence from 61 peer-reviewed studies to evaluate how different green infrastructure (GI) interventions influence land surface temperature (LST) across diverse built environments. The findings indicate that GI is an effective strategy for reducing surface heat exposure; however, its performance is strongly context-dependent, varying with urban morphology, vegetation characteristics, and climatic conditions. Across the reviewed literature, tree-based GI and urban parks consistently demonstrate the most substantial and spatially extensive reductions in LST, primarily driven by shading and evapotranspiration processes. These effects are generally more pronounced during daytime peak heat periods. In contrast, green roofs and vertical greenery systems provide more localized cooling benefits, particularly in high-density urban areas where horizontal greening is constrained. Blue–green infrastructure exhibits strong cooling potential in certain climatic contexts, especially where vegetation–water interactions enhance latent heat exchange, although performance is influenced by water availability and maintenance conditions.

The review further highlights that the effectiveness of GI interventions is mediated by urban form and spatial configuration. Large and contiguous green spaces tend to

produce broader and more stable cooling effects in low- and medium-density environments, whereas compact urban areas require carefully integrated design strategies that balance shading, ventilation, and spatial constraints. Evidence from informal settlements suggests that small-scale, strategically implemented GI interventions can contribute to reducing localized surface heat exposure, with potential implications for improving thermal conditions in heat-vulnerable areas, although systematic evidence on equity outcomes remains limited.

Methodologically, the literature is characterized by a predominance of satellite-based LST assessments, complemented by simulation models and field measurements. While these approaches provide valuable insights into spatial patterns of surface temperature, the review identifies key limitations, including methodological heterogeneity, limited long-term evaluation, and insufficient integration of LST-based analysis with human-centered thermal comfort indicators and socio-environmental factors.

Overall, this review advances understanding from generalized assertions of GI effectiveness toward a more context-sensitive and design-oriented perspective on urban heat mitigation. By systematically organizing evidence across intervention types, urban forms, and climatic conditions, the study provides a transparent foundation for interpreting GI performance in different settings. Future research should prioritize longitudinal and comparative studies, expand representation across underrepresented regions, and integrate LST analysis with air temperature measurements and human thermal comfort indices (e.g., PET, UTCI) to support more comprehensive and human-centered assessments. Such efforts will strengthen the role of green infrastructure as a key component of climate-responsive and resilient urban planning.

## 6. ACKNOWLEDGMENTS

The authors would like to express their sincere gratitude to all the researchers whose work was reviewed and synthesized in this study. Their valuable contributions have shaped the knowledge base surrounding LST and green infrastructure. We also acknowledge the use of VOSviewer software for the bibliometric analysis, which provided critical insights into thematic trends and research networks. Special thanks are extended to the institutions and libraries that provided access to academic databases, making the systematic literature review possible. Finally, the authors appreciate the support and feedback received from colleagues and academic peers throughout the development of this manuscript.

## 7. REFERENCES

Abuwaer, N., Ullah, S. and Al-Ghamdi, S.G., 2023. Building Climate Resilience Through Urban Planning: Strategies, Challenges, and

- Opportunities. In: *Sustainable Cities in a Changing Climate*, pp.185–206. Wiley. [DOI]
- Al-Zghoul, S. and Al-Homoud, M., 2025. GIS-Driven Spatial Planning for Resilient Communities: Walkability, Social Cohesion, and Green Infrastructure in Peri-Urban Jordan. *Sustainability*, 17(14). [DOI]
- Amani-Beni, M., Zhang, B., Xie, G.-D. and Odgaard, A.J., 2021. Impacts of the Microclimate of a Large Urban Park on Its Surrounding Built Environment in the Summertime. *Remote Sensing*, 13(22). [DOI]
- Anderson, V., Gough, W.A., Zgela, M., Milosevic, D. and Dunjic, J., 2022. Lowering the Temperature to Increase Heat Equity: A Multi-Scale Evaluation of Nature-Based Solutions in Toronto, Ontario, Canada. *Atmosphere*, 13(7). [DOI]
- Arruda, H., 2022. VOSviewer and Bibliometrix. *Journal of the Medical Library Association : JMLA*, 110(3), pp.392–395. [DOI]
- Azmeer, A., Tahir, F. and Al-Ghamdi, S.G., 2024. Progress on green infrastructure for urban cooling: Evaluating techniques, design strategies, and benefits. *Urban Climate*, 56. [DOI]
- Bachofen, C., Peillon, M., Meili, N., Bourgeois, I. and Grossiord, C., 2025. High transpirational cooling by urban trees despite extreme summer heatwaves. *Urban Forestry & Urban Greening*, 107. [DOI]
- Back, Y., Bach, P.M., Jasper-Toennies, A., Rauch, W. and Kleidorfer, M., 2021. A rapid fine-scale approach to modelling urban bioclimatic conditions. *Science of the Total Environment*, 756. [DOI]
- Battista, G., Carnielo, E. and De Lieto Vollaro, R., 2016. Thermal impact of a redeveloped area on localized urban microclimate: A case study in Rome. *Energy and Buildings*, 133, pp.446–454. [DOI]
- Bera, M. and Nag, P.K., 2025. AN OVERVIEW OF BIOCLIMATIC DESIGN STRATEGIES OF THE BUILT ENVIRONMENT. *Environmental Engineering and Management Journal*, 24(4). [DOI]
- Berger, N.K., Kollmann, J. and Teixeira, L.H., 2026. Native plant communities designed for infiltration swales can tolerate urban heat, flooding, and pollution. *Basic and Applied Ecology*, 90, pp.89–98. [DOI]
- Bouzouidja, R., Cannavo, P., Bodenan, P., Gulyas, A., Kiss, M., Kovacs, A., Bechet, B., Chancibault, K., Chantoiseau, E., Bournet Pierre-Emmanuel and Bouzidi, R., Guenon, R., Lebeau, T., Musy, M. and Rodriguez, F., 2021. How to evaluate nature-based solutions performance for microclimate, water and soil management issues - Available tools and methods from Nature4Cities European project results. *Ecological Indicators*, 125. [DOI]
- Bowen, X. Hui, L., Jingmao, W. and Shuo, C. 2024. Ecological design approach for urban green space in arid and semi-arid areas in Northwest China based on habitat and vegetation characteristics thereof. *Landscape Architecture*, 31(6), pp.12-18. [DOI]
- Chaudhuri, S. and Kumar, A., 2022. Urban greenery for air pollution control: a meta-analysis of current practice, progress, and challenges. *Environmental Monitoring and Assessment*. [DOI]
- Chen, D., Zhang, F., Zhang, M., Meng, Q., Jim, C.Y., Shi, J., Tan, M.L. and Ma, X., 2022. Landscape and vegetation traits of urban green space can predict local surface temperature. *Science of the Total Environment*, 825, p.154006. [DOI]
- Chen, M., Dang, A., Li, X., Huang, J. and Weng, Y., 2024. Identifying climate-ready plants for urban environment: Integrating machine learning with traditional plant selection tools. *Urban Forestry & Urban Greening*, 102. [DOI]
- Chen, P. and Hanlon, B., 2025. Urban green space, respiratory health and rising temperatures: An examination of the complex relationship between green space and adult asthma across racialized neighborhoods in Los Angeles County. *Landscape and Urban Planning*, 258. [DOI]
- Chen, Y., Wang, Y. and Zhou, D., 2021. Knowledge map of urban morphology and thermal comfort: A bibliometric analysis based on CiteSpace. *Buildings*, 11(10), p.427. [DOI]
- Chen, Y., Zhang, R., Alekouei, S.A. and Amani-Beni, M., 2024. Nonlinear impacts of landscape and climatological interactions on urban thermal

- environment during a hot and rainy summer. *Ecological Indicators*, 166. [DOI]
- Cheung, P.K. and Jim, C.Y., 2018. Subjective outdoor thermal comfort and urban green space usage in humid-subtropical Hong Kong. *Energy and Buildings*, 173, pp.150-162. [DOI]
- Cheung, P.K., Livesley, S.J. and Nice, K.A., 2021. Estimating the cooling potential of irrigating green spaces in 100 global cities with arid, temperate or continental climates. *Sustainable Cities and Society*, 71. [DOI]
- Chow, W.T., Akbar, S.N.A.B.A., Heng, S.L. and Roth, M., 2016. Assessment of measured and perceived microclimates within a tropical urban forest. *Urban Forestry & Urban Greening*, 16, pp.62-75. [DOI]
- Chu, Z., Li, S., Li, T., Qian, H. and Liu Chuan and Yan, Z., 2024. Numerical simulation of layout and landscape elements on the thermal environment of urban squares. *Ecological Informatics*, 82. [DOI]
- Chun, B., Hur, M. and Won, J., 2021. Impacts of Thermal Environments on Health Risk: A Case Study of Harris County, Texas. *International Journal Of Environmental Research And Public Health*, 18(11). [DOI]
- Cuce, P.M., Cuce, E. and Santamouris, M., 2025. Towards Sustainable and Climate-Resilient Cities: Mitigating Urban Heat Islands Through Green Infrastructure. *Sustainability*, 17(3). [DOI]
- de Oliveira, J.A.P., Bellezoni, R.A., Shih, W.Y. and Bayulken, B., 2022. Innovations in Urban Green and Blue Infrastructure: Tackling local and global challenges in cities. *Journal of Cleaner Production*, 362, p.132355. [DOI]
- De Quadros, B.M. and Mizgier, M.G.O., 2023. Urban green infrastructures to improve pedestrian thermal comfort: A systematic review. *Urban Forestry & Urban Greening*, 88, p.128091. [DOI]
- Del Rosario, L., Astell-Burt, T., Olsen, J.R., Thompson, S., Mitchell, R. and Feng, X., 2025. Walkable by nature? Adding green and blue space to walkability indices and assessment of socioeconomic inequalities. *Environmental Research*, 284. [DOI]
- Derkzen, M.L., van Teeffelen, A.J.A. and Verburg, P.H., 2017. Green infrastructure for urban climate adaptation: How do residents' views on climate impacts and green infrastructure shape adaptation preferences? *Landscape and Urban Planning*, 157, 106-130. [DOI]
- Dybå, T. and Dingsøyr, T., 2008. Empirical studies of agile software development: A systematic review. *Information and Software Technology*, 50(9-10), pp.833-859. [DOI]
- Esperon-Rodriguez, M., Sharmin, M., Rodriguez, D.E., Messier, C., Svenning, J.-C., Moore, S. and Tjoelker, M.G., 2025. Socio-economic factors, climate, and people's behaviours determine urban tree health. *Urban Forestry & Urban Greening*, 107. [DOI]
- Evola, G., Gagliano, A., Fichera, A., Marletta, L., Martinico, F., Nocera, F. and Pagano, A., 2017. UHI effects and strategies to improve outdoor thermal comfort in dense and old neighbourhoods. *Energy Procedia*, 134, pp.692-701. [DOI]
- Fang, X., Li, J. and Ma, Q., 2023. Integrating green infrastructure, ecosystem services and nature-based solutions for urban sustainability: A comprehensive literature review. *Sustainable Cities and Society*, 98, p.104843. [DOI]
- Feng, S., 2022. Land Use/Land Cover Mapping Based on GEE for the Monitoring of Changes in Ecosystem Types in the Upper Yellow River Basin over the Tibetan Plateau. *Remote Sensing*, 14(21). [DOI]
- Fong, C.S., Manavvi, S., Priya, R.S., Ramakreshnan, L., Sulaiman, N.M. and Aghamohammadi, N., 2023. Traits of Adaptive Outdoor Thermal Comfort in a Tropical Urban Microclimate. *Atmosphere*, 14(5). [DOI]
- Frosini, G., Amato, A., Mugnai, F. and Cinelli, F., 2024. The Impact of Trees on the UHI Effect and Urban Environment Quality: A Case Study of a District in Pisa, Italy. *Atmosphere*, 15(1). [DOI]
- Fu, J., Dupre, K., Tavares, S., King, D. and Banhalmi-Zakar, Z., 2022. Optimized greenery configuration to mitigate urban heat: A decade systematic review. *Frontiers of Architectural Research*, 11(3), pp.466-491. [DOI]
- Fuentes, S., Tongson, E. and Gonzalez Viejo, C., 2021. Urban Green Infrastructure Monitoring Using Remote Sensing from Integrated Visible and Thermal Infrared Cameras Mounted on a Moving Vehicle. *Sensors*, 21(1). [DOI]
- Gherri, B., 2023. The Role of Urban Vegetation in Counteracting Overheating in Different Urban Textures. *Land*, 12(12). [DOI]
- Gorai, N., Bandyopadhyay, J., Halder, B., Ahmed, M.F., Molla, A.H. and Lei, T.M.T., 2024. Spatio-Temporal Variation in Landforms and Surface Urban Heat Island in Riverine Megacity. *Sustainability*, 16(8), p.3383. [DOI]
- Graef, M., Pucher, B., Hietz, P., Hofbauer, K., Allabashi, R., Pitha, U., Hood-Nowotny, R. and Stangl, R., 2022. Application of leaf analysis in addition to growth assessment to evaluate the suitability of greywater for irrigation of *Tilia cordata* and *Acer pseudoplatanus*. *Science of the Total Environment*, 836. [DOI]
- Guan, Z. and Zhang, H., 2024. A Quantitative Analysis of the Complex Response Relationship between Urban Green Infrastructure (UGI) Structure/Spatial Pattern and Urban Thermal Environment in Shanghai. *Sustainability*, 16(16). [DOI]
- Halder, N. and Kumar, D., 2025. Towards Reviving Vernacular Architecture in India: Insights into the Indigenous Building Traditions. *Journal of the International Society for the Study of Vernacular Settlements (ISVS e-Journal)*, 12(5), pp.40-54. [DOI]
- Halder, N., Kumar, M., Deepak, A., Abuwaer, N.M., Kanwal, Q., Nurdiawati, A. and Al-Ghamdi, S.G., 2026. Urban Greenery and Outdoor Thermal Comfort: Cooling Indicators, Mechanisms, and Equity Implications. *Environmental Research Communications*. [DOI]
- Halder, N., Kumar, M., Deepak, A., Kumar, D., Yasmin, N., Ullah, S. and Al-Ghamdi, S.G., 2025. Spatiotemporal Assessment of Urban Thermal Discomfort in Kolkata, India: Insights from Cloud-Based Remote Sensing. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 1-17. [DOI]
- Halder, N., Kumar, M., Deepak, A., Mandal, S.K., Azmeer, A., Mir, B.A., Nurdiawati, A. and Al-Ghamdi, S.G., 2025. The Role of Urban Greenery in Enhancing Thermal Comfort: Systematic Review Insights. *Sustainability*, 17(6), p.2545. [DOI]
- Helmreich, B., Deeb, M., Eben, P., Egerer Monika and Kollmann, J., Schulz, S., Pauleit, S., Weisser, W.W. and Schlöter, M., 2025. Converting infiltration swales to sustainable urban drainage systems can improve water management and biodiversity. *Frontiers in Environmental Science*, 12. [DOI]
- Hogewij, M., Ploegmakers, H., Raaphorst, K., Bongers, C. and van der Krabben, E., 2025. It's not easy being cool: Residents' perceived cooling ability in relation to different types of green infrastructure in their neighbourhood. *Urban Forestry & Urban Greening*, 104. [DOI]
- Huang, C., Liu, K., Ma, T., Xue, H., Wang, P. and Li, L., 2025. Analysis of the impact mechanisms and driving factors of urban spatial morphology on urban heat islands. *Scientific Reports*, 15(1), p.18589. [DOI]
- Huang, J., Hao, T., Wang, Y. and Jones, P., 2022. A street-scale simulation model for the cooling performance of urban greenery: Evidence from a high-density city. *Sustainable Cities and Society*, 82, p.103908. [DOI]
- Hunter, R.F., Christian, H., Veitch, J., Astell-Burt, T., Hipp, J.A. and Schipperijn, J., 2015. The impact of interventions to promote physical activity in urban green space: a systematic review and recommendations for future research. *Social Science & Medicine*, 124, pp.246-256. [DOI]
- Hurdud, A., Ermida, S.L. and DaCamara, C.C., 2024. On the Suitability of Different Satellite Land Surface Temperature Products to Study Surface Urban Heat Islands. *Remote Sensing*, 16(20), p.3765. [DOI]
- Ibric, A., Bostenaru Dan, M. and Craciun, C., 2024. Adapting Street Profile Design by Using Nature-Based Solutions in New Neighbourhoods and the Retrofit of Buildings. *Buildings*, 14(7). [DOI]
- Irfeey, A.M.M., Chau, H.-W., Sumaiya, M.M.F., Wai, C.Y., Muttill, N. and Jamei, E., 2023. Sustainable Mitigation Strategies for Urban Heat Island Effects in Urban Areas. *Sustainability*, 15(14). [DOI]

- Jiang, Y., Huang, J., Shi, T. and Wang, H., 2021. Interaction of urban rivers and green space morphology to mitigate the urban heat island effect: Case-based comparative analysis. *International Journal of Environmental Research and Public Health*, 18(21). [DOI]
- Jimenez, J.L.S. and de Adana, M.R., 2024. Assessment of Outdoor Thermal Comfort in a Hot Summer Region of Europe. *Atmosphere*, 15(2). [DOI]
- Jin, J., Chen, W.Y., Jia, B. and Wang, C., 2024. Cooling effect of urban greenery: A bibliometric analysis. *Urban Forestry & Urban Greening*, 99, p.128453. [DOI]
- Kabano, P., Harris, A. and Lindley, S., 2021. Sensitivity of Canopy Phenology to Local Urban Environmental Characteristics in a Tropical City. *Ecosystems*, 24(5), pp.1110–1124. [DOI]
- Kanga, S., Meraj, G., Johnson, B.A., Singh, S.K., PV, M.N., Farooq, M., Kumar, P., Marazi, A. and Sahu, N., 2022. Understanding the Linkage between Urban Growth and Land Surface Temperature—A Case Study of Bangalore City, India. *Remote Sensing*, 14(17), p.4241. [DOI]
- Karimi, A., Moreno-Rangel, D. and Garcia-Martinez, A., 2025. Granular mapping of UHI and heatwave effects: Implications for building performance and urban resilience. *Building and Environment*, 273. [DOI]
- Kikon, N., Kumar, D. and Ahmed, S.A., 2023. Quantitative assessment of land surface temperature and vegetation indices on a kilometer grid scale. *Environmental Science and Pollution Research*, 30(49), pp.107236–107258. [DOI]
- Kirby, A., 2023. Exploratory Bibliometrics: Using VOSviewer as a Preliminary Research Tool. *Publications*, 11(1). [DOI]
- Kotharkar, R. and Bagade, A., 2018. Evaluating urban heat island in the critical local climate zones of an Indian city. *Landscape and Urban Planning*, 169, pp.92–104. [DOI]
- Kranti Kumar Maurya, D.K., 2026. Towards Net Zero: A Review of Embodied Carbon in Residential Building Construction. In: Fulena Rajak, Ajay Kumar, Bijay Kumar Das, Ravish Kumar, Kranti Kumar Maurya, Sandeep Kumar, Shubham Meena, Abhishek Kumar Bittu, & Preksha Gupta (Eds.), *International Conference on Urban Resilience and Sustainable Architecture*, pp.86–91. Department of Architecture and Planning, NIT Patna.
- Kumar, D., Maurya, K.K., Mandal, S.K., Halder, N., Mir, B.A., Nurdiawati, A. and Al-Ghamdi, S.G., 2025. A Whole-Life Carbon Assessment of a Single-Family House in North India Using BIM-LCA Integration. *Buildings*, 15(13), p.2195. [DOI]
- Kumar, D., Maurya, K.K., Mandal, S.K., Mir, B.A., Nurdiawati, A. and Al-Ghamdi, S.G., 2025. Life Cycle Assessment in the Early Design Phase of Buildings: Strategies, Tools, and Future Directions. *Buildings*, 15(10), p.1612. [DOI]
- Kumar, D. and Shukla, B., 2022. Urban Green Spaces For Promoting Healthy Living And Wellbeing: Prospects For Housing. *ECS Transactions*, 107(1), pp.18835–18857. [DOI]
- La Rosa, D. and Li, J., 2023. High-Resolution Greening Scenarios for Urban Climate Regulation Based on Physical and Socio-Economical Factors. *Sustainability*, 15(9). [DOI]
- Lazovic, I.M., Nikezic, D.P., Markovic, Z.J., Eric, M., Zivkovic, M., Ramadani, U., Tasic, G. and Tasic, V., 2025. Assessment of Traffic-Induced Air Pollution and Its Effects on Intensity of Urban Heat Islands. *Applied Sciences-Basel*, 15(20). [DOI]
- Lee, L.S. and Jim, C.Y., 2018. Thermal-cooling performance of subtropical green roof with deep substrate and woodland vegetation. *Ecological Engineering*, 119, pp.8-18. [DOI]
- Li, J., Mao, Y., Ouyang, J. and Zheng, S., 2022. A review of urban microclimate research based on CiteSpace and VOSviewer analysis. *International Journal of Environmental Research and Public Health*, 19(8), p.4741. [DOI]
- Liu, H., Zhou, Z., Wen, Q., Chen, J. and Kojima, S., 2024. Spatiotemporal Land Use/Land Cover Changes and Impact on Urban Thermal Environments: Analyzing Cool Island Intensity Variations. *Sustainability*, 16(8). [DOI]
- Liu, J., Wu, X., Pan, L. and Hsieh, C.-M., 2025. Multi-Scale Analysis of the Mitigation Effect of Green Space Morphology on Urban Heat Islands. *Atmosphere*, 16(7). [DOI]
- Liu, Z., Cheng, K.Y., Sinsel, T., Simon, H., Jim, C.Y., Morakinyo, T.E., He, Y., Yin, S., Ouyang, W., Shi, Y. and Ng, E., 2023. Modeling microclimatic effects of trees and green roofs/façades in ENVI-met: Sensitivity tests and proposed model library. *Building and Environment*, 244. [DOI]
- Liu, Z., Cheng, W., Jim, C.Y., Morakinyo, T.E., Shi, Y. and Ng, E., 2021. Heat mitigation benefits of urban green and blue infrastructures: A systematic review of modeling techniques, validation and scenario simulation in ENVI-met V4. *Building and Environment*, 200, p.107939. [DOI]
- Lu, L., Weng, Q., Xiao, D., Guo, H., Li, Q. and Hui, W., 2020. Spatiotemporal Variation of Surface Urban Heat Islands in Relation to Land Cover Composition and Configuration: A Multi-Scale Case Study of Xi'an, China. *Remote Sensing*, 12(17). [DOI]
- Luo, H. and Wu, J., 2021. Effects of urban growth on the land surface temperature: a case study in Taiyuan, China. *Environment, Development and Sustainability*, 23(7), pp.10787–10813. [DOI]
- Makido, Y., Hellman, D. and Shandas, V., 2019. Nature-Based Designs to Mitigate Urban Heat: The Efficacy of Green Infrastructure Treatments in Portland, Oregon. *Atmosphere*, 10(5). [DOI]
- Mittermüller, J., Erlwein, S., Bauer, A., Trokai, T., Duschinger, S. and Schönmann, M., 2021. Context-specific, user-centred: Designing urban green infrastructure to effectively mitigate urban density and heat stress. *Urban Planning*, 6(4), pp.40–53. [DOI]
- Mohan, M. and Kandya, A., 2015. Impact of urbanization and land-use/land-cover change on diurnal temperature range: A case study of tropical urban airshed of India using remote sensing data. *Science of The Total Environment*, 506–507, 453–465. [DOI]
- Nath, R., 2025. Modeling Urban Thermal Dynamics and Land Use Change in Guwahati: A Machine Learning-Based Analysis of Lst and Urfvi Trends (2000–2040). [DOI]
- Ni, Y., Lin, L., Xia, H. and Wang, X., 2025. Urban Street Greening and Resident Comfort: An Integrated Approach Based on High-Precision Shadow Distribution and Facade Visual Assessment. *Sustainability*, 17(3). [DOI]
- O'Donnell, E.C., Lamond, J.E. and Thorne, C.R., 2017. Recognising barriers to implementation of Blue-Green Infrastructure: a Newcastle case study. *Urban Water Journal*, 14(9), pp.964–971. [DOI]
- O'Keeffe, J., Pluchinotta, I., De Stercke, S., Hinson, C., Puchol-Salort, P., Mijic, A., Zimmermann, N. and Collins, A.M., 2022. Evaluating natural capital performance of urban development through system dynamics: A case study from London. *Science of the Total Environment*, 824. [DOI]
- Okumus, D.E. and Terzi, F., 2023. Ice floes in urban furnace: Cooling services of cemeteries in regulating the thermal environment of Istanbul's urban landscape. *Urban Climate*, 49. [DOI]
- Oraiopoulos, A., Wieser, M., Verdieri, M., Lambert, R., Fennell, P. and Ruyssevelt, P., 2026. Improving habitability in informal settlements in the Global South: Exploring the impact of community urban green infrastructure on outdoor heat stress. *Building and Environment*, 287(A). [DOI]
- Ornelas, A., Cordeiro, A. and Lameiras, J.M., 2023. Thermal comfort assessment in urban green spaces: Contribution of thermography to the study of thermal variation between tree canopies and air temperature. *Land*, 12(8), p.1568. [DOI]
- Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E. and Chou, R., 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *bmj*, 372. [DOI]

- Pandey, B., Banerjee, R. and Sharma, A., 2021. Coupled EnergyPlus and CFD analysis of PCM for thermal management of buildings. *Energy and Buildings*, 231, p.110598. [DOI]
- Pattanaik, N., Rahman, M.A. and Pauleit, S., 2025. Urban vegetation structure and composition influence pedestrian thermal comfort in public spaces: A case study in Munich. *Urban Forestry & Urban Greening*, 114. [DOI]
- Perini, K. and Magliocco, A., 2014. Effects of vegetation, urban density, building height, and atmospheric conditions on local temperatures and thermal comfort. *Urban Forestry & Urban Greening*, 13(3), pp.495-506. [DOI]
- Pradeep Kumar, B., Anusha, B.N., Raghu Babu, K. and Padma Sree, P., 2023. Identification of climate change impact and thermal comfort zones in semi-arid regions of AP, India using LST and NDBI techniques. *Journal of Cleaner Production*, 407, p.137175. [DOI]
- Pradeep Kumar Kori, K.K.M.D.K., 2026. Towards Net Zero: A Systematic Review of Operational Emissions in Residential Building Construction. In: Fulena Rajak, Ajay Kumar, Bijay Kumar Das, Ravish Kumar, Kranti Kumar Maurya, Sandeep Kumar, Shubham Meena, Abhishek Kumar Bittu, & Preksha Gupta (Eds.), *International Conference on Urban Resilience and Sustainable Architecture*, pp.148-153. Department of Architecture and Planning, NIT Patna.
- Pragati, S., Priya, R.S., Pradeepa, C. and Senthil, R., 2023. Simulation of the Energy Performance of a Building with Green Roofs and Green Walls in a Tropical Climate. *Sustainability*, 15(3). [DOI]
- Pritipadmaja, Garg, R.D. and Sharma, A.K., 2023. Assessing the Cooling Effect of Blue-Green Spaces: Implications for Urban Heat Island Mitigation. *Water*, 15(16). [DOI]
- Rahman, Md. M., Avtar, R., Yunus, A.P., Dou, J., Misra, P., Takeuchi, W., Sahu, N., Kumar, P., Johnson, B.A., Dasgupta, R., Kharrazi, A., Chakraborty, S. and Agustiono Kurniawan, T., 2020. Monitoring Effect of spatial growth on land surface temperature in Dhaka. *Remote Sensing*, 12(7), p.1191. [DOI]
- Ramadan, M.S., Tariq, A., Abdo, H.G. and Al Hosani, N., 2025. GEE-based analysis of spatiotemporal patterns of land cover and LST in an arid urban environment a case study of Abu Dhabi. *Frontiers in Environmental Science*, 13. [DOI]
- Rathod, V., Rupapara, S., Agrawal, H., Manek, P., Halder, N. and Kumar, D., 2026. Nature-Based Solutions vs Conventional Infrastructure: Cost-benefit and Sustainability Evaluations in Built Environments. *Asian Journal of Environment & Ecology*, 25(3), pp.169-200. [DOI]
- Revell, G. and Anda, M., 2014. Sustainable Urban Biophilia: The Case of Greenskins for Urban Density. *Sustainability*, 6(8), pp.5423-5438. [DOI]
- Reyhani, M., Santolini, E., Torreggiani, D. and Tassinari, P., 2022. Assessing the environmental performance of plastic-based and felt-based green wall systems in a life-cycle perspective. *Science of the Total Environment*, 822. [DOI]
- Roversi, R. and Longo, D., 2025. Regenerative and Connective Green Cells to Address Fragmentation and Climate Change in Cities: The TALEA Project Integrated Solution. *Sustainability*, 17(7). [DOI]
- Sabatino, S. Di, Barbano, F., Brattich, E. and Pulvirenti, B., 2020. The Multiple-Scale Nature of Urban Heat Island and Its Footprint on Air Quality in Real Urban Environment. *Atmosphere*, 11(11). [DOI]
- Salvalai, G., Cadena, J.D.B., Sparvoli Gessica and Bernardini, G. and Quagliarini, E., 2022. Pedestrian Single and Multi-Risk Assessment to SLODs in Urban Built Environment: A Mesoscale Approach. *Sustainability*, 14(18). [DOI]
- Schaffernicht, S.K., Tuerk, A., Kogler, M., Berger, A., Scharf, B., Clementschitsch, L., Hammer, P., Holzer, P., Formayer, H., Koenig, B. and Haluza, D., 2023. Heat vs. Health: Home Office under a Changing Climate. *Sustainability*, 15(9). [DOI]
- Seidu, S., Edwards, D.J., Owusu-Manu, D., Chan, D.W.M., Abdulai, S.F. and Buerter, J.I.T., 2025. Achieving multifunctionality in green infrastructure projects: a fuzzy evaluation and Gini index of Key drivers in developing countries. *Environment Development And Sustainability*. [DOI]
- Seto, K.C., Güneralp, B. and Hutyra, L.R., 2012. Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences*, 109(40), pp.16083-16088. [DOI]
- Shan, Z., An, Y., Xu, L. and Yuan, M., 2022. High-Temperature Disaster Risk Assessment for Urban Communities: A Case Study in Wuhan, China. *International Journal Of Environmental Research And Public Health*, 19(1). [DOI]
- Shao, H., Kim, G., Li, Q. and Newman, G., 2021. Web of science-based green infrastructure: a bibliometric analysis in CiteSpace. *Land*, 10(7), p.711. [DOI]
- Shin, Y., Kim, S., Lee, S.-W. and An, K., 2020. Identifying the Planning Priorities for Green Infrastructure within Urban Environments Using Analytic Hierarchy Process. *Sustainability*, 12(13). [DOI]
- Sobrinho, O.M., Rudke, A.P., de Moraes, M.V.B. and Martins, L.D., 2023. Meteorological Effects of Green Infrastructure on a Developing Medium Latin American City: A Numerical Modeling Assessment. *Sustainability*, 15(2). [DOI]
- Su, W., Yang, G., Chen, S. and Yang, Y., 2012. Measuring the Pattern of High Temperature Areas in Urban Greenery of Nanjing City, China. *International Journal of Environmental Research and Public Health*, 9(8), pp.2922-2935. [DOI]
- Rupapara, S., Rathod, V., Rupapara, H., Halder, N. and Kumar, D., 2025. Evaluating the benefits of urban greenery in Urban Heat Island mitigation: Methods, indicators and gaps. *Nature Environment and Pollution Technology*, 24(4), p.D1812. [DOI]
- Toparlar, Y., Blocken, B., Maiheu, B. and van Heijst, G.J.F., 2017. A review on the CFD analysis of urban microclimate. *Renewable and Sustainable Energy Reviews*, 80, pp.1613-1640. [DOI]
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kaźmierczak, A., Niemela, J. and James, P., 2007. Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and Urban Planning*, 81(3), pp.167-178. [DOI]
- Ullah, S., Aldossary, A., Ullah, W. and Al-Ghamdi, S.G., 2024. Augmented human thermal discomfort in urban centers of the Arabian Peninsula. *Scientific Reports*, 14(1). [DOI]
- Ullah, S. and Al-Ghamdi, S.G., 2023. Advancing Urban Resilience and Sustainability Through the WRF-Urban Model: Bridging Numerical Modeling and Real-World Applications. In: *Sustainable Cities in a Changing Climate*, pp.17-27. Wiley. [DOI]
- Urban, J., Piki, M., Zemek, F. and Novotny, J., 2022. Using Google Street View photographs to assess long-term outdoor thermal perception and thermal comfort in the urban environment during heatwaves. *Frontiers in Environmental Science*, 10. [DOI]
- van Eck, N.J., 2010. Software survey: VOSviewer, a computer program for bibliometric mapping. *Scientometrics*, 84(2), pp.523-538. [DOI]
- Vulova, S., Meier, F., Rocha, A.D., Quanz Justus and Nouri, H. and Kleinschmit, B., 2021. Modeling urban evapotranspiration using remote sensing, flux footprints, and artificial intelligence. *Science of the Total Environment*, 786. [DOI]
- Vyas, A., Shastri, B. and Joshi, Y., 2014. Spatio-temporal analysis of UHI using geo-spatial techniques: A case study of Ahmedabad city. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*, 40(8), pp.997-1002. [DOI]
- Weng, Q., Lu, D. and Schubring, J., 2004. Estimation of land surface temperature-vegetation abundance relationship for urban heat island studies. *Remote Sensing of Environment*, 89(4), pp.467-483. [DOI]
- Wong, N.H., Tan, C.L., Kolokotsa, D.D. and Takebayashi, H., 2021. Greenery as a mitigation and adaptation strategy to urban heat. *Nature Reviews Earth & Environment*, 2(3), pp.166-181. [DOI]

- Yang, H.-R., Li, Y.-H., Wu, W.-J., Zhao, A.-L. and Zhang, H., 2025. Exploring the Complex Relationships Between Influential Factors of Urban Land Development Patterns and Urban Thermal Environment: A Study on Downtown Shanghai. *Sustainability*, 17(19). [DOI]
- Yang, J., Zhao, L. and Oleson, K., 2023. Large humidity effects on urban heat exposure and cooling challenges under climate change. *Environmental Research Letters*, 18(4). [DOI]
- Yang, S. and La Roche, P., 2025. Microclimate Analysis of Tree Canopies and Green Surface Combinations for Urban Heat Island Mitigation in Los Angeles and Phoenix. *Buildings*, 15(9). [DOI]
- Yang, S., Wang, L.L., Stathopoulos, T. and Marey, A.M., 2023. Urban microclimate and its impact on built environment—a review. *Building and Environment*, 238, p.110334. [DOI]
- Yang, X. and Ma, Y., 2024. Design and thermal-optical environment simulation of double-slope greenhouse roof structure based on Ecotect. *Agriculture*, 14(8), p.1410. [DOI]
- Yazdi, H., Moser-Reischl, A., Roetzer, T., Petzold, F. and Ludwig, F., 2024. Machine learning-based prediction of tree crown development in competitive urban environments. *Urban Forestry & Urban Greening*, 101. [DOI]
- Ying, J., 2022. Green infrastructure: systematic literature review. *Economic Research-Ekonomska Istrazivanja*, 35(1), pp.343–366. [DOI]
- Zakrzewska, A., Kopec, D., Ochtyra, A. and Potuckova, M., 2023. Can canopy temperature acquired from an airborne level be a tree health indicator in an urban environment? *Urban Forestry & Urban Greening*, 79. [DOI]
- Zhang, H., Cen, X., An, H. and Yin, Y., 2024. Quantitative assessment and driving factors analysis of surface urban heat island of urban agglomerations in China based on GEE. *Environmental Science and Pollution Research*, 31(34), pp.47350–47364. [DOI]
- Zhang, X., Friedl, M.A., Schaaf, C.B., Strahler, A.H., Hodges, J.C.F., Gao, F., Reed, B.C. and Huete, A., 2003. Monitoring vegetation phenology using MODIS. *Remote Sensing of Environment*, 84(3), pp.471–475. [DOI]
- Zhang, Y., Li, S., Fu, X. and Dong, R., 2021. Quantification of urban greenery using hemisphere-view panoramas with a green cover index. *Ecosystem Health And Sustainability*, 7(1). [DOI]
- Zhou, W., Huang, G. and Cadenasso, M.L., 2011. Does spatial configuration matter? Understanding the effects of land cover pattern on land surface temperature in urban landscapes. *Landscape and Urban Planning*, 102(1), pp.54–63. [DOI]